

CANDIDATE ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM

SCIENTIFIC NAME: *Korthalsella degeneri*

COMMON NAME: Hulumoa

LEAD REGION: Region 1

INFORMATION CURRENT AS OF: February 2003

STATUS/ACTION (Check all that apply):

New candidate

Continuing candidate

Non-petitioned

Petitioned - Date petition received: ____

90-day positive - FR date: ____

12-month warranted but precluded - FR date: ____

Is the petition requesting a reclassification of a listed species?

Listing priority change

Former LP: ____

New LP: ____

Latest date species first became a Candidate: ____

Candidate removal: Former LP: ____ (Check only one reason)

A - Taxon more abundant or widespread than previously believed or not subject to a degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status.

F - Range is no longer a U.S. territory.

M - Taxon mistakenly included in past notice of review.

N - Taxon may not meet the Act's definition of "species."

X - Taxon believed to be extinct.

ANIMAL/PLANT GROUP AND FAMILY: Plant, Viscaceae

HISTORICAL STATES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, island of Oahu

CURRENT STATES/COUNTIES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, island of Oahu

LEAD REGION CONTACT (Name, phone number): Scott McCarthy, 503-231-6131

LEAD FIELD OFFICE CONTACT (Office, name, phone number): Pacific Islands (Ecological Services), Christa Russell, 808-541-3441

BIOLOGICAL INFORMATION (Describe habitat, historic vs. current range, historic vs. current population estimates (# populations, #individuals/population), etc.):

This species is found only on the island of Oahu, in Makua Valley. It is parasitic on two species of native trees, *Sapindus oahuensis* and *Nestigis sandwicensis*, which are found only in mesic forests. Native mesic forest communities are increasingly rare and are under constant threat of degradation from development, alien plant species, goats, and other factors (Marie Bruegmann, Service, pers. comm., 1997). Prior to human arrival on Oahu, native mesic forests were widespread and abundant and it is assumed that this species was relatively abundant within this habitat type. Recent surveys indicate that the species is known only from one population of 1,000 individuals.

THREATS (Describe threats in terms of the five factors in section 4 of the ESA providing specific, substantive information. If this is a removal of a species from candidate status or a change in listing priority, explain reasons for change):

A. The present or threatened destruction, modification, or curtailment of its habitat or range. All mesic forests on Oahu have been degraded by goats and alien plant species, which threaten the continued existence of this species (Joel Lau, Nature Conservancy; John Obata, private consultant; Winona Char, private consultant; Bill Garnett, private consultant, pers. comms., 1996).

As early as 1778, European explorers introduced livestock, which became feral, increased in number and range, and caused significant changes to the natural environment of Hawaii. Past and present activities of introduced alien mammals are the primary factor altering and degrading vegetation and habitats on Oahu. Feral ungulates trample and eat native vegetation and disturb and open areas. This causes erosion and allows the entry of alien plant species (Cuddihy and Stone 1990; Wagner *et al.* 1990). The goat (*Capra hircus*), a species originally native to the Middle East and India, was successfully introduced to the Hawaiian Islands in 1792. Currently, populations exist on Kauai, Oahu, Maui, and Hawaii. Goats are managed in Hawaii as a game animal, but many herds populate inaccessible areas where hunting has little effect on their numbers (Hawaii Heritage Program 1990c). Goat hunting is allowed year-round or during certain months, depending on the area Hawaii Department of Land and Natural Resources n.d.-a, n.d.-b, n.d.-c, 1990). Goats browse on introduced grasses and native plants, especially in drier and more open ecosystems. Feral goats eat native vegetation, trample roots and seedlings, cause erosion, and promote the invasion of alien plants. They are able to forage in extremely rugged terrain and have a high reproductive capacity (Clarke and Cuddihy 1980; Cuddihy and Stone 1990; Culliney 1988; Scott *et al.* 1986; Tomich 1986; van Riper and van Riper 1982).

The original native flora of Hawaii consisted of about 1,000 species, 89 percent of which were endemic. Of the total native and naturalized Hawaiian flora of 1,817 species, 47 percent were introduced from other parts of the world and nearly 100 species have become pests (Smith 1985; Wagner *et al.* 1990). Naturalized, introduced species compete with native plants for space, light, water, and nutrients (Cuddihy and Stone 1990). Some of these species were brought to Hawaii by various groups of people, including the Polynesian immigrants, for food or cultural reasons. Plantation owners, alarmed at the reduction of water resources for their crops caused by the

destruction of native forest cover by grazing feral animals, supported the introduction of alien tree species for reforestation. Ranchers intentionally introduced pasture grasses and other species for agriculture, and sometimes inadvertently introduced weed seeds as well. Other plants were brought to Hawaii for their potential horticultural value (Cuddihy and Stone 1990; Scott *et al.* 1986). Many of these introduced alien plant taxa are highly invasive, out-competing and displacing native plants.

B. Overutilization for commercial, recreational, scientific, or educational purposes.

None known.

C. Disease or predation.

None known.

D. The inadequacy of existing regulatory mechanisms.

The State of Hawaii does not recognize this species as endangered until it is federally listed as endangered.

E. Other natural or manmade factors affecting its continued existence.

This species is threatened by fire (J. Lau, J. Obata, W. Char, B. Garnett, pers. comms., 1996). The U.S. Army uses Makua Valley as a firing range and, as a result, fires periodically occur, especially during the dry summer months. Because Hawaiian plants were subjected to fire during their evolution only in areas of volcanic activity and from occasional lightning strikes, they are not adapted to recurring fire regimes and do not quickly recover following a fire. Alien plants are often better adapted to fire than native plant species, and some fire-adapted grasses have become widespread in Hawaii. Native shrubland and dry forest can thus be converted to land dominated by alien grasses. The presence of such species in Hawaiian ecosystems greatly increases the intensity, extent, and frequency of fire, especially during drier months or drought. Fire-adapted alien plant taxa can reestablish in a burned area, resulting in a reduction in the amount of native vegetation after each fire. Fire can destroy dormant seeds as well as plants, even in steep or inaccessible areas. Fires may result from natural causes, or they may be accidentally or purposely started by humans (Cuddihy and Stone 1990). With only one remaining population, the threat from fire and natural events, such as hurricanes and landslides, is very high.

FOR RECYCLED PETITIONS:

- a. Is listing still warranted? ___
- b. To date, has publication of a proposal to list been precluded by other higher priority listing actions? ___
- c. Is a proposal to list the species as threatened or endangered in preparation? ___
- d. If the answer to c. above is no, provide an explanation of why the action is still precluded.

LAND OWNERSHIP (Estimate proportion Federal/state/local government/private, identify non-private owners): The population is located on public land in Makua Valley, in an area

operated as a firing range by the U.S. Army.

PRELISTING (Describe status of conservation agreements or other conservation activities):
The U.S. Army has initiated efforts to exclude goats from portions of Makua Valley through fencing. However, the completed fences will not create complete enclosure areas, they will simply make it more difficult for the goats to get to the areas with rare plant species.

REFERENCES (Identify primary sources of information (e.g., status reports, petitions, journal publications, unpublished data from species experts) using formal citation format):

The information in this form is based on the results of a meeting of 20 botanical experts held by the Center for Plant Conservation in December of 1995, and has been updated by personal communication with Joel Lau of The Nature Conservancy and Winona Char, private consultant. Based on the IUCN Red Plant Data Book rarity categories, this species is recognized as Rare (could be considered at risk) by Wagner, Herbst, and Sohmer (1990). Additional sources of information include:

Clarke, G., and L.W. Cuddihy. 1980. A botanical reconnaissance of the Na Pali coast trail: Kee Beach to Kalalau Valley (April 9-11, 1980). Division of Forestry and Wildlife, Department of Land and Natural Resources, Hilo, Hawaii.

Corn, C.A., G. Clarke, L. Cuddihy, and L. Yoshida. 1979. A botanical reconnaissance of Kalalau, Honopu, Awaawapuhi, Nualolo and Milolii Valleys and shorelines--Na Pali, Kauai. Unpublished report. Division of Forestry and Wildlife, Department of Land and Natural Resources, Endangered Species Program, Honolulu. 14 pp.

Cuddihy, L.W., and C.P. Stone. 1990. Alteration of native Hawaiian vegetation; effects of humans, their activities and introductions. Coop. Natl. Park Resources Stud. Unit, Hawaii. 138 pp.

Culliney, J.L. 1988. Islands in a far sea; nature and man in Hawaii. Sierra Club Books, San Francisco. 410 pp.

Scott, J.M., S. Mountainspring, F.L. Ramsey, and C.B. Kepler. 1986. Forest bird communities of the Hawaiian Islands: Their dynamics, ecology, and conservation. Studies in Avian Biology 9:1-429. Cooper Ornithological Society, Los Angeles.

Smith, C.W. 1985. Impact of alien plants on Hawai'i's native biota: in Stone, C.P., and J.M. Scott (eds.), Hawai'i's terrestrial ecosystems: preservation and management. Coop. Natl. Park Resources Stud. Unit, Univ. Hawaii, Honolulu, pp. 180-250.

Wagner, W.L., D.R. Herbst, and S.H. Sohmer. 1990. Manual of the flowering plants of Hawai'i. University of Hawaii Press and Bishop Museum Press, Honolulu. Bishop Mus. Spec. Publ. 83:1-1853.

LISTING PRIORITY (* after number)

THREAT

Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2
		Subspecies/population	3
	Non-imminent	Monotypic genus	4
		Species	5 *
		Subspecies/population	6
Moderate to Low	Imminent	Monotypic genus	7
		Species	8
		Subspecies/population	9
	Non-imminent	Monotypic genus	10
		Species	11
		Subspecies/population	12

Rationale for listing priority number:

Magnitude:

Imminence:

APPROVAL/CONCURRENCE: Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes to the candidate list, including listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all additions of species to the candidate list, removal of candidates, and listing priority changes.

Approve: Rowan Gould March 6, 2003
Regional Director, Fish and Wildlife Service Date

Concur: _____
Director, Fish and Wildlife Service Date

Do not concur: _____
Director, Fish and Wildlife Service Date

Director's Remarks:

-

-

Date of annual review: 2/03

Conducted by: _____

Comments:

-

-