

CANDIDATE ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM

SCIENTIFIC NAME: *Ochrosia haleakalae*

COMMON NAME: Holei

LEAD REGION: Region 1

INFORMATION CURRENT AS OF: February 2003

STATUS/ACTION (Check all that apply):

New candidate

Continuing candidate

Non-petitioned

Petitioned - Date petition received: ____

90-day positive - FR date: ____

12-month warranted but precluded - FR date: ____

Is the petition requesting a reclassification of a listed species?

Listing priority change

Former LP: ____

New LP: ____

Latest date species first became a Candidate: ____

Candidate removal: Former LP: ____ (Check only one reason)

A - Taxon more abundant or widespread than previously believed or not subject to a degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status.

F - Range is no longer a U.S. territory.

M - Taxon mistakenly included in past notice of review.

N - Taxon may not meet the Act's definition of "species."

X - Taxon believed to be extinct.

ANIMAL/PLANT GROUP AND FAMILY: Plant, Apocynaceae (Dogbane family)

HISTORICAL STATES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, islands of Maui and Hawaii

CURRENT STATES/COUNTIES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, islands of Maui and Hawaii

LEAD REGION CONTACT (Name, phone number): Scott McCarthy, 503-231-6131

LEAD FIELD OFFICE CONTACT (Office, name, phone number): Pacific Islands (Ecological Services), Christa Russell, 808-541-3441

BIOLOGICAL INFORMATION (Describe habitat, historic vs. current range, historic vs. current population estimates (# populations, #individuals/population), etc.):

This species is currently known from one population totaling 150-200 individuals on East Maui and 1 population of a few individuals on Hawaii (Perlman and Wood 1996; Art Medeiros, USGS Biological Resources Division, pers. comm., 1995, 1999). Typical habitat is dry to mesic forest, often on lava (Wagner *et al.* 1990). On Maui, the species occurs in diverse dry forest in the Auwahi area (Medeiros *et al.* 1986; A. Medeiros, pers. comm., 1995). *Ochrosia halaekalae* was only recently discovered on the island of Hawaii, in degraded *Metrosideros polymorpha/Pisonia sandwicensis* (‘ohi‘a/papala kepau) mesic forest in the Kohala Mountains (Perlman and Wood 1996).

THREATS (Describe threats in terms of the five factors in section 4 of the ESA providing specific, substantive information. If this is a removal of a species from candidate status or a change in listing priority, explain reasons for change):

A. The present or threatened destruction, modification, or curtailment of its habitat or range. *Ochrosia halaekalae* is highly and imminently threatened by cattle and feral goats on Maui (Medeiros *et al.* 1986; A. Medeiros, pers. comm., 1995). On the island of Hawaii, pigs are the major threat (Perlman and Wood 1996). As early as 1778, European explorers introduced livestock, which became feral, increased in number and range, and caused significant changes to the natural environment of Hawaii. Past and present activities of introduced alien mammals are the primary factors in altering and degrading vegetation and habitats on the Hawaiian islands. Feral ungulates trample and eat native vegetation and disturb and open areas. This causes erosion and allows the entry of alien plant taxa (Cuddihy and Stone 1990; Wagner *et al.* 1990).

Cattle (*Bos taurus*), the wild progenitor of which was native to Europe, northern Africa, and southwestern Asia, were introduced to the Hawaiian Islands in 1793. Large feral herds developed as a result of restrictions on killing cattle decreed by King Kamehameha I. While small cattle ranches were developed on Kauai, Oahu, and West Maui, very large ranches of tens of thousands of acres were created on East Maui and Hawaii. Much of the land used in these private enterprises was leased from the State or was privately owned and considered Forest Reserve and/or Conservation District land. Feral cattle can presently be found on the island of Hawaii, and ranching is still a major commercial activity there. Hunting of feral cattle is no longer allowed in Hawaii (Hawaii Department of Land and Natural Resources 1985). Cattle eat native vegetation, trample roots and seedlings, cause erosion, create disturbed areas into which alien plants invade, and spread seeds of alien plants in their feces and on their bodies. The forest in areas grazed by cattle becomes degraded to grassland pasture, and plant cover is reduced for many years following removal of cattle from an area. Several alien grasses and legumes purposely introduced for cattle forage have become noxious weeds (Cuddihy and Stone 1990; Tomich 1986).

The goat (*Capra hircus*), a species originally native to the Middle East and India, was successfully introduced to the Hawaiian Islands in 1792. Currently, populations exist on Kauai, Oahu, Maui, and Hawaii. On Maui, feral goats have been present in drier, more rugged areas since the 1800s. Goats are managed in Hawaii as a game animal, but many herds populate inaccessible areas where hunting has little effect on their numbers (Hawaii Heritage Program 1990c). Goat hunting is allowed year-round or during certain months, depending on the area

(Hawaii Department of Land and Natural Resources(DLNR) n.d.-a, n.d.-b, n.d.-c, 1990). Goats browse on introduced grasses and native plants, especially in drier and more open ecosystems. Feral goats eat native vegetation, trample roots and seedlings, cause erosion, and promote the invasion of alien plants. They are able to forage in extremely rugged terrain and have a high reproductive capacity (Clarke and Cuddihy 1980; Cuddihy and Stone 1990; Culliney 1988; Scott *et al.* 1986; Tomich 1986; van Riper and van Riper 1982).

Although many plant species survive on steep cliffs inaccessible to goats, the original range of these plants was probably much larger. These species are now vulnerable to the long-term, indirect effects of goats, such as large-scale erosion (Corn *et al.* 1979). The habitat on leeward east Maui has been damaged in the past by goats, and these effects are still apparent in the form of alien vegetation and erosion (Clarke and Cuddihy 1980; Culliney 1988; Scott *et al.* 1986; van Riper and van Riper 1982).

The pig (*Sus scrofa*) is originally native to Europe, northern Africa, Asia Minor, and Asia. European pigs, introduced to Hawaii by Captain James Cook in 1778, became feral and invaded forested areas, especially wet and mesic forests and dry areas at high elevations. They are currently present on Kauai and four other islands, and inhabit rain forests and grasslands. Pig hunting is allowed on all islands either year-round or during certain months, depending on the area (DLNR n.d.-a, n.d.-b, n.d.-c, 1990). While rooting in the ground in search of the invertebrates and plant material they eat, feral pigs disturb and destroy vegetative cover, trample plants and seedlings, and threaten forest regeneration by damaging seeds and seedlings. They disturb soil and cause erosion, especially on slopes. Alien plant seeds are dispersed on their hooves and coats as well as through their digestive tracts, and the disturbed soil is fertilized by their feces, helping these plants to establish. Pigs are a major vector in the spread of many introduced plant species (Cuddihy and Stone 1990; Medeiros *et al.* 1986; Scott *et al.* 1986; Smith 1985; Stone 1985; Tomich 1986; Wagner *et al.* 1990).

B. Overutilization for commercial, recreational, scientific, or educational purposes.

None known.

C. Disease or predation.

None known.

D. The inadequacy of existing regulatory mechanisms.

There is no Federal or State protection for this taxon. The State of Hawaii does not recognize this species as endangered until it is federally listed as endangered.

E. Other natural or manmade factors affecting its continued existence.

Ochrosia haleakalae is threatened by fire and alien plant species (Medeiros *et al.* 1986; Perlman and Wood 1996; A. Medeiros, pers. comm., 1995). Because Hawaiian plants were subjected to fire during their evolution only in areas of volcanic activity and from occasional lightning

strikes, they are not adapted to recurring fire regimes and do not quickly recover following a fire. Alien plants are often better adapted to fire than native plant species, and some fire-adapted grasses have become widespread in Hawaii. Native shrubland and dry forest can thus be converted to land dominated by alien grasses. The presence of such species in Hawaiian ecosystems greatly increases the intensity, extent, and frequency of fire, especially during drier months or drought. Fire-adapted alien plant taxa can reestablish in a burned area, resulting in a reduction in the amount of native vegetation after each fire. Fire can destroy dormant seeds as well as plants, even in steep or inaccessible areas. Fires may result from natural causes, or they may be accidentally or purposely started by humans (Cuddihy and Stone 1990).

FOR RECYCLED PETITIONS:

- a. Is listing still warranted? ___
- b. To date, has publication of a proposal to list been precluded by other higher priority listing actions? ___
- c. Is a proposal to list the species as threatened or endangered in preparation? ___
- d. If the answer to c. above is no, provide an explanation of why the action is still precluded.

LAND OWNERSHIP (Estimate proportion Federal/state/local government/private, identify non-private owners): This species is found on private land and State-owned land on Maui, and State-owned land on Hawaii.

PRELISTING (Describe status of conservation agreements or other conservation activities): The Service has funded the USGS Biological Resources Division to fence the Maui population of *Ochrosia haleakalae* on private land.

REFERENCES (Identify primary sources of information (e.g., status reports, petitions, journal publications, unpublished data from species experts) using formal citation format):

The information in this form is based on the results of a meeting of 20 botanical experts held by the Center for Plant Conservation in December 1995 and November 1996, and has been updated with information from a Service funded survey by National Tropical Botanical Garden in late 1995 and from Art Medeiros of USGS Biological Resources Division in 1999. Based on the IUCN Red Plant Data Book rarity categories, this species is recognized as Rare (could be considered at risk) by Wagner, Herbst, and Sohmer in the 1990 Manual of Flowering Plants of Hawai'i.

Cuddihy, L.W., and C.P. Stone. 1990. Alteration of native Hawaiian vegetation; effects of humans, their activities and introductions. Coop. Natl. Park Resources Stud. Unit, Hawaii. 138 pp.

Culliney, J.L. 1988. Islands in a far sea; nature and man in Hawaii. Sierra Club Books, San Francisco. 410 pp.

Hawaii, Department of Land and Natural Resources. N.d.-a. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Oahu. Division of Forestry and Wildlife, Honolulu. 2 pp.

- Hawaii, Department of Land and Natural Resources. N.d.-b. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Molokai. Division of Forestry and Wildlife, Honolulu. 2 pp.
- Hawaii, Department of Land and Natural Resources. N.d.-c. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Maui. Division of Forestry and Wildlife, Honolulu. 2 pp.
- Hawaii, Department of Land and Natural Resources. 1985. Hunting in Hawaii, fourth revision. Division of Forestry and Wildlife, Honolulu, 32 pp.
- Medeiros, A.C., Jr., L.L. Loope, and R.A. Holt. 1986. Status of native flowering plant species on the south slope of Haleakala, East Maui, Hawaii. Coop. Natl. Park Resources Stud. Unit, Hawaii, Techn. Rept. 59:1-230.
- Perlman, S. And K. Wood. 1996. Kohala Mountains Survey. Prepared for the U.S. Fish and Wildlife Service by National Tropical Botanical Garden, January 1996.
- Scott, J.M., S. Mountainspring, F.L. Ramsey, and C.B. Kepler. 1986. Forest bird communities of the Hawaiian Islands: Their dynamics, ecology, and conservation. Studies in Avian Biology 9:1-429. Cooper Ornithological Society, Los Angeles.
- Stone, C.P. 1985. Alien animals in Hawai'i's native ecosystems: toward controlling the adverse effects of introduced vertebrates: in Stone, C.P., and J.M. Scott (eds.), Hawai'i's terrestrial ecosystems: preservation and management. Coop. Natl. Park Resources Stud. Unit, Univ. Hawaii, Honolulu, pp. 251-197.
- Tomich, P.Q. 1986. Mammals in Hawai'i; a synopsis and notational bibliography. Bishop Museum Press, Honolulu. 375 pp.
- van Riper, S.G., and C. van Riper III. 1982. A field guide to the mammals in Hawaii. The Oriental Publishing Company, Honolulu. 68 pp.
- Wagner, W.L., D.R. Herbst, and S.H. Sohmer. 1990. Manual of the flowering plants of Hawai'i. University of Hawaii Press and Bishop Museum Press, Honolulu. Bishop Mus. Spec. Publ. 83:1-1853.
- Wenkam, R. 1969. Kauai and the park country of Hawaii. Sierra Club, San Francisco. 160 pp.

LISTING PRIORITY (* after number)

THREAT

Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2 *
		Subspecies/population	3
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/population	6
Moderate to Low	Imminent	Monotypic genus	7
		Species	8
		Subspecies/population	9
	Non-imminent	Monotypic genus	10
		Species	11
		Subspecies/population	12

Rationale for listing priority number:

Magnitude:

Imminence:

APPROVAL/CONCURRENCE: Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes to the candidate list, including listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all additions of species to the candidate list, removal of candidates, and listing priority changes.

Approve: Rowan Gould March 6, 2003
Regional Director, Fish and Wildlife Service Date

Concur: _____
Director, Fish and Wildlife Service Date

Do not concur: _____
Director, Fish and Wildlife Service Date

Director's Remarks:

-

-

Date of annual review: 2/03

Conducted by: _____

Comments:

-

-