

CANDIDATE ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM

SCIENTIFIC NAME: *Stenogyne cranwelliae*

COMMON NAME: No common name

LEAD REGION: Region 1

INFORMATION CURRENT AS OF: February 2003

STATUS/ACTION (Check all that apply):

New candidate

Continuing candidate

Non-petitioned

Petitioned - Date petition received: ____

90-day positive - FR date: ____

12-month warranted but precluded - FR date: ____

Is the petition requesting a reclassification of a listed species?

Listing priority change

Former LP: ____

New LP: ____

Latest date species became a Candidate: ____

Candidate removal: Former LP: ____ (Check only one reason)

A - Taxon more abundant or widespread than previously believed or not subject to a degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status.

F - Range is no longer a U.S. territory.

M - Taxon mistakenly included in past notice of review.

N - Taxon may not meet the Act's definition of "species."

X - Taxon believed to be extinct.

ANIMAL/PLANT GROUP AND FAMILY: Plant, Lamiaceae (mint family)

HISTORICAL STATES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, island of Hawaii

CURRENT STATES/COUNTIES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, island of Hawaii

LEAD REGION CONTACT (Name, phone number): Scott McCarthy, 503-231-6131

LEAD FIELD OFFICE CONTACT (Office, name, phone number): Pacific Islands (Ecological Services), Christa Russell, 808-541-3441

BIOLOGICAL INFORMATION (Describe habitat, historic vs. current range, historic vs. current population estimates (# populations, #individuals/population), etc.):

Stenogyne cranwelliae is known from six populations of 100 individuals. Historically found in the Kohala Mts. on the island of Hawaii, this species was thought to be extinct until rediscovered during surveys of the Kohala Mts in 1995. Typical habitat is wet forest dominated by *Metrosideros polymorpha* (ōhi`a) (Perlman and Wood 1996; Weller and Sakai 1990).

THREATS (Describe threats in terms of the five factors in section 4 of the ESA providing specific, substantive information. If this is a removal of a species from candidate status or a change in listing priority, explain reasons for change):

A. The present or threatened destruction, modification, or curtailment of its habitat or range. This species is highly and imminently threatened by feral pigs (Perlman and Wood 1996). As early as 1778, European explorers introduced livestock, which became feral, increased in number and range, and caused significant changes to the natural environment of Hawaii. Past and present activities of introduced alien mammals are the primary factor altering and degrading vegetation and habitats on Hawaii. Feral ungulates trample and eat native vegetation and disturb and open areas. This causes erosion and allows the entry of alien plant species (Cuddihy and Stone 1990; Wagner *et al.* 1990).

The pig (*Sus scrofa*) is originally native to Europe, northern Africa, Asia Minor, and Asia. European pigs, introduced to Hawaii by Captain James Cook in 1778, became feral and invaded forested areas, especially wet and mesic forests and dry areas at high elevations. They are currently present on Hawaii and four other islands, and inhabit rain forests and grasslands. Pig hunting is allowed on all islands either year-round or during certain months, depending on the area (Hawaii Department of Land and Natural Resources n.d.-a, n.d.-b, n.d.-c, 1990). While rooting in the ground in search of the invertebrates and plant material they eat, feral pigs disturb and destroy vegetative cover, trample plants and seedlings, and threaten forest regeneration by damaging seeds and seedlings. They disturb soil and cause erosion, especially on slopes. Alien plant seeds are dispersed on their hooves and coats as well as through their digestive tracts, and the disturbed soil is fertilized by their feces, helping these plants to establish. Pigs are a major vector in the spread of many introduced plant species (Cuddihy and Stone 1990; Medeiros *et al.* 1986; Scott *et al.* 1986; Smith 1985; Stone 1985; Tomich 1986; Wagner *et al.* 1990).

B. Overutilization for commercial, recreational, scientific, or educational purposes.

None known.

C. Disease or predation.

Rats are a potential threat to *Stenogyne cranwelliae* (Perlman and Wood 1996). Of the four species of rodents that have been introduced to the Hawaiian Islands, the species with the greatest impact on the native flora and fauna is probably *Rattus rattus* (black or roof rat), that now occurs on all the main Hawaiian Islands around human habitations, cultivated fields, and forests. Black rats, and to a lesser extent *Mus musculus* (house mouse), *Rattus exulans* (Polynesian rat), and *R. norvegicus* (Norway rat), eat the fruits of some native plants, especially those with large, fleshy fruits. Many native Hawaiian plants produce fruit over an extended period of time, thus producing a prolonged food supply for rodent populations. Black rats strip bark from some native plants, and eat the fleshy stems and fruits of plants (Cuddihy and Stone 1990; Tomich 1986).

D. The inadequacy of existing regulatory mechanisms.

There is no Federal or State protection for this taxon. The State of Hawaii does not recognize this species as endangered until it is federally listed as endangered.

E. Other natural or manmade factors affecting its continued existence.

This species is threatened by several alien plant species. With only 100 individuals in six populations of restricted distribution, reduced reproductive vigor and extinction due to naturally occurring events, such as hurricanes or landslides, are also major threats (Perlman and Wood 1996). The original native flora of Hawaii consisted of about 1,000 species, 89 percent of which were endemic. Of the total native and naturalized Hawaiian flora of 1,817 taxa, 47 percent were introduced from other parts of the world and nearly 100 species have become pests (Smith 1985; Wagner *et al.* 1990). Naturalized, introduced species compete with native plants for space, light, water, and nutrients (Cuddihy and Stone 1990). Some of these species were brought to Hawaii by various groups of people, including the Polynesian immigrants, for food or cultural reasons. Plantation owners, alarmed at the reduction of water resources for their crops caused by the destruction of native forest cover by grazing feral animals, supported the introduction of alien tree species for reforestation. Ranchers intentionally introduced pasture grasses and other species for agriculture, and sometimes inadvertently introduced weed seeds as well. Other plants were brought to Hawaii for their potential horticultural value (Cuddihy and Stone 1990; Scott *et al.* 1986; Wenkam 1969).

Introduced species that threaten *Stenogyne cranwelliae* include *Setaria palmifolia* (palm grass), *Hedychium* spp. (gingers), and *Tibouchina herbacea* (glorybush) (Perlman and Wood 1996). *Setaria palmifolia* (palmgrass), native to tropical Asia, has become naturalized in mesic valleys, wet forests, and along streams on Oahu, Lanai, Maui, and Hawaii. First collected in 1903, major infestations can now be found in the O'laa area and the windward side of the island of Hawaii (Cuddihy and Stone 1990; O'Connor 1990).

Hedychium coronarium (white ginger) was introduced to Hawaii in the late 1800s, probably by Chinese immigrants. It escaped from cultivation and is found in wet and mesic forests on most of the main Hawaiian islands. The large, vigorous herbs mainly reproduce vegetatively, forming very dense stands that exclude all other growth. *Hedychium gardnerianum* (kahili ginger) was introduced to Hawaii before 1940 from the Himalayas, and now has major infestations on the islands of Hawaii, Maui, and Kauai. This species is considered a more serious threat to native forests because it produces abundant fruit (Cuddihy and Stone 1990; Wagner *et al.* 1990). *Tibouchina herbacea* (glorybush) first became established on the island of Hawaii in the late 1970s and, by 1982, was collected in Lanilili on West Maui (Almeda 1990). Although the disruptive potential of this alien plant is not fully known, *Tibouchina herbacea* appears to be invading mesic and wet forests of Hawaii and Maui (Cuddihy and Stone 1990).

FOR RECYCLED PETITIONS:

- a. Is listing still warranted? ____
- b. To date, has publication of a proposal to list been precluded by other higher priority listing actions? ____
- c. Is a proposal to list the species as threatened or endangered in preparation? ____

d. If the answer to c. above is no, provide an explanation of why the action is still precluded.

LAND OWNERSHIP (Estimate proportion Federal/state/local government/private, identify non-private owners): All six populations occur on State land.

PRELISTING (Describe status of conservation agreements or other conservation activities): None.

REFERENCES (Identify primary sources of information (e.g., status reports, petitions, journal publications, unpublished data from species experts) using formal citation format):

The information in this form is based on the results of a meeting of 20 botanical experts held by the Center for Plant Conservation in December of 1995 and November of 1996, and has been updated by personal communications as cited in the text.

Almeda, F. 1990. Melastomataceae: in Wagner, W.L., D.R. Herbst, and S.H. Sohmer, Manual of the flowering plants of Hawai'i. University of Hawaii Press and Bishop Museum Press, Honolulu. Bishop Mus. Spec. Publ. 83:903-917.

Cuddihy, L.W., and C.P. Stone. 1990. Alteration of native Hawaiian vegetation; effects of humans, their activities and introductions. Coop. Natl. Park Resources Stud. Unit, Hawaii. 138 pp.

Hawaii, Department of Land and Natural Resources. N.d.-a. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Oahu. Division of Forestry and Wildlife, Honolulu. 2 pp.

Hawaii, Department of Land and Natural Resources. N.d.-b. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Molokai. Division of Forestry and Wildlife, Honolulu. 2 pp.

Hawaii, Department of Land and Natural Resources. N.d.-c. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Maui. Division of Forestry and Wildlife, Honolulu. 2 pp.

Medeiros, A.C., Jr., L.L. Loope, and R.A. Holt. 1986. Status of native flowering plant species on the south slope of Haleakala, East Maui, Hawaii. Coop. Natl. Park Resources Stud. Unit, Hawaii, Techn. Rept. 59:1-230.

O'Connor, P.J. 1990. Poaceae: in Wagner, W.L., D.R. Herbst, and S.H. Sohmer, Manual of the flowering plants of Hawai'i. University of Hawaii Press and Bishop Museum Press, Honolulu. Bishop Mus. Spec. Publ. 83:1481-1604.

Perlman, S. and K. Wood. 1996. Kohala Mountains surveys. Prepared by National Tropical Botanical Garden for U.S. Fish and Wildlife Service, Honolulu.

Scott, J.M., S. Mountainspring, F.L. Ramsey, and C.B. Kepler. 1986. Forest bird communities of the Hawaiian Islands: Their dynamics, ecology, and conservation. Studies in Avian

- Biology 9:1-429. Cooper Ornithological Society, Los Angeles.
- Smith, C.W. 1985. Impact of alien plants on Hawai'i's native biota: in Stone, C.P., and J.M. Scott (eds.), Hawai'i's terrestrial ecosystems: preservation and management. Coop. Natl. Park Resources Stud. Unit, Univ. Hawaii, Honolulu, pp. 180-250.
- Stone, C.P. 1985. Alien animals in Hawai'i's native ecosystems: toward controlling the adverse effects of introduced vertebrates: in Stone, C.P., and J.M. Scott (eds.), Hawai'i's terrestrial ecosystems: preservation and management. Coop. Natl. Park Resources Stud. Unit, Univ. Hawaii, Honolulu, pp. 251-197.
- Tomich, P.Q. 1986. Mammals in Hawai'i; a synopsis and notational bibliography. Bishop Museum Press, Honolulu. 375 pp.
- Wagner, W.L., D.R. Herbst, and S.H. Sohmer. 1990. Manual of the flowering plants of Hawai'i. University of Hawaii Press and Bishop Museum Press, Honolulu. Bishop Mus. Spec. Publ. 83:1-1853.
- Weller, S.G. and A.K. Sakai. 1990. *Stenogyne*: *in* Wagner, W.L., D.R. Herbst, and S.H. Sohmer, Manual of the flowering plants of Hawaii. Univ. of Hawaii Press and Bishop Museum Press; Honolulu. Bishop Mus. Spec. Pub. 83:831-843.
- Wenkam, R. 1969. Kauai and the park country of Hawaii. Sierra Club, San Francisco. 160 pp.

LISTING PRIORITY (* after number)

THREAT

Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2 *
		Subspecies/population	3
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/population	6
Moderate to Low	Imminent	Monotypic genus	7
		Species	8
		Subspecies/population	9
	Non-imminent	Monotypic genus	10
		Species	11
		Subspecies/population	12

Rationale for listing priority number:

Magnitude:

Imminence:

APPROVAL/CONCURRENCE: Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes to the candidate list, including listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all additions of species to the candidate list, removal of candidates, and listing priority changes.

Approve: Rowan Gould March 6, 2003
Acting Regional Director, Fish and Wildlife Service Date

Concur: _____
Director, Fish and Wildlife Service Date

Do not concur: _____
Director, Fish and Wildlife Service Date

Director's Remarks:

-

-

Date of annual review: 2/03
Conducted by: _____

Comments:

-

-