

5-YEAR REVIEW

Short Form Summary

Species Reviewed: Poouli (*Melamprosops phaeosoma*)

Current Classification: Endangered

Federal Register Notice announcing initiation of this review:

[USFWS] U.S. Fish and Wildlife Service. 2012. Endangered and threatened wildlife and plants; 5-year status reviews of 46 species in Idaho, Oregon, Washington, Nevada, Montana, Hawaii, Guam, and the Northern Mariana Islands. Federal Register 77(44):13248-13251.

Lead Region/Field Office:

Region 1/Pacific Islands Fish and Wildlife Office (PIFWO), Honolulu, Hawaii

Name of Reviewer(s):

Jay Nelson, Fish and Wildlife Biologist, PIFWO

Michelle Bogardus, Maui Nui and Hawaii Island Team Manager, PIFWO

Marie Bruegmann, Plant Recovery Coordinator, PIFWO

Kristi Young, Programmatic Deputy Field Supervisor, PIFWO

Methodology used to complete this 5-year review:

This review was conducted by staff of the Pacific Islands Fish and Wildlife Office of the U.S. Fish and Wildlife Service (USFWS), beginning on May 5, 2015. The review was based on a review of current, available information since the last 5-year review for poouli (USFWS 2010). The evaluation by Jay Nelson, Fish and Wildlife Biologist, was reviewed by the Island Team Manager, followed by the Plant Recovery Coordinator. It was subsequently reviewed and approved by the Programmatic Deputy Field Supervisor.

Background:

For information regarding the species listing history and other facts, please refer to the Fish and Wildlife Service's Environmental Conservation On-line System (ECOS) database for threatened and endangered species at: http://ecos.fws.gov/tess_public.

Review Analysis:

Please refer to the previous 5-year review for poouli (*Melamprosops phaeosoma*) published on August 27, 2010 (available at: http://ecos.fws.gov/docs/five_year_review/doc3861.pdf) for a complete review of the species' status, threats, and management efforts. No significant new information regarding the species' biological status has come to light since listing to warrant a change in the Federal listing status of poouli.

The poouli is a medium sized, stocky Hawaiian honeycreeper recognized by its brown plumage and characteristic black mask framed by a gray crown and white cheek patch (USFWS 2006).

New status information:

- Surveys for Hawaiian forest birds using the variable circular-plot method as previously conducted by Scott *et al.* (1986) were conducted in forest areas on east and west Maui from 2010-2012 in areas with historical occurrence of poouli (R. Camp, U.S. Geological Survey, pers. comm. 2015). Poouli were not detected during these surveys.
- Elphick *et al.* (2010) estimated the extinction of the poouli to have occurred in 2005 using a method by which the predicted probability of extinction increases as a function of the time since a species was last observed. Using 2004 as the last reliable observation record for poouli, the authors determined the year 2008 as the upper 95% confidence bound for species extinction. This approach for establishing extinction probability however is problematic when applied to extremely rare species such as poouli that are potentially distributed over a large area because the absence of observation records may be the result of inadequate survey effort and the few if any visits by qualified observers to remote areas where rare and potentially extinct species may still exist.

New threats:

- Climate change destruction or degradation of habitat – Hawaiian honeycreepers are known to be highly susceptible to introduced avian disease, particularly avian malaria (*Plasmodium relictum*) (Atkinson *et al.* 1995; Atkinson *et al.* 2000; Yorinks and Atkinson 2000; Banko and Banko 2009). According to some climate change projections, temperature increases could present an additional threat specific to Hawaiian forest birds by causing an increase in the elevation below which regular transmission of avian malaria occurs, potentially reducing the remaining suitable habitat for these species. In Hawaii, the threshold temperature for transmission of avian malaria has been estimated to be 13 degrees Celsius (55 degrees Fahrenheit), whereas peak *P. relictum* prevalence in wild mosquitoes occurs in mid-elevation forest where the mean ambient summer temperature is 17 degrees Celsius (64 degrees Fahrenheit) (Benning *et al.* 2002). Benning *et al.* (2002) used GIS simulation to show that an increase in temperature of 2 degrees Celsius (3.6 degrees Fahrenheit), which is within the range predicted by some climate models (e.g., IPCC 2013; ICAP 2010), would result in 100 years in a 50 percent decrease in the land area for poouli where malaria transmission currently is only periodic. Lia *et al.* (2015) assessed how global climate change will affect future malaria risk for native Hawaiian bird populations and expect high elevation areas to remain mosquito free only to mid-century due to combined factors of increased rainfall and increasing temperatures. If climate change were to reduce the remaining suitable habitat for poouli as predicted, it would likely contribute to the extinction of this species over time.

New management actions:

- Surveys / inventories – Forest bird surveys were conducted on Maui in 2010-2012, but no birds were detected.

Synthesis:

We applied a standard of demonstrated species absence at the 95% confidence level to recommend species status change from unknown to “delisting” due to extinction. This standard means we assumed a species is potentially extant until survey or other information shows the species is absent with 95% confidence from suitable habitat and areas it was known to have occupied historically during the roughly last half century. The poouli was last seen in the wild in 2004 (USFWS 2006). We believe the status of the poouli is “unknown,” based on low survey effort for poouli subsequent to its last sighting. There are instances where rare Hawaiian birds have been rediscovered after they were presumed extinct or have been found in larger populations than expected (Reynolds and Snetsinger 2001). The large area on East Maui with suitable habitat (approximately 11,715 hectares [28,948 acres]) (USFWS 2012), and the many remote areas within this that are only rarely visited by qualified observers, increase the potential that a small population of poouli could still exist on Maui. The rough terrain on Maui and frequent wet weather make surveys difficult, and numerous steep valleys create many small pockets of habitat where poouli could still exist. In addition, all threats are not being sufficiently managed throughout all of the populations (Table 2).

Recommendations for Future Actions:

- Surveys / inventories
 - One of the most important recovery actions for the poouli is to intensively and systematically search areas of forest habitat where the species occurred historically (USFWS 2006). Statewide surveys of Hawaiian forest bird populations are conducted along widely spaced transects (Scott *et al.* 1986) and do not cover all areas where extremely rare Hawaiian forest birds are most likely to be. Additionally, these surveys do not spend the lengths of time needed to maximize the probability that extremely rare Hawaiian forest birds will be detected or rediscovered. Therefore, we recommend that an intensive search for poouli be conducted on Maui using similar methodologies as those employed during the Hawaii Rare Bird Search 1994-1996 (Reynolds and Snetsinger 2001).
 - In addition, we recommend that autonomous recording units, or ARUs (Fitzpatrick 2002; Wallace 2010), be deployed in suitable habitats for this species. These field recording units record vocalizations of forest birds. The tapes are then analyzed using computer programs to determine if the target species is present in the area. Use of this technology would greatly increase the amount of search time for this species.
- Threats – disease control research – Of particular concern to the continued survival of many Hawaiian forest birds (particularly Hawaiian honeycreepers) is avian disease. Existing tools and approaches have proved largely ineffective in addressing this problem given mosquito dispersal distance and the abundance of mosquito breeding sites in most wet native forest habitats (LaPointe *et al.* 2009). Opportunities are emerging however based on new genetic tools as part of the fields of synthetic biology and genomic technology that have the potential to assist Hawaiian forest birds in developing genetic resistance to avian disease (LaPointe *et al.* 2009). In addition, recent progress has been made with the development of genetically modified mosquitoes for disease control. Several of

these techniques have achieved proof-of-principle in laboratory studies, while other transgenic insect techniques, including self-sustaining technologies to achieve long-term transmission control are anticipated to advance to field testing in the near future. We encourage continued research in the fields of genomic technologies and genetically modified mosquitoes for disease control and their field application as a conservation strategy for Hawaiian forest birds.

- Habitat and natural process management and restoration –
 - Poouli would benefit in the long-term from habitat restoration to assist other endangered birds on Maui. We therefore recommend continued habitat management in areas where the species may still exist (USFWS 2006).
 - Hawaiian forest birds susceptible to avian disease may become extinct following a drastic reduction in disease free habitat, but ultimately forest might expand into higher elevations maintaining disease free refugia for some species. Acquisition and management of transmission-free high-elevation habitat is crucial to the preservation and restoration of native Hawaiian forest birds (Lapointe *et al.* 2009). As a long-term contingency against a warming scenario, we recommend securing deforested and pasture lands on Maui at high elevations adjacent to protected refugia and managing these areas for forest growth to provide suitable habitat for poouli and other Hawaiian forest birds.
- Captive propagation for reintroduction and genetic storage – Should poouli be rediscovered we recommend the Rare Bird Discovery Protocol in the revised recovery plan for Hawaiian forest birds (USFWS 2006) be followed with regard to decisions for whether to attempt to establish a captive population for propagation and reintroduction or to manage rediscovered population(s) in situ.

Table 1. Trends in status of poouli since listing.

Date	Number wild individuals	Number released	Key Recovery Actions	Actions Accomplished
1975 (listing)	Very rare	0	See below	
1980 (Hawaii forest bird survey)	140 ± 280 (95% CI)	0	See below	
1984 (first recovery plan)	141	0	Improve habitat conditions; decrease threat of avian disease; systematically search area of suitable forest habitat;	Improve habitat conditions – Yes Decrease threat of avian disease – No Systematically search area of suitable forest habitat – Yes
1994 (Hawaii Rare Bird Search)	5-6	0	See above	See above

Date	Number wild individuals	Number released	Key Recovery Actions	Actions Accomplished
2002	3	0	See above	See above
2006 (revised recovery plan)	Unknown	0	Improve habitat conditions; decrease threat of avian disease; systematically search area of suitable forest habitat;	Improve habitat conditions – Yes Decrease threat of avian disease – No Systematically search area of suitable forest habitat – No
2010 (5-year review)	Unknown	0	Improve habitat conditions; decrease threat of avian disease; systematically search area of suitable forest habitat;	Improve habitat conditions – Yes Decrease threat of avian disease – No Systematically search area of suitable forest habitat – No
2015(5-year review)	Unknown	0	Improve habitat conditions; decrease threat of avian disease; systematically search area of suitable forest habitat;	Improve habitat conditions – Yes Decrease threat of avian disease – No Systematically search area of suitable forest habitat – No

Table 2. Threats to pououli and ongoing conservation efforts.

Threat	Listing factor	Current Status	Conservation/ Management Efforts
Ungulates – degradation of habitat and herbivory	A, C, E	Ongoing	Partially, some habitat areas fenced
Invasive introduced plants	A, E	Ongoing	Partially, some habitat areas managed
Low numbers	E	Ongoing	None
Climate change	A, E	Increasing	Partially, reforestation of some high elevation areas

References:

See previous 5-year review for a full list of references (USFWS 2010).

Atkinson, C. T., R. J. Dusek, K. L. Woods, and W. M. Iko. 2000. Pathogenicity of avian malaria in experimentally-infected amakihi. *Journal of Wildlife Disease* 36:197-204.

Atkinson, C. T., K. L. Woods, R. J. Dusek, L. S. Sileo, and W. M. Iko. 1995. Wildlife disease and conservation in Hawaii: Pathogenicity of avian malaria (*Plasmodium relictum*) in experimentally infected iiwi (*Vestiaria coccinea*). *Parasitology* 111:S59-S69.

Banko, W. E., and P. C. Banko. 2009. Historic Decline and Extinction. Pages 27-58 in T. K. Pratt, C. T. Atkinson, P. C. Banko, J. D. Jacobi, and B. L. Woodworth (eds.). *Conservation biology of Hawaiian forest birds: Implications for island avifauna*. Yale University Press, New Haven and London.

Benning, T. L., D. LaPointe, C. T. Atkinson, and P. M. Vitousek. 2002. Interactions of climate change with biological invasions and land use in the Hawaiian Islands: Modeling the fate of endemic birds using a geographic information system. *PNAS* 99:14246-14249.

Elphick, C. S., D. L. Roberts, and J. M. Reed. 2010. Estimated dates of recent extinctions for North American and Hawaiian birds. *Biological Conservation* 143: 617-624.

Fitzpatrick, J. W. Ivory-bill Quest. *Birdscope* (Spring 2002) 16(2):1-2. Available online at: http://static.birds.cornell.edu/Publications/Birdscope/Spring2002/Ivory-bill_Search.html.

ICAP. 2010. Hawaii's changing climate: Briefing sheet, 2010. Fletcher, C. (ed.). University of Hawaii Sea Grant College Program, Center for Island Climate Adaptation and Policy. Available online at: http://www.soest.hawaii.edu/coasts/publications/ClimateBrief_low.pdf.

IPCC. 2013. Summary for policymakers. *In: Climate Change 2013: The physical science basis. Contribution of the Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Stocker, T. F., D. Qin, G. -K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P. M. Bidgley (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, New York.

LaPointe, D. A., Atkinson, C. T., and S. I. Jarvi. 2009. Managing disease. Pages 405-424 in T. K. Pratt, C. T. Atkinson, P. C. Banko, J. D. Jacobi, and B. L. Woodworth (eds.). *Conservation biology of Hawaiian forest birds: Implications for island avifauna*. Yale University Press, New Haven and London.

Lia, W., O. E. Timm, C. Zhang, C. T. Atkinson, D. A. LaPointe, and M. D. Samuel. 2015. Will a warmer and wetter future cause extinction of native Hawaiian forest birds? Available on line at:
<http://onlinelibrary.wiley.com/doi/10.1111/gcb.13005/pdf>.

Reynolds, M. H., and T. J. Snetsinger. 2001. The Hawaii rare bird search 1994-1996. *Studies in Avian Biology* 22:133-143.

Scott, J. M., S. Mountainspring, F. L. Ramsey, and C. B. Kepler. 1986. Forest bird communities of the Hawaiian Islands: their dynamics, ecology, and conservation. *Studies in Avian Biology* 9:1-431

[USFWS] U.S. Fish and Wildlife Service. 2006. Revised recovery plan for Hawaiian forest birds. Region 1, Portland, Oregon. 622 pages.

[USFWS] U.S. Fish and Wildlife Service. 2010. Poouli (*Melamprosops phaeosoma*) 5-year Review Summary and Evaluation. U.S. Fish and Wildlife Service, Pacific Islands Fish and Wildlife Office, Honolulu, Hawaii. 13 pages.

[USFWS] U.S. Fish and Wildlife Service. 2012. Endangered and threatened wildlife and plants; listing 38 Species on Molokai, Lanai, and Maui as endangered and designating critical habitat on Molokai, Lanai, Maui and Kahoolawe for 135 species. *Federal Register* 77:34464-34775.

Wallace, G. E. 2010. Using autonomous recording units to monitor endangered Hawaiian seabirds. *Connect* (August 2010): 26. Available on line at:
https://www.aza.org/uploadedFiles/Conservation/Funding/AZA_Conservation_Endowment_Fund/Information_for_Applicants/CEF_Previous_Awards/07-744UseofAutonomousRecordingUnitstoDetectRareHawaiianBirdsAugust10.pdf.

Yorinks, N., and C. T. Atkinson. 2000. Effects of malaria on activity budgets of experimentally infected juvenile apapane (*Himatione sanguinea*). *The Auk* 117:731-738.

Personal communication:

Camp, Rick. 2015. U.S. Geological Survey. E-mail communication with Jay Nelson, Pacific Islands Fish and Wildlife Office, dated May 6, 2015. Subject: Recent forest bird surveys – islands and area(s) surveyed.

**U.S. FISH AND WILDLIFE SERVICE
SIGNATURE PAGE for 5-YEAR REVIEW of
Poouli (*Melamprosops phaeosoma*)**

Pre-1996 DPS listing still considered a listable entity? N/A

Recommendation resulting from the 5-year review:

- Delisting
- Reclassify from Endangered to Threatened status
- Reclassify from Threatened to Endangered status
- No Change in listing status

Appropriate Listing/Reclassification Priority Number, if applicable: _____

for Programmatic Deputy Field Supervisor, Pacific Islands Fish and Wildlife Office

Maria M. Bruegmann

Date 2015-08-20