

**DEVELOPMENT OF A FRAMEWORK APPLICABLE FOR MANAGED WETLANDS:  
STATE-AND-TRANSITION MODELS FOR WETLAND HABITAT ASSESSMENT AND  
MANAGEMENT**

**PROCEDURES MANUAL**



Photo by E. S. Peckham

U.S. Department of the Interior  
27650B South Valley Road  
Lima, Montana 59739

Phone: (406)276-3536

4<sup>th</sup> edition  
June 2018

## **TABLE OF CONTENTS**

Introduction.....	1
1. General Information.....	5
Safety.....	5
Training.....	5
Preparations to be Made Before Going Afield.....	8
Equipment to be Taken Afield.....	9
2. Surveys and Monitoring.....	10
Sampling Design.....	10
Timing of Surveys.....	12
Vegetation Surveys.....	12
Abiotic Variables.....	17
Wetland Hydrology.....	22
3. Data Management.....	23
Data Forms and Data Storage.....	23
Data Entry and Editing.....	23
Literature Cited.....	24
Appendix I. Data Forms.....	26
Appendix II. How to Build a 1m × 1m Plot Frame.....	29
Appendix III. How to Calibrate a YSI 63 Water Quality Meter.....	31
Appendix IV. Description of Wetland Soil Analysis.....	39
Appendix V. Double Sampling.....	42
Appendix VI. Guide to Wetland Indicator Species.....	43

## INTRODUCTION

This manual is intended to assist participants in understanding the purpose of this study as well as provide information and rationale behind the techniques employed. We understand that this manual cannot be comprehensive, but hope that it will serve as a useful guide. If you find areas of the manual confusing, don't hesitate to ask for an explanation. Comments and constructive criticism regarding the manual are welcomed. To help prepare this manual, we "borrowed liberally" from the Prairie Habitat Joint Venture Assessment Program: Procedures Manual, 7<sup>th</sup> edition (Institute for Wetland and Waterfowl Research, Ducks Unlimited Canada, Stonewall, Manitoba, Canada).

### **Background**

Intermountain West wetlands provide important habitat for migratory birds and other wetland-dependent wildlife. Similar to wetland habitats in other regions of North America, agriculture and urban development have resulted in the loss of approximately 57% of Intermountain West wetlands to drainage (Ratti and Kadlec 1992). The significance of this loss is magnified due to the region's largely arid landscape. Moreover, anthropogenic modification of wetlands has altered historical wetland function; however the extent and degree of altered functions has not been quantified. Wetland loss and degradation, and the importance of these wetland resources to migratory birds, has led to the establishment and active management of some of the Intermountain West's largest wetland complexes, such as National Wildlife Refuges (NWRs) (e.g., Alamosa-Monte Vista, Malheur, Red Rock Lakes, and Ruby Lake), state Wildlife Management Areas (WMAs) (e.g., Farmington Bay, Market Lake), and private duck clubs (e.g., Ambassador Duck Club, Newstate Duck Club).

Objectives for managing wetland habitat for migratory birds and other wetland-dependent wildlife within the Intermountain West typically fall into two classes: wetland function based on a desired plant community, or habitat for a specified population size and/or life-history requirements of a focal wildlife species or suite of species. In the former case, a wetland is commonly managed to maintain a desired reference submerged aquatic vegetation (SAV) community based on 1) historical accounts of SAV community composition, and/or 2) relative quality of SAV species necessary for migratory birds (commonly waterfowl). For the latter scenario, habitat objectives for SAV could be based upon bioenergetics models defining the necessary biomass of species or communities of SAV providing forage to support population objective(s) of migratory birds (again, commonly waterfowl).

Building a common management framework encompassing these, and other, types of wetland objectives is challenging and has not been accomplished to date. Much of the difficulty lies in incorporating ecological processes that define a potential range of states of a wetland considering management, i.e., fitting the objective within a site's potential. A wetland will have multiple potential states whose expression can be influenced by management. Therefore, it is essential to 1) determine the potential range of states a wetland can express, 2) define which species or group of species and associated life cycles are to be managed for, and 3) link 1 and 2 as a basis to determine management prescriptions that provide the preferred state within dynamic and productive wetland cycles for the target species or group of species.


Wetland management actions largely center on mimicking natural disturbance processes to maintain ecological function of managed wetlands in the Intermountain West. The primary process of management interest is the dynamic wet/dry hydrological cycle, which is a key driver of wetland productivity and vegetation community structure (see reviews in Murkin et al. 1997, Mitsch and Gosselink 2007). The ability to manipulate the timing and duration of flooding in managed wetlands permits controlling, to some degree, the primary wetland disturbance regime. Water-level manipulations can be considered a perturbation, where the ability to predict the outcome of such manipulations varies dependent upon the knowledge of the system being manipulated. Importantly, ecological systems (including wetlands) commonly respond to perturbations in a non-linear fashion with multiple states possible (Drake 1990, van der Valk 1981, Zweig and Kitchens 2009, Smith 2012). This can be contrasted with linear succession to a climax seral community as initially espoused by Clements (1936). Application of linear climax theory to management has proven largely unfruitful (Stringham et al. 2003), and led to the development of non-linear state and transition models (Westoby et al. 1989).

State and transition models (STMs) provide a framework to address the needs described above. An STM depicts the current knowledge of ecological dynamics on a site, identifying the range of potential vegetation communities, i.e., states, which could exist. The STM also identifies the conditions, disturbances, and management actions that may cause a site to transition among states (e.g., from submerged aquatic vegetation to emergent vegetation in a wetland) or simply shift among phases of a state (e.g., from milfoil-dominated to pondweed-dominated within the submerged aquatic vegetation state). Therefore, STMs can assist in making management decisions by identifying actions to maintain a current state, or those that would likely result in a transition to a more preferred state.

We have developed a draft STM for semi-permanently flooded wetlands in the Intermountain West (Fig. 1). It is based on conceptual ecological models that include drivers, stressors, and effects on performance measures (Busch and Trexler 2003). Conceptual models will assist in another important step – identifying possible ecological, utility, and decision thresholds (Martin et al. 2009). Ecological thresholds are commonly defined as a point or zone along a continuum of a system variable (or suite of variables); when it is crossed, there is a sudden transition in the system state (Huggett 2005, Bennetts et al. 2007). Utility thresholds are values of system state or key drivers of system state at which small changes result in significant response in the management outcome (Martin et al. 2009).

A multi-region FWS working group is also currently developing a bottom-up, empirically-driven STM using data from two distinct sites: Lower Red Rock and Malheur lakes. The working group believes that these 2 approaches to STMs will provide 1) a more universally-applicable framework to apply across stations, WMAs, and regions, 2) a more comprehensive understanding of the process of STM development, and 3) a broader tool-set of analyses useful for creating and validating STMs for individual stations. A top-down development of a wetland STM will help elucidate common ecological drivers and processes that influence the expression of wetland plants.





**Figure 1.** Draft state-and-transition model (STM) for semi-permanently flooded wetlands. Six states are defined in the STM, with state 1.0 Diverse SAV & Tall Emergent the reference state. Within the reference state six vegetation community phases exist. Arrows between states represent transition (‘Txx’) and restoration (‘Rxx’) pathways. Dashed lines represent 1) unintended outcomes of restoration efforts, i.e., restoration pathways to altered states that could result from management actions intended to return a wetland to the reference state, or 2) partial restorations, e.g., removing artificial nutrient inputs, but not restoring hydrology. Bestelmeyer et al. (2010) is an excellent resource for clarification on the concepts and terminology of STMs.

## Objectives

The objectives of the field work to be conducted in 2015 vary by participating station, and include:

- ◆ Conduct an inventory of vegetation, hydrologic conditions, and water quality within semi-permanently flooded wetland habitats on priority units of participating stations.
- ◆ Re-sample semi-permanently flooded wetlands on selected units evaluated in 2014.

- ◆ Implement refined methodologies for collecting consistent wetland vegetation data for characterizing a wetland's current condition across diverse landscapes in the Intermountain West and western Prairie Pothole Region.
- ◆ Describe, qualitatively and quantitatively, the ecological and abiotic conditions of wetland states and vegetation community phases to inform future management actions and the current draft state-and-transition model at multiple scales.
- ◆ Identify preliminary indicator species and abiotic variables for states and phases for streamlining future monitoring efforts to quantify wetland status and assess response to management actions.

### **Study Area**

The 2015 field work will be conducted across eight NWRs within USFWS regions one and six. Participating stations include: Bear Lake NWR, Bowdoin NWR, Camas NWR, Fish Springs NWR, Grays Lake NWR, Lee Metcalf NWR, Malheur NWR, and Medicine Lake NWR. Each station selected priority management units for sampling.

## CHAPTER 1

### GENERAL INFORMATION

#### Safety

Watercraft will be used frequently during fieldwork. Personal flotation devices (PFDs) **MUST BE WORN AT ALL TIMES WHILE IN A WATERCRAFT**. All other FWS field and safety protocols should be strictly followed.

#### Training

It is very important for training to take place prior to field sampling to 1) maximize the repeatability of field data collection, i.e., minimize among-observer variation, 2) ensure consistent data collection among participating stations, and 3) make field data collection as efficient as possible. The two most common sources of observer error in a project such as this are 1) mis-identification of wetland plants and 2) differences in ocular estimates of percent cover. An investment in training before going to the field can greatly reduce both these types of errors. Stations are generally far from the next nearest station participating in this project, so it will be rare that training will occur with staff from multiple stations. We have tried to provide detailed instructions for training so that training occurs similarly across participating stations. We have also tried to make the training as efficient and effective as possible.

*Plant Identification*—It will be imperative to spend time reviewing common wetland species you expect to encounter on your unit(s). Local knowledge of wetland plant communities is very helpful; if you are new to an area it will be worth spending time looking at plant inventories available in vegetation maps, reports, plans (e.g., Comprehensive Conservation Plans), theses, etc. If you've sampled wetlands on your station previously you may already have a list of common species for the area. The Consortium of Pacific Northwest Herbaria (<http://www.pnwherbaria.org/>) is also a useful resource for species lists in your county (and adjacent counties). These resources will be very helpful when keying out wetland plants, but don't try too hard to make an unknown plant key out to one on your list (many of us have made that mistake before!). When keying out plants there are numerous manuals to choose from; a brief list is provided below. Some of the manuals listed are not current with plant scientific names. The Integrated Taxonomic Information System (<http://www.itis.gov/>) should be used for accepted common and scientific names; the US Department of Agriculture's PLANTS database (<http://plants.usda.gov/java/>) is the source for standardized plant species symbols used on field data forms (e.g., Northwest Territory sedge, *Carex utriculata*, CAUT). The codes for bare ground/open water (BASU) and residual vegetation (RESID) are included on the datasheet. Lastly, if you can't identify a plant while in the field collect a specimen for later identification (specimens should be placed in a small sealable plastic bag with a small amount of water; try to collect a full plant, i.e., roots or rhizomes should be included).

Useful identification manuals for wetland vegetation in western North America:

Chadde, S. W. 2012. Wetland plants of the Northern Great Plains: a complete guide to the wetland and aquatic plants of North and South Dakota, Nebraska, eastern Montana, and eastern Wyoming. CreateSpace Independent Publishing Platform. (Update to Larson 1993.)

DiTomaso and Healy 2003. Aquatic and riparian weeds of the West. University of California

Fassett, N. C. 1957. A manual of aquatic plants. University of Wisconsin Press, Madison, Wisconsin.

Guard, B. J. 1995. Wetland plants of Oregon and Washington. Lone Pine Publishing, Auburn, Washington.

Hamel, K., J. Parsons, M. Boule, S. Feldman, I. Wertz, and L. Zempke. 2001. An aquatic plant identification manual for Washington's freshwater plants. Washington Department of Ecology Publication No. 01-10-032. <http://www.ecy.wa.gov/PROGRAMS/wq/plants/plantid2/index.html>

Hitchcock and Cronquist 1973. Flora of the Pacific Northwest: an illustrated manual. University of Washington Press, Seattle, Washington.

Hotchkiss, N. 1972. Common marsh plants of the United States and Canada. Dover Publications, Inc. New York, New York.

Hurd, E. G., S. Goodrich, and N. L. Shaw. 1994. Field guide to intermountain rushes. US Department of Agriculture Forest Service General Technical Report INT-306. <http://www.treeseearch.fs.fed.us/pubs/24234>

Hurd, E. G., N. L. Shaw, J. Mastrogiuseppe, L. C. Smithman, and S. Goodrich. 1994. Field guide to intermountain sedges. US Department of Agriculture Forest Service General Technical Report RMRS-GTR-10. [http://www.fs.fed.us/rm/pubs/rmrs\\_gtr010/rmrs\\_gtr010\\_01intro.pdf](http://www.fs.fed.us/rm/pubs/rmrs_gtr010/rmrs_gtr010_01intro.pdf)

Larson, G. E. 1993. Aquatic and wetland vascular plants of the northern Great Plains. US Department of Agriculture Forest Service General Technical Report RM-238. <http://www.treeseearch.fs.fed.us/pubs/30420>

Lesica, P., and P. Husby. 2001. Field guide to Montana's wetland vascular plants. A non-technical key to the genera with keys to the species of sedges and rushes. Montana Wetlands Trust, Helena, Montana.

Also, a great reference for relationships among some abiotic conditions and wetland plants based largely on data from the Midwest, but applicable to this project, is the US Geological Survey Scientific Investigations Report 'Literature review and database of relations between salinity and



aquatic biota: applications to Bowdoin National Wildlife Refuge, Montana', which is available at <http://pubs.usgs.gov/sir/2009/5098/pdf/sir2009-5098.pdf>.

If you need references for assistance with plant terminology the following may help:

Beentje, H. J. 2010. The Kew plant glossary: an illustrated dictionary of plant terms. Royal Botanic Gardens, Kew, U.K.

Harris, J. G., and M. W. Harris. 2001. Plant identification terminology: an illustrated glossary. Springer Lake Pub.

*Ocular Cover Estimation*—‘Training’ your eye to consistently estimate percent canopy cover of a species in a quadrat is an important step in preparing for field data collection. This can be done with simple cutouts in various shapes and sizes of a known area. Preparing a set of cutouts can be done with cardboard, card stock, construction paper, etc., and the cutouts can be saved for training in subsequent years. Start by cutting out the shapes listed in Table 1 and labeling them as one of five species. You could simply use ‘Spp. 1’, ‘Spp. 2’, etc., or you could pick five species you expect to find and label the cutouts using actual species names. Next, print out several data sheets to record estimates on. Place a PVC plot frame (see Appendix II for instructions to build a frame) on the ground and have someone put a series of cutouts in the quadrat for each person to record. Have everyone participating record their observations on a data sheet following the protocol described in the Vegetation Surveys section of chapter two. Compare everyone’s results with the ‘truth’ and discuss differences among observers. Repeat this with different combinations of cutouts until the observers have ‘repeatable’ results across a broad diversity of cutout combinations.

**Table 1.** Inventory of cutouts to be used for training observers for ocular estimation of percent canopy cover prior to field data collection.

Shape	Size	Area (m <sup>2</sup> )	Dimensions <sup>a</sup> (cm)	No. of Cutouts
Circle	Large	0.126	20	5 <sup>b</sup>
Circle	Medium	0.031	10	11 <sup>c</sup>
Circle	Small	0.005	4	12 <sup>d</sup>
Rectangle	Extra Large	0.780	100 × 78	1
Rectangle	Large	0.550	100 × 55	1
Rectangle	Medium	0.360	90 × 40	2
Rectangle	Small	0.05	25 × 20	5 <sup>b</sup>
Square	Large	0.640	80 × 80	1
Square	Medium	0.410	64 × 64	3
Square	Small	0.090	30 × 30	5 <sup>b</sup>

<sup>a</sup>Radius provided for circles.

<sup>b</sup>Labelled such that one species has two cutouts, three species have one.

<sup>c</sup>Labelled such that one species has five cutouts, one species has three, and three species have one.

<sup>d</sup>Labelled such that one species has five circles, one species has four circles, and three species have one circle.

After the observers are trained using the known percent cover cutouts, practice cover estimates of **both** submerged and tall emergent species in the field. Have the cutouts available to discuss discrepancies among observers.

*GPS*— Please spend some time familiarizing yourself with the GPS unit you'll be using. It will be necessary to quickly and efficiently navigate to sampling points using a GPS unit while in the field. If you do not have experience navigating to points, or want a refresher, work with other staff at the station to practice finding points. For example, one person can mark the location of an object (e.g., a beach ball) in tall emergent vegetation and the other person can locate it using the GPS unit. It is important that you set your GPS unit to the same datum of the GRTS sample points (see Chapter 2), which is **WGS 1984**. You should have been provided a .csv file that contains location data for each GRTS sample point by sampling unit. If you need assistance importing that data into your GPS unit you can contact Jenny Barnett ([jenny\\_barnett@fws.gov](mailto:jenny_barnett@fws.gov); 509.380.6479) or Jeff Warren ([jeffrey\\_warren@fws.gov](mailto:jeffrey_warren@fws.gov); 406.276.3536 ext. 106).

*Water Quality*—It is important to be familiar with your water quality meter and to calibrate it following the manufacturer's recommendations. Calibration buffer (pH and conductivity) will be needed, so plan ahead and make certain you have those on hand prior to calibrating your unit for sampling. Instructions for calibrating a YSI 63 meter, a common meter used for collecting water quality data, are provided in Appendix III. Note that pH should be calibrated at least each year, while specific conductivity will rarely need recalibrated. Follow calibration and storage instructions for your meter and sensors. The complete operations manual for a YSI 63 can be downloaded at <http://www.ysi.com/media/pdfs/031178-YSI-Model-63-Operations-Manual-RevC.pdf>. The water quality meter and other equipment should be cleaned between units and/or stations.

*Soil Texture*—In addition to becoming familiar with the soil texture diagrams in Figures 6 and 7, it is helpful to practice identifying soil type by texture before the sampling effort. In 2014, the SE Idaho field crew had access to lab-calibrated samples and a NRCS soil scientist mid-season, and reported that this level of training greatly increased their accuracy and confidence in the field.

### **Preparations to be Made Before Going Afield**

The bulleted list below is intended to help you get ready for going into the field to collect data. There are a few items that will need to be purchased if you don't already have them on hand, e.g., a water quality meter. If you need to borrow equipment for sampling this year call Jeff because there are some items on the list below that are available for loaning out to stations.

- Plant identification training
- Soil texture identification/calibration training
- Ocular cover estimation training
- GPS training
- Create and print maps of GRTS points (datum WGS 1984)
- Print data forms (Appendix I)

- Build PVC plot frame for sampling vegetation (Appendix II)
- Calibrate water quality meter (Appendix IV for model YSI 63)
- Attach secchi disc to 5–30 m fiberglass tape (attach end of tape using eyebolt on disc)
- Build rake sampler (see Figs. X and X; text pg X)
- Build/purchase aqua-viewer (<http://www.wildco.com/Fieldmaster-Aquavue.html>)
- Purchase soil corer ([http://www.forestry-suppliers.com/product\\_pages/Products.asp?mi=31233&title=Oakfield+Model+L+Tube+Sampler+Soil+Probe&itemnum=77117](http://www.forestry-suppliers.com/product_pages/Products.asp?mi=31233&title=Oakfield+Model+L+Tube+Sampler+Soil+Probe&itemnum=77117))
- Organize PPE (e.g., PFDs, polarized glasses, sun block, radio, etc.)

### **Equipment to be Taken Afield**

- Maps
- Data forms
- Vegetation manuals
- Hand lens
- GPS and spare batteries
- PVC plot frame
- Water quality meter
- Secchi disc with fiberglass tape
- Rake sampler
- Aqua-viewer
- Soil corer
- Plastic bags to collect unidentified species samples
- Pencils
- Permanent marker (for labeling vegetation sample, soil sample bags, etc)
- Compass
- Field notebook
- PPE (e.g., PFDs, polarized glasses, sun block, insect repellent, radio, etc.)
- Materials to anchor your boat/canoe (in open water habitat)

**WE MAY SEEM OVERLY CONCERNED WITH HOW THINGS GET DONE. HOWEVER, THIS IS NECESSARY BECAUSE MANY DIFFERENT PEOPLE WILL CONTRIBUTE TO THE PROJECT OVER MANY YEARS. WE MUST ACHIEVE CONSISTENT DATA COLLECTION TO ACHIEVE COMPARABLE RESULTS.**

## CHAPTER 2

### SURVEYS AND MONITORING

#### Sampling Design

Our sampling frame is the **entire collection of geographic areas across all stations classified as potential semi-permanently flooded wetland**. Each station can be considered a distinct geographic "strata." Within each station we have multiple delineated areas (hereafter, "unit" = primary sampling unit), and within each area we will have plots where vegetation and covariates are measured (hereafter, "points" = secondary sampling units). As the units were not selected randomly from a list of possible units in 2014, we can think of this design as a nested stratified design with strata1 = stations and strata 2 = units that are nested within strata 1 or stations.

A generalized random tessellation stratified (**GRTS**) sample of points was created for each unit identified as a sampling priority by a station. The GRTS algorithm provides a spatially-balanced random sample of ordered points to be visited for data collection. We used the `grts()` function within the `spsurvey` R package (Kincaid 2013). Sampling intensity, defined as points per acre  $\times 100\%$  (e.g., 30 points/100 acres  $\times 100\% = 30\%$  sampling intensity), was specified at 30% within most units (i.e., those 100–300 acres in area) with a minimum of 30 points for smaller units (20–100 acres), 90 points for units between 300 and 500 acres, and a maximum of 120 points for larger units (>500 Acres). Units within a station were stratified to produce an independent GRTS sample for each unit within each station.

The output of the GRTS sample was saved as a .csv file with GRTS point locations ("xcoord" and "ycoord") provided in decimal degrees, datum **WGS 1984**. The first column, **GRTS\_ID**, is a concatenation of several pieces of information found in the .csv file. The first 3 letters are a literal for your station (e.g., LMC for Lee Metcalf), followed by an abbreviation for the unit (e.g., PO10 for Pond 10), the panel (i.e., PO for PanelOne or OS for OverSamp), and finally a consecutive numbering of points for each panel. The highest point number within a panel for a unit is the total number of suggested survey points. The **panel** column will be either PanelOne or OverSamp; the former comprises the primary set of GRTS points for sampling and the latter are replacement points if a PanelOne point needs to be excluded from the sample (see **Dropping Points** below). The .csv file therefore contains *twice* as many GRTS points as necessary if all of the PanelOne points are sampled. The **siteID** column in these files has a consecutive numbering of points across the entire station. The unit name is listed in the **stratum** column with the station literal preceding it (e.g., LMC – Pond 10).

#### ***Order of GRTS Points***

As mentioned above, the GRTS sample is a spatially-balanced random sample of *ordered* points. If you can sample the complete set of PanelOne GRTS points in a unit the ordering is not relevant to your field effort. However, if you determine that you can only sample a subset of GRTS points, it is important to select an ordered subset for sampling. For example, if you have 30 GRTS points for a unit, but determine *a priori* that you only have time to sample 20, your

subset should be 1–20 (or some similarly ordered subset). This will maintain the spatially-balanced nature of the sampling, minimizing over- or under-sampling areas of the unit. Once you determine the ordered subset of GRTS points to be sampled in a unit, the actual order in which points are visited in the field is irrelevant and should be organized for efficiency. If PanelOne points need to be dropped from the sample (see ***Dropping Points*** below) you will need to sample OverSamp points, also in order. For example, if you dropped three PanelOne points (*any* three) you would use OverSamp points 1–3 as replacements. If one of the first three OverSamp points needed to be dropped, you would continue in order, i.e., move to point 4. Replacement points will likely not be close to the dropped points due to the spatial balance of the sampling design.

We want stations to focus on complete sampling of priority units. For example, if a station has 6 units with 50 points each, and you only have time to sample 150 points, please sample 50 points at the 3 highest priority units rather than 25 points at 6 units.

### ***Dropping Points***

It is important to keep good records of each point, identifying if the point was surveyed or not and if not surveyed, recording the reason. This record keeping for both the PanelOne and OverSample points is critical for adjustment to the design-weights after field surveys are conducted.

There are several reasons why data may not be collected at a point sampled but most likely because the point was not in the appropriate habitat type (see below) or it was logistically inaccessible. If a point was visited and deemed non-target (e.g., point fell within uplands) or in the office it was deemed non-target, it needs to be denoted as such on the data sheet (see header). No vegetation data needs to be recorded for that point, although **it may be helpful to list recognizable species and give them a canopy cover of 0**. This will help validate a point determined in the field to be non-target due to wetland permanency class. Standard codes are: TS = target and sampled, NT = non-target, IN = inaccessible, and NotEval = not evaluated. Within the EvalReason column denote why a point was deemed non-target (NT) or inaccessible (IN). EvalReason codes include:

UPL (upland e.g., sagebrush, lodgepole pine) note habitat type

TF (temporarily flooded, e.g., tufted hairgrass) note dominant vegetation

SF (seasonally flooded, e.g., Baltic rush, sedge) note dominant vegetation

PF (deep > 3 m permanently flooded)

IN (inaccessible) note why a point was inaccessible

OTHER please explain (e.g., muskrat house, swan nest platform, vegetation uprooted by foraging)

NOTE: A semi-permanently flooded area that is dry at the time of sampling **IS** a target point and **SHOULD** be sampled.

If you are not familiar with vegetation types in wetlands with different flooding regimes, work with station staff to identify these habitat types at the station. A breakdown of indicator species by habitat type is provided in Appendix VI. Another great reference for plant species as indicators of wetland permanency class is Stewart and Kantrud (1971), *Classification of Natural Ponds and Lakes in the Glaciated Prairie Region*. Appendix A of this publication has a list of characteristic plants species in prairie wetlands. A pdf of this resource publication is available online at <http://esrd.alberta.ca/lands-forests/shorelands/documents/ClassificationPondsLakesPrairie-Part1.pdf>.

An example of an SOP that sampled using GRTS that might be useful:

Yeo, J.J., T.J. Rodhouse, G.H. Dicus, K.M. Irvine, L.K. Garrett. 2009. 1 Upper Columbia Basin Network sagebrush steppe vegetation monitoring protocol: Standard operating procedures version 1.0. Natural Resource Report NPS/UCBN/NRR---2009/142. National Park Service, Fort Collins, Colorado.

### **Timing of Surveys**

*When* surveys should be conducted is as important as *where* they are conducted, the latter discussed in detail above. When surveys should be conducted can be considered at two temporal scales, 1) season within a year, and 2) period within a season. First, the season within a year that sampling should occur is during peak growing season for wetland vegetation. This allows species with varying phenology to reach their full growth potential, as well as ensure seeds, or other characteristic reproductive parts, are present to assist with species identification. For most stations participating in this project this will be July through August, with lower elevation/warmer sites having an earlier phenology than higher elevation/cooler sites. The phenology at a station among years may also vary; a warm, dry spring may result in peak growing season occurring several weeks earlier than a cool, wet spring. When to conduct sampling within the growing season is driven by local conditions that may influence your ability to detect aquatic species. For example, heavy thunderstorms could result in localized and short-term turbid conditions on a unit to be sampled. Postponing sampling a few days to let the turbidity settle out would be warranted in this situation.

### **Vegetation Surveys**


Vegetation surveys will consist of 1 × 1 m quadrats positioned such that the GRTS point location demarcates the northwest corner of the quadrat. To not bias placement of the plot, do not look at the vegetation before placing the plot. For surveys conducted from a boat the quadrat is placed on the east side of the boat (Fig. 2) and ocular estimates of percent cover will be recorded for each species (see *Ocular observations* below). If the water in a unit is too turbid to see the bottom of the quadrat, rake subsamples will be taken to characterize SAV. These subsamples



comprise three juxtaposed  $1 \times 0.35$  m rake samples collected within a  $1 \times 1$  m quadrat (Fig. 3; see *Rake observations* below) (Yin et al. 2000).

All surveys should be conducted with 1 observer and 1 recorder. The observer may sit in the bow or stern of the boat, and will locate each GRTS point. The boat should be oriented with the stern due south of the bow. The boat should be steadied either with an anchor at each end, or, in shallow water when using a canoe, paddles pushed into the substrate will often suffice to hold position.

### Submerged Aquatic Vegetation Survey for Clear Water



**Figure 2.** Placement of submerged aquatic vegetation survey quadrat in relation to the GRTS point and canoe or boat.

### *Aquatic Vegetation*

*Ocular observations*—For each quadrat, percent canopy cover will be recorded for each species present. Quadrats will be surveyed from canoe or boat when possible, or on foot in areas of low water and/or high density emergent vegetation. Ocular estimates should be collected first after arriving at a sample point in order to reduce the potential of stirring up bottom sediment that can obscure vegetation. Localized turbidity may preclude you from observing submerged vegetation at some points in a unit that has otherwise clear water. This should be obvious when you first arrive at a point, i.e., if you can't see the bottom substrate due to turbidity (**not because SAV**

**obscures the bottom substrate**). In this situation the point should be sampled using a rake (see *Rake observations* and *Secchi disc depth* below). The data can be used to define vegetation community phases because that uses presence/absence data (assuming detection probability does not differ significantly between the two methods). We will have to exclude those points from unit-level summaries of canopy cover, however.

Once the sampling frame is positioned, the observer should assess the canopy cover of **each species, residual vegetation (i.e., last year's standing dead vegetation), and bare substrate** in the quadrat. It is imperative that the observer uses polarized glasses at this step to maximize visibility into the water column. An aqua viewer should also be used to view SAV and ensure that all vegetation species within the plot are recorded (<http://www.wildco.com/Fieldmaster-Aquavue.html>; they can also be relatively easily built). Slowly sweep the aqua viewer through the water to examine the vegetation present, careful not to disturb vegetation and bottom sediment. Start with species that occur as a single plant in the plot and work up to the species that dominate the quadrat, recording the percent of the plot that is bare substrate last. **For every species present, percent cover should be estimated to the nearest 1%.** Species that occur as a single plant should be recorded as having at least 1% canopy cover; they could have more than that, but not less (i.e., it is o.k. to record 3% canopy cover for a single plant, but not 0.5%). When you have recorded the percent canopy cover for all species present and bare substrate, **be sure to check that the total sums to 100%.**

In 2015, we are separating “Residual vegetation” into the following three categories:

- RESID – TE; Tall emergent (e.g. hardstem or softstem bulrush, cattail),
- RESID – SE; Short emergent (e.g. alkali bulrush, Northwest Territory sedge), and
- RESID – Su; Submerged (**only** for current-year SAV on a mudflat).

Please record the category of residual cover at each point. If more than one residual vegetation category is present within a plot, a separate percent cover should be recorded for each category. Tall and short emergent categories are named in reference to potential species height, not the actual height of the residual vegetation. For example, residual cattail that is lying at or near the water's surface should be recorded as tall emergent residual cover. The residual SAV category is **only** for the situation where dead SAV from the current year is found on a mudflat.

Floating-leaved species (e.g., duckweed (*Lemna* spp.), floating pondweed (*Potamogeton natans*), bladderwort (*Utricularia* spp.)) may obscure the observers ability to see submerged species in the water column. Observers should record aerial coverage of each plant species as it looks from above, even if the floating-leaved species obscures other species. Once canopy cover has been estimated, please record obscured species in the notes for that point (you do *not* need to record canopy cover of the obscured species, simply their presence).

We have modified the sampling protocol to better address filamentous algae based on feedback from field crews and preliminary results from the 2014 field season. Filamentous algae will be treated similarly to waterlily (*Nuphar* spp.; see below), i.e., the canopy cover of filamentous algae will not be included in the canopy cover estimate of wetland plants. Filamentous algae should be recorded in one of two categories – FILAL-FL (floating) or FILAL-WC (water column; includes algae on the surface of SAV). Observers should sequentially evaluate and

remove the algae layer(s) until they can estimate SAV canopy cover. Algae can be relatively easily moved out of the way using your canoe paddle. At points with considerable algae cover on the SAV you won't be able to clear it all away – infer SAV canopy cover assuming the FILAL is not there. Lastly, at a site with deeper water, it may not be feasible to move aside all of the algae in the water column at depth. This percentage cover should be recorded as “FILAL” in the SAV stratum and will be summed as part of the total cover of that plot.

You will also likely encounter floating plant parts from current year growth in your frame during surveys, often the result of grazing by waterbirds. These species should **not** be included in your estimate of aerial cover in your frame; remove them if necessary to get a clear view of the frame. Be certain that you do not exclude non-rooted species such as coon's tail (*Ceratophyllum demersum*) or common bladderwort (*Utricularia macrorhiza*) as floating plant parts.

Waterlily (*Nuphar* spp.) presents a unique issue for sampling due to the growth form of this genus. If waterlily is present, ocular estimates will be made for 2 strata: 1) vegetation at or above the surface of the water, and 2) vegetation within the water column, including all species (SAV, waterlily stems, etc). For total cover, the ocular estimate of waterlily above the water column will not be included. The percent cover of the strata within the water column should still sum to 100%. Ocular estimates for sites where smartweed (*Persicaria* spp.) occurs at or above the surface of an otherwise open water site should similarly be assessed in two different strata.

Any unidentified species should be collected and placed in a labeled plastic bag with a small amount of water for keying out after the survey.


Rake observations—In units where water turbidity precludes obtaining ocular estimates of canopy cover it will be necessary to collect rake samples to characterize aquatic vegetation. Canopy cover of visible SAV, emergent, and floating-leaved species should be estimated and recorded similar to ocular observation surveys before rake samples are collected. The rake sampler is the same as that described by Yin et al. (2000), which is modified from Jessen and Lound (1962) and Deppe and Lathrop (1992). The rake sampler comprises two square-headed garden rakes welded (or bolted, Fig. 4) together such that the teeth of each faces out. Rake heads should be 35 cm in width with 14 5-cm long teeth. Teeth are marked in 20% increments for estimating total plant density for each sample (Fig. 5). For species that occur as one plant in the rake sample, record the plant density as a 1 (Fig. 5).

Prior to sampling the gunwale of the boat should be marked at the 1) ends of the 1 × 1 m quadrat edges, and 2) center of each 1 × 0.35 m subsample (Fig. 3).

Rake subplot sampling should start with R1 and proceed from north to south until all three subplots are sampled (Fig. 3). Extend the rake out from the boat to the bottom of the wetland at the outer boundary of the imaginary 1 × 0.35 m rectangle within the 1 × 1 m quadrat. Drag the rake along the bottom for 1 m. Twist the rake 180° as it is lifted off of the bottom to bring the vegetation into the boat. Twisting minimizes the loss of plants from the rake; twisting more than 180° could cause plants to fall off. Plants hanging off of the rake head should be added to the teeth and those hanging from the handle should be ignored (Yin et al. 2000). Gently sweep the rake head in the water using a figure eight motion to compact and rinse vegetation. Rate the total

density of plants on the rake head according to the rake density categories in Table 2, then record the density of **each species** such that it sums to the total plant density recorded. Repeat these steps for rake subplots R2 and R3 (Fig. 3). To identify all the species in the rake sample, this may require poking through the rake sample with your fingers to better see plants present. Deppe and Lathrop (1992) state that separation of plants to assess individual coverage on the rake may be necessary, particularly when plants are entwined with each other.

### Submerged Aquatic Vegetation Survey for Turbid Water




**Figure 3.** Placement of the three rake sub-samples in relation to the GRTS point and canoe or boat.

**Table 2.** Rake density plant rating categories.

Density Rating	Percent Rake Teeth Filled
1	1% (i.e., a single plant)
2	1–20%
3	21–40%
4	41–60%
5	61–80%
6	81–100%
7	>100%

Any unidentified species should be collected and placed in a labeled plastic bag with a small amount of water for keying out after the survey.



**Figure 4.** Rake samplers can be made from two square-headed garden rakes that are 35cm wide with 14 5-cm long teeth. Rake heads can be welded or clamped together.



**Figure 5.** Rake sampler used to sample aquatic vegetation in turbid-water units (photo from Yen and Kreiling 2011). Arrows show the lines used to indicate increments used to estimate total plant density. Vegetation hanging from the rake teeth would be added to the rake before estimating total SAV density.

### **Abiotic Variables**

Abiotic characteristics known to influence wetland plant communities will be recorded during vegetation surveys. The method of survey (boat or foot), water depth ( $\pm 1$  cm), Secchi disc depth

( $\pm 1$  cm), temperature ( $\pm 0.1^\circ\text{C}$ ), pH ( $\pm 0.01$  units), salinity ( $\pm 0.1$  ppt), relative conductivity ( $\pm 1$   $\mu\text{S}/\text{cm}$ ), and soil texture (Table 3) will be measured at each GRTS point after vegetation data are recorded.

Survey method—Record how the survey was conducted, i.e., from a boat (B) or on foot (F).

Water depth—Lower the Secchi disc attached to the fiberglass tape to the bottom of the wetland. If dense vegetation prevents the disc from reaching the bottom, gently separate the vegetation to allow the Secchi disc to sit flat on the substrate. If you are still unable to get the Secchi disc to the substrate use a steel tape measure. In wetlands with unconsolidated bottoms it may be hard to determine where the water column ends and the substrate begins, but do your best. Measure water depth to the nearest 1 cm. **If the site is dry a water depth of ‘0’ should be recorded on the datasheet and in the database.** Circle ‘Other’ in the Depth section of the data form if you used something other than a Secchi disc for measuring depth. This should **only** occur when the vegetation is too dense to successfully get the Secchi disc to the bottom substrate.

Secchi disc depth—While the disc is on the bottom of the wetland look to see if it is visible. If it is, the Secchi disc depth should be recorded the same as the water depth. If the disc is not visible, slowly raise it until it is visible. Move the disc up and down to determine the point at which it disappears from view. Record the depth at which that occurs ( $\pm 1$  cm) as the Secchi disc depth. **If the site is dry, please enter a Secchi disc depth of 0. If you entered ‘Other’ for how depth was measured we assume you will not be able to record a Secchi disc depth.**

Temperature, pH, salinity, and specific conductivity—Using a water quality meter (e.g., YSI Pro 1030) with the probe at the mid-point of the water column, record water temperature ( $\pm 0.1^\circ\text{C}$ ), pH, salinity ( $\pm 0.1$  ppt), and specific conductivity ( $\pm 1$   $\mu\text{S cm}^{-1}$ ). Ensure that the entire length of the probe is submerged, as the conductivity sensors are located near the top. If the water depth at a site is less than the length of the probe, but greater than 5 cm, water quality should still be measured. This can be done by situating the probe in the water horizontally. At some tall emergent sites, residual vegetation may need to be parted in order to submerge the probe in the water.

Soil texture—Soil texture by feel will be recorded for each plot. Use a soil corer to collect soil (~25 grams, slightly larger than a ping-pong ball) from the top 4 inches of the substrate. For deeper SAV sites, an Ekman dredge in combination with a canoe paddle can be used to obtain a soil sample. Use the flow chart in Figure 6 and/or the soil textural triangle in Figure 7 to determine the soil texture and record the appropriate code from Table 3 on the datasheet.

Soil chemical and physical characteristics—**Optional for 2015.** Soil samples will be collected, air-dried within 24 hours and stored in a cool dark place until they can be sent to a lab for analyses (see Appendix IV). No funding from I&M is available, so if stations have funding available and are interested in analysis of soil samples, see Appendix IV for variables and estimated cost (based on prices from Utah State University Analytic Lab).


Soils should be collected from the first 10 (or 20) GRTS points in each habitat type (e.g., tall emergent, SAV). Approximately 2 cups of dry soil will be needed for a lab to analyze all the



parameters in Appendix IV. It is important to note that depending on substrate saturation levels, a greater amount of soil is often required from the field in order to yield 2 cups post air-drying. Some of the open water sites at Bear Lake NWR contained highly saturated loamy soils that required collection of up to a 1 gallon bag of substrate in order to acquire the required amount of dry sample. Soil from the top 6 inches of the core should be placed in a plastic bag. After sampling, immediately take the soils to an area (no direct sunlight) where they can be spread out on a plastic tray to air dry. In 2014, the SE Idaho crew used paint strainers arranged on a corrosion-free metal grate table. Fans can be used to increase air circulation drying speed, but ensure that fans do not blow away fine soils or light organic matter. If soils are collected at a plot, circle “yes” for soil sample collected on the vegetation data sheet.


*Soil moisture and temperature*—**Optional for 2015.** For plots with no standing water, measure soil moisture and temperature at two locations in the soil profile: 1) in the top 3 cm of the soil surface and 2) at 12 inches below the soil surface. Measure soil temperature to the nearest 1 °C using with a soil thermometer (e.g., [http://www.forestry-suppliers.com/product\\_pages/Products.asp?mi=64641&title=Soil+Thermometer&itemnum=89027](http://www.forestry-suppliers.com/product_pages/Products.asp?mi=64641&title=Soil+Thermometer&itemnum=89027)).

Due to the high clay and organic matter likely present in most sampling unit, volumetric water content will be calculated from permittivity values measured by a soil moisture meter such as the Campbell Scientific H2S (<http://www.campbellsci.com/hs2-overview>). For each sampling unit where soil moisture is collected, a linear relationship between permittivity and volumetric water content will have to be developed. We are currently working on this methodology. If you are interested in collecting data on soil moisture during 2015, contact Adonia Henry ([adoniarhenry@gmail.com](mailto:adoniarhenry@gmail.com)).



**Figure 6.** Flow chart for estimating soil texture by feel. Note that most soils collected during the vegetation sampling will be wet. If the soil is too wet and puddles, let the water drain off. There is no silt in this diagram but pure silt is uncommon and the difference between silt and silt loam is inconsequential in most routine wetland work (adapted from Thein 1979 by Richardson and Vepraskas 2001). Soil codes from Table 3 included in red.

## Soil Textural Triangle



**Figure 7.** The USDA Soil Textural Triangle.

**Table 3.** Soil texture classes.

Class	Code	Characteristics <sup>a</sup>
<b>Organic</b> (peat, mucky peat, or muck)	Org	Does not feel gritty or sticky, is easily compressed, has no internal strength, almost always black
<b>Rhizomes</b>	Rhiz	Dense mat of rhizomes > 2 feet deep with no other discernable substrate type present
<b>Mineral</b>		Feel gritty or sticky, resists compression, usually brown, red, yellow, or gray
Sand	Sa	Does not remain in a ball when squeezed
Loamy sand	LoSa	Remains in a ball when squeezed but does not form a ribbon
Sandy loam	SaLo	Makes a weak ribbon and feels gritty
Silt loam	SiLo	Makes a weak ribbon and feels very smooth
Loam	Lo	Makes a weak ribbon, neither grittiness nor smoothness dominant
Sandy clay loam	SaClLo	Makes a medium ribbon and feels gritty
Silty clay loam	SiClLo	Makes a medium ribbon and feels smooth
Clay loam	ClLo	Makes a medium ribbon, neither grittiness nor smoothness dominant
Sandy clay	SaCl	Makes a long ribbon and feels gritty
Silty clay	SiCl	Makes a long ribbon and feels very smooth
Clay	Cl	Makes a long ribbon, neither grittiness nor smoothness dominant

<sup>a</sup>See Figure 6 for detailed characteristics of measuring soil texture in the field

### **Wetland Hydrology**

*Depth and duration of flooding*—A staff gauge or water level data-logger should be installed in each sampling unit if not already present. Water levels should be recorded at least weekly following ice-out through the end of October. **Water levels at the staff gauge (or data-logger) should also be recorded on each day vegetation sampling is completed within that unit.** For guidance on installation of staff gauges, see Appendix IV.

## CHAPTER 3

### DATA MANAGEMENT

#### **Data Forms and Data Storage:**

Most of the data you collect will be recorded on data forms (Appendix I). Please record all data in pencil. Complete data forms following the instructions provided. If you are unsure about how you should record something, spell it out in sufficient detail so that there can be no doubt about what you mean. Then contact one of the project leads for an explanation as soon as possible.

A database will be provided to you for entering data collected in the field.

**PLEASE SEND 1) ELECTRONIC AND 2) HARD COPIES OF YOUR DATA FORMS TO JEFF AT THE END OF THE SEASON.**

### LITERATURE CITED

- Bennetts, R. E., J. E. Gross, K. Cahill, C. McIntyre, B. B. Bingham, A. Hubbard, L. Cameron, and S. Carter. 2007. Linking monitoring to management and planning: assessment points as a generalized approach. *George Wright Forum* 24:59-77.
- Bestelmeyer, B. T., K. Moseley, P. L. Shaver, H. Sanchez, D. D. Briske, and M. E. Fernandez-Gimenez. 2010. Practical Guidance for Developing State-and-Transition Models. *Rangelands* 32:23-30.
- Busch, D. E. and J. C. Trexler (eds). 2003. *Monitoring ecosystems*. Island Press, Washington D.C., USA. 447 pp.
- Clements, F. E. 1936. Nature and structure of the climax. *Journal of Ecology* 24:252-284.
- Deppe, E. R., and R. C. Lathrop. 1992. A comparison of two rake sampling techniques for sampling aquatic macrophytes. *Wisconsin Department of Natural Resources Research Management Findings* 32.
- Drake, J. A. 1990. The mechanics of community assembly and succession. *Journal of Theoretical Biology* 147:213- 233.
- Euliss, N., J. LaBaugh, L. Fredrickson, D. Mushet, M. Laubhan, G. Swanson, T. Winter, D. Rosenberry, and R. Nelson. 2004. The wetland continuum: A conceptual framework for interpreting biological studies. *Wetlands* 24:448-458.
- Huggett, A. J. 2005. The concept and utility of 'ecological thresholds' in biodiversity conservation. *Biological Conservation* 124:301-310.
- Jessen, R., and R. Lound. 1962. An evaluation of a survey technique for submerged aquatic plants. *Minnesota Department of Conservation Game Investigation Report No. 6*.
- Kenkel, N. C., and J. Podani. 1991. Plot Size and Estimation Efficiency in Plant Community Studies. *Journal of Vegetation Science* 2:539-544.
- Kindcaid, T and Olsen, T. 2013 spsurvey: Spatial survey design and analysis, R package version 2.6 <http://www.epa.gov/nheerl/arm/>
- Martin, J., M. C. Runge, J. D. Nichols, B. C. Lubow, and W. L. Kendall. 2009. Structured decision making as a conceptual framework to identify thresholds for conservation and management. *Ecological Applications* 19:1079-1090.
- Mitsch, W. J., and J. G. Gosselink. 2007. *Wetlands*. John Wiley and Sons, Inc. Hoboken, NJ.



- Murkin, H. R., E. J. Murkin, and J. P. Ball. 1997. Avian habitat selection and prairie wetland dynamics: a 10-year experiment. *Ecological Applications* 7:1144-1159.
- Ratti, J. T., and J. A. Kadlec. 1992. Concept plan for the preservation of wetland habitat of the Intermountain West. Unpublished report. Department of the Interior, U. S. Fish and Wildlife Service, Region 1, Portland, OR.
- Richardson, J. L. and M. J. Vepraskas. 2001. Wetland soils genesis, hydrology, landscapes, and classification. Lewis Publishers, CRC Press LLC, Boca Raton, FL.
- Smith, S. D. P. 2012. Identifying and evaluating causes of alternative community states in wetland plant communities. *Oikos* 121:675-686.
- Stringham, T. K., W. C. Krueger, and P. L. Shaver. 2003. State and transition modeling: an ecological process approach. *Journal of Range Management* 56:106-113.
- Thein, S. J. 1979. A flow diagram for teaching texture-by-feel analysis. *Journal of Agronomic Education* 8:54-55.
- van der Valk, A. G. 1981. Succession in wetlands: a Gleasonian approach. *Ecology* 62:688-696.
- Westoby, M. 1980. Elements of a theory of vegetation dynamics in arid rangelands. *Israel Journal of Botany* 28:169-194.
- Yin, Y., J. S. Winkelman, and H. A. Langrehr. 2000. Longterm resource monitoring program procedures: aquatic vegetation monitoring. Pages 22 *in* U. S. Geological Survey, LTRMP95-P002-7, La Crosse, Wisconsin.
- Zweig, C. L., and W. M. Kitchens. 2009. Multi-state succession in wetlands: a novel use of state and transition models. *Ecology* 90:1900-1909.

**APPENDIX I**  
**DATA FORMS**

Date \_\_\_\_/\_\_\_\_/\_\_\_\_ (mm/dd/yy) Observer \_\_\_\_\_ Recorder \_\_\_\_\_ Time: Start \_\_\_\_\_ End \_\_\_\_\_  
 Station: \_\_\_\_\_ Unit Name \_\_\_\_\_ Water Level \_\_\_\_\_

SAV/TALL EMERGENT WETLAND VEGETATION SAMPLING DATA SHEET  
**OCULAR SURVEY**

Point (SiteID) \_\_\_\_\_

EvalStatus: TS NT IN Turbid NotEval (circle)

Survey Method: B F (circle)

EvalReason: (if IN or NT) UPL TF SF PF IN\* OTHER\* (circle)

Explain\* (IN or Other): \_\_\_\_\_

OCULAR VEGETATION SURVEY (1m x 1m plot)						ABIOTIC
Spp Code	% Cover	Comments	Spp Code	% Cover	Comments	Depth (cm) <sup>c</sup>
BASU						Water Depth _____
RESID <sup>b</sup> -						Secchi Depth _____
						Secchi Other (circle)
						Water Chemistry
						pH (units) _____
						Sp. Cond (µs/cm) _____
						Salinity (ppt) _____
						Temp (°C) _____
						Soil Characteristics
						Texture _____
						(texture by feel)
						Soil Sample Collected?
						Y N (circle)
Total Cover			Total Cover			

Comments<sup>a</sup>: \_\_\_\_\_

Point (SiteID) \_\_\_\_\_

EvalStatus: TS NT IN Turbid NotEval (circle)

Survey Method: B F (circle)

EvalReason: (if IN or NT) UPL TF SF PF IN\* OTHER\* (circle)

Explain\* (IN or Other): \_\_\_\_\_

OCULAR VEGETATION SURVEY (1m x 1m plot)						ABIOTIC
Spp Code	% Cover	Comments	Spp Code	% Cover	Comments	Depth (cm) <sup>c</sup>
BASU						Water Depth _____
RESID <sup>b</sup> -						Secchi Depth _____
						Secchi Other (circle)
						Water Chemistry
						pH (units) _____
						Sp. Cond (µs/cm) _____
						Salinity (ppt) _____
						Temp (°C) _____
						Soil Characteristics
						Texture _____
						(texture by feel)
						Soil Sample Collected?
						Y N (circle)
Total Cover			Total Cover			

Comments<sup>a</sup>: \_\_\_\_\_

<sup>a</sup>For sites with that require measurements from more than one stratum (e.g. FILAL, waterlily, aquatic morph smartweed), please enter "surface" stata (e.g. FILAL - surface, FILAL - water column, waterlily - above) data in the site comments section

<sup>b</sup>Identify to category: 1 - tall emergent 2 - short emergent 3 - submerged

<sup>c</sup>NA ≠ 0

### RAKE AND OCCULAR SURVEY

**EvalReason:** (if IN or NT) UPL TF SF PF IN\* OTHER\* (circle)

Explain\* (IN or Other):

OCULAR VEGETATION SURVEY (1m x 1m plot)						ABIOTIC
Spp Code	% Cover	Comments	Spp Code	% Cover	Comments	Depth (cm) <sup>c</sup>
BASU						Water Depth _____
RESID <sup>b</sup> -						Secchi Depth _____
						Secchi    Other (circle)
						<b>Water Chemistry</b>
						pH (units) _____
						Sp. Cond (μs/cm) _____
						Salinity (ppt) _____
						Temp (°C) _____
						<b>Soil Characteristics</b>
						Texture _____
						(texture by feel)
						<b>Soil Sample Collected?</b>
Total Cover			Total Cover			Y    N (circle)
Comments <sup>a</sup> :						

RAKE VEGETATION SURVEY (1m × 0.35m sub-plots)						Rake Density Categories	
R1		R2		R3			
Spp Code	Density	Spp Code	Density	Spp Code	Density	Density	%Filled
<i>Total</i>		<i>Total</i>		<i>Total</i>		1	1% (single plant)
						2	1–20%
						3	21–40%
						4	41–60%
						5	61–80%
						6	81–100%
						7	>100%
Comments <sup>a</sup> :							

<sup>a</sup>For sites with that require measurements from more than one stratum (e.g. FILAL, waterlily, aquatic morph smartweed), please enter "surface" stata (e.g. FILAL - surface, FILAL - water column, waterlily - above) data in the site comments section

<sup>b</sup>identify to category:    1 - tall emergent    2 - short emergent    3 - submerged

 $c_{NA} \neq 0$

## APPENDIX II


### HOW TO BUILD A 1M × 1M PVC PLOT FRAME

#### Supplies

¾ inch PVC pipe (cut to desired lengths)	PVC primer
4 PVC elbows	PVC cement
2 or 4 PVC in-line couplers	Silicone or expansive foam
4 eyebolts (¼ x 1 ½ inch)	Sharpie or electrical tape (cut thin)
2 wing nuts	Bright-colored paracord
2 nylon locking washers	Drill and hacksaw

Cut and assemble plot frame using PVC pipe, elbows, and in-line couplers as shown below. The plot frame can be made with 0, 2, or 4 in-line couplers, depending on how small you want it to be able to break down for easy transport. We suggest that it should at least split in half (e.g., using 2 couplers) to facilitate putting the frame over tall emergent vegetation. Using 4 couplers will allow it to break down further, but the more joints that are not glued, the less sturdy it will be in the field. Seal inside of PVC pipe, couplers, and elbows with silicone or expansive foam to ensure water does not get into the PVC. PVC joints should be glued with PVC primer and cement (follow instructions on the cans). All elbow joints should be glued!!

Note: Larger diameter PVC may be used to build a sturdier frame. Ensure that the eye-bolts are long enough to go through the size PVC you use.



Characteristics of all plot frames (regardless of the number of glued joints):

1. Seal inside of PVC pipe, couplers, and elbows with silicone or expansive foam to ensure water does not get into the PVC.
2. **Interior** dimensions of 1 meter x 1 meter. PVC elbows will overlap 1 inch of the PVC pipe, but may not be exactly the same. THEREFORE IT IS NECESSARY TO MEASURE WITH THE PARTS YOU ARE USING.
3. Marks every 10 cm along the frame (measured along the inside of the frame) to facilitate accurate and consistent estimates of percent cover.
4. One end of bright-colored paracord should be permanently tied to the eye bolt with the nylon locking nut. The other end should be attached to a wing nut that can be secured onto the eyebolt on the opposite side of the frame once the frame is placed at sample points in tall emergent and floating leaved vegetation. Note: both ends of the paracord can stay permanently attached to the frame when in SAV/open water.



## APPENDIX III

### CALIBRATING A YSI 63 PORTABLE WATER METER

#### 4.2 pH Calibration

The YSI Model 63 *MUST* be calibrated before making pH measurements. Calibration may be performed at 1, 2 or 3-points (at pH 7, 4 and 10, or at pH 6.86, 4.01 and 9.18). Perform a 1-point calibration (at pH 7 or at pH 6.86) *ONLY* if a previous 2 or 3-point calibration has been performed recently. In most cases, a 2-point pH calibration will be sufficient for accurate pH measurements, but if the general range of pH in the sample is not known, a 3-point calibration may be necessary. 3-point calibration assures accurate pH readings regardless of the pH value of the sample. See *9.1 pH* for more details.

**WARNING: Calibration reagents may be hazardous to your health. Refer to *Appendix B - Health and Safety* for more information.**

Before calibrating the YSI Model 63, complete the procedures discussed in the *Preparing the Meter* and *Preparing the Probe* chapters of this manual.

The user can choose from two sets of pH buffer values for 3-point calibration. The first set consists of the standard YSI pH buffer values of pH 7 (YSI 3822), pH 4 (YSI 3821) and pH 10 (YSI 3823). The second set available is the NIST pH 6.86, 4.01 and 9.18. **Note that the first calibration point must be either pH 7 or pH 6.86.** Calibration is performed as follows:

1. Turn the instrument on by pressing the **ON/OFF** key. Press the **MODE** key until pH is displayed.
2. Rinse the probe with deionized or distilled water, then carefully dry the probe (or rinse it with some of the pH buffer solution to be used for calibration).
3. Place 30 to 35 mL of the pH buffer you have chosen to calibrate the system with (pH 7 or 6.86) in the 100 mL graduated cylinder. The graduated cylinder minimizes the amount of solution needed. Immerse the probe making sure that both the pH and temperature sensors are covered by the solution (see *Figure 5* on the following page).

**For best results:**

- Calibrate as close as possible to the sample temperature.
- After storage in pH 4 buffer/KCl solution, place the pH sensor in pH 7 (6.86) buffer and allow to acclimate before calibrating (5 to 10 minutes).
- Always give the pH and temperature sensors enough time to equilibrate with the temperature of the buffer.





Figure 5


4. To enter the calibration menu, use two fingers to press and release both the **UP ARROW** and **DOWN ARROW** keys at the same time. The Model 63 display will show **CAL** at the bottom, **STAND** will be flashing and the pH reading will show **7.00** (the buffer to be used to adjust the offset).



NOTE: If you will be calibrating with pH buffers of 6.86, 4.01 and 9.18 (instead of 7, 4 and 10), press both the **UP ARROW** and **DOWN ARROW** keys again. The display will change to 6.86.

NOTE: The Model 63 automatically accounts for the fact that the true pH of the buffers changes with temperature, therefore, the pH values displayed during calibration will vary with temperature. For example, pH 7 buffer at 20°C (rather than 25°C) has an actual pH of 7.02 and this number (rather than 7.00) will appear on the display when the probe is placed in the solution. See *Appendix C - pH Buffer Values*.


5. Press the **ENTER** key. The Model 63 display will show **CAL** at the bottom, **STAND** will stop flashing and the pH calibration value is shown with the middle decimal point flashing.



6. When the reading is stable (does not change by 0.01 pH in 10 seconds), the decimal point will stop flashing. Press and hold the **ENTER** key to save the calibration point. The Model 63 will flash **SAVE** on the display along with **OFS** to indicate that the offset value has been saved.



7. **SLOPE** will now appear on the display and be flashing. This indicates that the slope is ready to be set using a second pH buffer. The system is now calibrated at a single point. If you are only performing a single point calibration, press the **MODE** key to return to normal operation.




8. Rinse the probe with deionized or distilled water, then carefully dry the probe.


**STOP HERE IF PERFORMING A 1-POINT CALIBRATION.**

9. If you are performing a 2-point or 3-point calibration, fill a clean container with the second value pH buffer (pH 4 or 10, or pH 4.01 or 9.18) and immerse the probe into the solution. Make sure that the temperature sensor is immersed.

10. Press the **ENTER** key. The Model 63 should now show **CAL** at the bottom, **SLOPE** will stop flashing and the pH calibration value (automatically sensed by the instrument) is shown with one of the decimal points flashing.




- If the second pH buffer is less than the first buffer (which was used to adjust the offset; pH 7 or pH 6.86), the left decimal point will flash as shown above. If the second pH buffer is greater than the first, the right decimal point will flash as shown below.



11. When the reading is stable (does not change by 0.01 pH in 10 seconds), the decimal point will stop flashing. Press and hold the **ENTER** key to save the first **SLOPE**. The Model 63 will flash **SAVE** on the display along with SLP to indicate that the first slope value has been saved.



12. **SLOPE** will start flashing again indicating that the slope is ready to be set using a third pH buffer.




13. The system is now calibrated at two points. If you are only performing a two point calibration, press the **MODE** key to return to normal operation.
14. Rinse the probe with deionized or distilled water, then carefully dry the probe.

**STOP HERE IF PERFORMING A 2-POINT CALIBRATION.**


15. If you are performing a 3-point calibration, fill a clean container with the third value pH buffer (pH 4 or 10, or pH 4.01 or 9.18) and immerse the probe into the solution. Make sure that the temperature sensor is immersed.

NOTE: The third buffer must not be the same as the second buffer. For example; if the second buffer was less than pH 7, the third buffer must be greater than pH 7.

16. Press the **ENTER** key. The Model 63 display will now show **CAL** at the bottom, **SLOPE** will stop flashing and the pH calibration value (automatically sensed by the instrument) is shown with one of the decimal points flashing. If the third pH buffer is less than the first buffer (which was used to adjust the offset; usually pH 7), the left decimal point will flash. If the third pH buffer is greater than the first, the right decimal point will flash.



or



17. When the reading is stable (does not change by 0.01 pH in 10 seconds), the decimal point will stop flashing. Press and hold the **ENTER** key to save the second SLOPE. The Model 63 will flash **SAVE** on the display along with **SLP** to indicate that the second slope value has been saved.



The system is now calibrated at three points and will return to normal operation.

18. Rinse the probe with deionized or distilled water.

### 4.3 Conductivity Calibration

**IMPORTANT:** System calibration is rarely required because of the factory calibration of the YSI Model 63. However, from time to time it is wise to check the system calibration and make adjustments when necessary.

**Prior to calibration of the YSI Model 63, it is important to remember the following:**

1. Always use clean, properly stored, NIST traceable calibration solutions (see *12 Accessories and Replacement Parts*). When filling a calibration container prior to performing the calibration procedures, make certain that the level of calibrant buffers is high enough in the container to cover the entire probe. Gently agitate the probe to remove any bubbles in the conductivity cell.
2. Rinse the probe with distilled water (and wipe dry) between changes of calibration solutions.
3. During calibration, allow the probe time to stabilize with regard to temperature (approximately 60 seconds) before proceeding with the calibration process. The readings after calibration are only as good as the calibration itself.
4. Perform conductivity calibration at a temperature as close to 25°C as possible. This will minimize any temperature compensation error.


**Follow these steps to perform an accurate calibration of the YSI Model 63:**

1. Turn the instrument on and allow it to complete its self test procedure.
2. Select a calibration solution which is most similar to the sample you will be measuring.
  - For sea water choose a 50 mS/cm conductivity standard (YSI Catalog# 3169)
  - For fresh water choose a 1 mS/cm conductivity standard (YSI Catalog# 3167)
  - For brackish water choose a 10 mS/cm conductivity standard (YSI Catalog # 3168)
3. Place at least 7 inches of solution in the plastic container or a clean glass beaker.

**NOTE: Do NOT use the 100 mL graduated cylinder.** The diameter of the cylinder is too small for accurate conductivity measurements.

4. Use the **MODE** key to advance the instrument to display conductivity.
5. Insert the probe into the solution deep enough to completely cover the probe. Both conductivity ports must be submerged (see *Figure 6* on the following page).
6. Allow at least 60 seconds for the temperature reading to become stable.
7. Move the probe vigorously from side to side to dislodge any air bubbles from the electrodes.
8. Press and release the **UP ARROW** and **DOWN ARROW** keys at the same time.

The **CAL** symbol will appear at the bottom left of the display to indicate that the instrument is now in Calibration mode.



9. Use the **UP ARROW** or **DOWN ARROW** key to adjust the reading on the display until it matches the value of the calibration solution you are using.
10. Once the display reads the exact value of the calibration solution being used (the instrument will make the appropriate compensation for temperature variation from 25°C), press the **ENTER** key. The word “**SAVE**” will flash across the display for a second indicating that the calibration has been accepted.

The YSI Model 63 is designed to retain its last conductivity calibration permanently. Therefore, there is no need to calibrate the instrument after battery changes or power down.



## APPENDIX IV

## DESCRIPTION OF WETLAND SOIL ANALYSIS

**Rationale**

The importance of substrate in determining the distribution of aquatic plants has been known since the early 1900s (i.e., Pearsall 1920, Misra 1938). Barko et al. (1986) list water temperature and sediment composition as 2 of the 4 most prevalent environmental factors affecting submergent vegetation communities. Soil properties that affect wetland vegetation dynamics include soil organic matter, texture, cation exchange capacity, pH, salinity, and nutrients (Sand-Jensen and Sondergaard 1979, Barko and Smart 1983, Smith and Kadlec 1983, Smith and Kadlec 1985, Barko and Smart 1986, Dunn and Scott 1987, Jackson and Charles 1988, Jackson et al. 1993, Madsen et al. 1993, Bini et al. 1999, Stolt et al. 2000). Individual and interrelated parameters of water chemistry also influence aquatic vegetation (Hutchinson 1975). Within a wetland, spatial variability in conductivity, pH, and soil parameters (pH, organic matter, CEC, and texture) may be associated with the distribution of different growth forms of aquatic vegetation (i.e., floating-leaved and submergent; Frodge et al. 1990, Reese and Moorhead 1996, Khedr and El-Demerdash 1997).

**Methods and Cost**

Soil core samples will be collected at a subsample of GRTS plots in each priority sample unit at participating stations. Soil samples will be placed in a Ziploc bag, labeled with the unit and plot number. When returning from the field, soil samples will be air-dried. Once dry, samples will be ground to pass through a 2 mm sieve, bagged, and sent to Utah State University Analytical Lab for analyses. Soil samples will be analyzed for the following characteristics (prices based on USU price list).

<b>Test</b>	<b>Description</b>	<b>Price (ea)</b>
<i>Basic</i>		
S6	pH, ECe (electrical conductivity, and SAR (sodium adsorption ratio)	\$14.00
S14a	Carbon (organic matter) and Total N	\$11.50
S23	Particle size by hydrometer	\$17.00
S17b	Cation exchange capacity	\$11.50
<b><i>Subtotal Basic</i></b>		<b>\$ 54.00</b>
<i>Nutrients</i>		
S7c	Phosphorus (available P) and Potassium	\$8.50
S8d	Ammonia N and Nitrate N	\$17.00
<b><i>Subtotal Nutrients</i></b>		<b>\$25.50</b>
<b>Total (per sample)</b>		<b>\$79.50</b>

A minimum of 10 soil core samples per aquatic plant growth form (e.g., tall emergent, floating-leaved, submerged aquatic) within a sampling unit should be collected. If funding is available, 20 samples per aquatic growth form are desired in order to analyze abiotic variables associated with individual vegetation species in addition to growth form. Cost scenarios for a unit with two aquatic plant growth forms and three priority units sampled at each station are listed below:

#Samples SAV	#Samples Tall Emergent	Qty	Price	Total Per Unit	#Units	Total PerStation
10	10	20	\$79.50	\$1,590.00	3	\$4,770.00
20	20	40	\$79.50	\$3,180.00	3	\$9,540.00

Other soil characteristics that will be measured in the field (or at the station after field collection) include: soil temperature, texture by feel, soil moisture (if sample plot has no standing water), and bulk density (see IWWWG Protocol Manual for methods).

#### LITERATURE CITED

- Barko, J. W., M. S. Adams, and N. L. Clesceri. 1986. Environmental factors and their consideration in the management of submersed aquatic vegetation: a review. *Journal of Aquatic Plant Management* 24:1-10.
- Barko, J. W. and R. M. Smart. 1983. Effects of organic matter additions to sediment on the growth of aquatic plants. *Journal of Ecology* 71:161-175.
- . 1986. Sediment-related mechanisms of growth limitation in submersed macrophytes. *Ecology* 67:1328-1340.
- Bini, L. M., S. M. Thomaz, K. J. Murphy, and A. F. M. Camargo. 1999. Aquatic macrophyte distribution in relation to water and sediment conditions in the Itaipu Reservoir, Brazil. *Hydrobiologia* 415:147-154.
- Dunn, C. P. and M. L. Scott. 1987. Response of wetland herbaceous communities to gradients of light and substrate following disturbance by thermal pollution. *Vegetatio* 70:119-124.
- Frodge, J. D., G. L. Thomas, and G. B. Pauley. 1990. Effects of canopy formation by floating and submergent aquatic macrophytes on the water quality of two shallow Pacific Northwest lakes. *Aquatic Botany* 38:231-248.
- Hutchinson, G. E. 1975. *A treatise on limnology*. John Wiley & Sons, New York, New York, USA. 660pp.
- Jackson, L. J., J. Kalff, and J. B. Rasmussen. 1993. Sediment pH and redox potential affect the bioavailability of Al, Cu, Fe, Mn, and Zn to rooted aquatic macrophytes. *Canadian Journal of Fisheries and Aquatic Sciences* 50:143-148.
- Jackson, S. T. and D. F. Charles. 1988. Aquatic macrophytes in Adirondack (New York) lakes: patterns of species composition in relation to environment. *Canadian Journal of Botany* 66:1449-1460.

- Khedr, A. H. A. and M. A. El-Demerdash. 1997. Distribution of aquatic plants in relation to environmental factors in the Nile Delta. *Aquatic Botany* 56:75-86.
- Madsen, J. D., R. M. Smart, L. W. Eichler, C. W. Boylen, J. W. Sutherland, and J. A. Bloomfield. 1993. Submersed littoral vegetation distribution: field quantification and experimental analysis of sediment types from Onondaga Lake, New York. *Aquatic Plant Control Research Program, Waterways Experiment Station, U. S. Army Corps of Engineers, Vicksburg, Mississippi, USA.* 50pp.
- Misra, R. D. 1938. Edaphic factors in the distribution of aquatic plants in the English Lakes. *Journal of Ecology* 26:411-451.
- Pearsall, W. H. 1920. The aquatic vegetation of the English Lakes. *Journal of Ecology* 8:163-201.
- Reese, R. E. and K. K. Moorhead. 1996. Spatial characteristics of soil properties along an elevational gradient in a Carolina bay wetland. *Soil Science Society of America Journal* 60:1273-1277.
- Sand-Jensen, K. and M. Sondergaard. 1979. Distribution and quantitative development of aquatic macrophytes in relation to sediment characteristics in oligotrophic Lake Kalgaard, Denmark. *Freshwater biology* 9:1-11.
- Smith, L. M. and J. A. Kadlec. 1983. Seed banks and their role during drawdown of a North American marsh. *Journal of Applied Ecology* 20:673-684.
- . 1985. Predictions of vegetation change following fire in a Great Salt Lake Marsh. *Aquatic Botany* 21:43-51.
- Stolt, M. H., M. H. Genthner, W. L. Daniels, V. A. Groover, S. Nagle, and K. C. Haering. 2000. Comparison of soil and other environmental conditions in constructed and adjacent palustrine reference wetlands. *Wetlands* 20:671-683.

## APPENDIX V

### DOUBLE-OBSERVER SAMPLING

We will continue to quantify variation among observers in 2015 at pre-selected stations/units. **Contact Jeff Warren if you are interested in assisting with double-observer efforts this year.**

*Double-observer Sampling*—The repeatability of our field methods will be assessed by estimating the amount of variation in ocular estimates among observers. We will use a double-observer approach to sample a subset of units for this purpose. The standard procedure is for one observer and one recorder when sampling wetland points. The double-observer method requires both individuals to record an estimate of canopy cover for each point. An observer's estimates must be independent from the second observer, therefore it is imperative that the observers **do not** discuss their estimates until both have recorded them. Also, observers **should not change** their estimates after discussing it with the other observer. The exception to this is for plant identification – it is allowable (recommended, in fact) for the two observers to discuss plant identification. This will eliminate variation due to detection probability and identification between observers, a source of variation we are not interested in at this time.

Stations participating in double-observer sampling **do not** need to conduct double-observer sampling on all GRTS points at their station. A general recommendation is to conduct double-observer sampling on  $\frac{1}{4}$  of the GRTS points as an ordered subset. For example, if a station was going to sample a total of 40 GRTS points, they would conduct double-observer sampling on GRTS points 1–10 and do the standard survey on the remaining points. In this example, if any points between 1–10 are dropped from the sample the double-observer subset can simply be extended using the PanelOne points.

Conducting double-observer sampling is most efficient for those stations or crews where multiple individuals will be trained in plant identification and expected to conduct vegetation surveys. For a station where a single individual will be the primary observer it is not likely worthwhile to attempt double-observer sampling. **Do not conduct double-observer points at different times** – that introduces spatial error associated with variation with GPS locations, something we are not interested in at this time.

When you enter double-observer observations please be sure to **1)** have the most experienced observer recorded as the 'primary observer', and **2)** check the 'Double-observer' box in the data entry form for the observations from the secondary observer. This allows us to query out the primary observer observations for standard analyses.

## APPENDIX VI

## GUIDE TO WETLAND INDICATOR SPECIES

## TARGET

<b>Semi-permanently Flooded Wetlands (SPF)</b>
<i>Surface water persists throughout the growing season in most years (CCGL79)</i>
<i>Deep Marsh: Maintain surface water throughout spring and summer, frequently into fall and winter, but in drought years often go dry by mid-summer (SK72)</i>
<i>For STM, the target also includes Intermittently Exposed where surface water is present throughout the year except during extreme drought (CCGL79)</i>

## Indicator Species

*If one or more of the emergent or SAV indicator species are present, regardless of cover (e.g., 1% or 100%) the plot should be sampled*

**Emergent Indicator Species**

	<b>Common Name</b>	<b>Reference</b>
Typha sp.	ALL cattail	SK72, CHHR72
Schoenoplectus heterochaetus	Slender bulrush	SK72
Bolboschoenus fluviatilis	River bulrush	SK72
Schoenoplectus acutus	Hardstem bulrush	SK72
Bolboschoenus maritimus	Alkali bulrush	SK72
Schoenoplectus tabernaemontani	Softstem bulrush	SK72
Schoenoplectus americanus	Olney bulrush	CHHR72
Phragmites australis	Common reed	SK72
Eleocharis acicularis (may be subm)	Needle spikerush	SK72

**SAV Indicator Species**

Stuckenia sp.	ALL narrow pondweeds	SK72
Potamogeton sp.	ALL pondweeds	SK72
Zannichelia palustris	Horned pondweed	SK72
Ruppia sp.	ALL wigeon grass	SK72, CHHR72
Callitriche sp.	ALL water-starwort	SK72
Hipparis vulgaris	Mare's tail	SK72
Ranunculus aquatilis	White water crowfoot	SK72
Utricularia sp.	ALL bladderwort	SK72, CHHR72
Myriophyllum sp.	ALL milfoil	SK72
Elodea sp.	ALL waterweed	SK72
Ceratophyllum demersum	Coontail	SK72, CHHR72
Najas sp.		CHHR72
Chara sp.	ALL muskgrass	SK72, CHHR72
Nitella sp.	ALL stoneworts	DeptEcol

## TARGET (con't)

Semi-permanently Flooded Wetlands (SPF)
<i>Surface water persists throughout the growing season in most years (CCGL79)</i>
<i>Deep Marsh: Maintain surface water throughout spring and summer, frequently into fall and winter, but in drought years often go dry by mid-summer (SK72)</i>
<i>For STM, the target also includes Intermittently Exposed where surface water is present throughout the year except during extreme drought (CCGL79)</i>

**Primary emergent shallow marsh species that can be SPF\***

*Shallow Marsh: Maintain surface water throughout spring and summer, but frequently dry during late summer and fall (SK72). NOTE: shallow marsh wetlands can be SPF or SF (CCGL79) and are often considered transitional between SPF and SF wetlands.*

**\*Because these species can occur in SPF or SF wetlands, they are not considered an indicator species. Therefore the decision as to whether or not to sample a plot with one or more of these species will depend on your knowledge of the flooding regime (e.g., water level management and/or NWI), and other species present. If one of more of these species covers 50% or more of the plot, the plot should be sampled. If these species represent a minor portion of the plot (< 50%) which is otherwise dominated by seasonally flooded species, it should not be sampled.**

Glyceria grandis	Tall mannagrass	SK72
Sparganium eurycarpum	Giant burreed	SK72, AKP03
Alisma triviale	Northern water plantain	SK72
Persicaria amphibia	Water smartweed	SK72
Beckmannia syzigachne	Slough grass	SK72
Scolochloa festucacea	River grass	SK72
Eleocharis palustris	Common spikerush	SK72, AKP03
Alisma gramineum	Narrowleaf waterplantain	SK72
Schoenoplectus pungens	Common threesquare	SK72
Puccinellia nuttalliana	Alkaligrass	SK72
Carex atherodes	Slough sedge	SK72
Carex utricularia	Northwest territory sedge	AKP03
Carex rostrata	Beaked sedge	WBLNO08

NOTE: If seed heads are not present, these sedge species are tall, robust sedges, greater than 2 ft and often up to 4 ft tall. This may be a useful characteristic to differentiate these species from those characteristic of seasonal or temporary wetlands.

Salicornia rubra	Red saltwort/swampfire	SK72
Distichlis spicata	Saltgrass	CHHR72

**TARGET (con't)**

<b>Semi-permanently Flooded Wetlands (SPF)</b>
<i>Surface water persists throughout the growing season in most years (CCGL79)</i>
<i>Deep Marsh: Maintain surface water throughout spring and summer, frequently into fall and winter, but in drought years often go dry by mid-summer (SK72)</i>
<i>For STM, the target also includes Intermittently Exposed where surface water is present throughout the year except during extreme drought (CCGL79)</i>

**Common Semi-permanent Drawdown Species**

*Annuals and/or bare substrate may occur during natural or managed drawdown phases of a SPF wetland and should be sampled.*

Chenopodium sp.

Hordeum jubatum

Eleocharis acicularis

Rumex sp.

Xanthium

Bidens sp.

Kochia sp.

Polygonum/Persicaria sp.

Echinochloa sp.

Kochia

Annual smartweeds

Millet

**NON-TARGET**

<b>Seasonally Flooded Wetlands (SF)</b>
<i>Surface water is present for extended periods especially early in the growing season, but is absent by the end of the season in most years (CCGL79)</i>
<i>Shallow marsh: Surface water is present for an extended period in spring and early summer, but frequently dry during late summer and fall (SK72)</i>

***Characteristic Species and Habitat Descriptions***

Typically dominated by grass and grass-like plants that are intermediate in height when compared to deep marsh and wet meadow/temporary wetlands (SK72)

***Grays Lake***

Juncus balticus	Baltic rush	AKP03
Mentha arvensis	Field mint	AKP03
Triglochin maritimum	Seaside arrowgrass	AKP03
Carex nebrascensis	Nebraska sedge	AKP03
Arnica chamissonis	Chamisso's arnica	AKP03
Argentina anserina	Silverweed cinquefoil	AKP03

*See also emergent shallow marsh species*



**NON-TARGET**

<b>Temporarily Flooded Wetlands (TF)</b>
<i>Surface water is present for brief periods during the growing season (CCGL79)</i>
<i>Wet meadow: Surface water usually persists for a few weeks during the spring and following precipitation events (SK72)</i>
<i>Wetland low prairie: surface water normally present in early spring for a few days or up to 2 weeks (SK72)</i>

***Characteristic Species and Habitat Descriptions***

Characterized by fine-texture grasses, rushes, and sedges of relatively low height with forbs common (SK72)

Characterized by plants that grow in uplands and wetlands (CCGL79).

***Grays Lake***

Deschampsia caespitosa	Tufted hairgrass	AKP03
Carex praegracilis	Clustered field sedge	AKP03
Aster campestris	Western aster	AKP03

***Prairie Pothole Region***

Poa palustris	fowl bluegrass	SK72
Carex praegracilis	Clustered field sedge	SK72
Symphyotrichum lanceolatum	lowland white aster	SK72
Calamagrostis stricta	Northern reedgrass	SK72
Spartina pectinata	Prairie cordgrass	SK72
Juncus balticus	Baltic rush	SK72
Hordeum jubatum	Foxtail barley	SK72
Distichlis spicata	Saltgrass	SK72

***Montane Wet Meadow Group (NVCS)***

Calamagrostis stricta	Northern reedgrass
Calamagrostis canadensis	Bluejoint reedgrass
Carex bolanderi	Bolander's sedge
Carex microptera	Small-wing sedge
Carex scopulorum	Mountain sedge
Deschampsia caespitosa	Tufted hairgrass
Juncus drummondii	Drummond's rush
Juncus nevadensis	Nevada rush
Camassia quamash	Camas
Rorippa alpina	Alpine yellowcress

*NOTE: Some of these species may also occur in seasonally flooded wetlands.*