PIPING PLOVER: 1989 SUMMARY

The observer network reported a total of 137 breeding pairs of Piping Plovers (Charadrius melodus) from 53 sites in Massachusetts in 1989 (Table 1), representing a second consecutive year of improvement. These results compare with 134 pairs in 1988 and 126 pairs in 1987, but remain below the 139 pairs estimated for 1986. Seventy-four sites were checked, with important sites-North Beach in Chatham and South Cape Beach in Mashpee-were not checked. Observer effort, as measured by the number of sites surveyed and the intensity of effort at each estimate reflects a true increase.

"Breeding pairs" are pairs that are observed with either a nest or unfledged chicks, or that exhibit site tenacity and evidence of pair bonding and territoriality. Single birds that are observed during the breeding season and exhibit agitated, site tenacious behavior or distraction displays when approached are generally considered indicative of a breeding pair at a site.

Breeding plovers were present at coastal sites from Plum Island south to the Rhode Island border and east to Cape Cod and Nantucket. Eighteen pairs (13%) occurred at sites on the North Shore (New Hampshire border to Boston); 7 pairs (5%) along the upper Cape (Boston to the Cape Cod Canal); 22 pairs (16%) on upper Cape Cod sites (Cape Cod Canal east to Brewster and Harwich); 34 pairs (25%) on lower Cape Cod sites (Chatham to Provincetown); 24 pairs (18%) on the west shore of Buzzards Bay outer islands (Elizabeths east to Nantucket). The North Shore Group showed the biggest overall gain owing to notable improvement at Crane Beach, Ipswich. The biggest overall drop occurred in the Buzzards Bay Group on account of predation problems at Little Beach, Dartmouth.

Mean number of pairs per site was 2.6 (range=i-14). Thirty-nine sites (74%) harbored 3 or fewer pairs, aggregating to 60 pairs (48% of the estimated population). Four sites harbored 34 distribution continues to dictate that management effort be directed at many different sites with relatively few pairs of plovers per site.

Overall mean productivity for the state was 1.59 chicks fledged (=reaching 25 days of age)/nesting pair, based on data reported for 123 breeding pairs (90% of state total) at 46 sites (87%). These results-up from 1.29 a year earlier and 1.10 in 1987--were certainly the best ever since statewide productivity estimates have been available. This favorable increase does seem

to be tied to very positive results obtained from the use of predator exclosure devices. Of 58 nests where devices were installed, 50 (86%) successfully hatched eggs!

We continue to be concerned that the intensity of observer effort mounted in recent years—both to census nesting beaches and to secure productivity data—may be unsustainable. Should this be the case, data quality would be compromised and year—to—year comparisons would be less meaningful. To address this concern, an attempt was made to assess the overall size of the breeding population in two "windows"—the first two weeks of May and the first two weeks of July. These indexes resulted in estimates of 98 and 105 pairs respectively. These numbers are interesting, but of dubious value, mainly owing to difficulty in quantifying observer effort and evenness of coverage.

Our thanks to all who contributed data and for your efforts during the past year. I welcome your comments and suggestions at any time.

Data compiled by: BRADFORD G. BLODGET, State Ornithologist, Massachusetts Division of Fisheries and Wildlife Field Headquarters, Route 135, Westboro, MA 01581.

Table 1. SUMMARY OF ABUNDANCE, DISTRIBUTION AND PRODUCTIVITY OF PIPING PLOVERS IN MASSACHUSETTS, 1989. An asterisk (*) denotes a fence surrounding a tern colony that functioned as a predator exclosure.

| Station Name | Numbe | er of B | Pairs | Chicks Fledged | Exclos N | sures NS |
|--|---------------|-------------|------------------|-------------------|----------------|------------------|
| NORTH SHORE GROUP | | | | nd | | |
| 2049 Crane Beach, Ipswich | 2 14 | 4 13 | 4 14 | 6 37 | 4 15 | 3 13 |
| SOUTH SHORE GROUP ACSO Third Cliff, Scituate ACSO Fourth Cliff, Scituate | 2 | 2 | 2 | 4 | 1* | 1 |
| 2052-Duxbury Beach, Duxbury S053 Plymouth Beach, Plymouth | 3 | 0 0 2 | 0 3 2 | 0 0 4 | - - - | - - - |
| BUZZARDS BAY GROUP Stony Point Dike, Wareham | 1 | 1 | 1 . | nd | _ | |
| ್ರಾಂತಿ Long Beach Point, Wareham | Ō | ō | Ō | 0 | | |
| West Island, Fairhaven | 0 | 0 | 0 | 0 | - | _ |
| <pre> Round Hill Beach, Dartmouth Salter's Pd:, Dartmouth </pre> | nd 1 | nd 1 | 1 1 | 1 | | |
| Little Beach, Dartmouth | 7 | 2 | 7 | 2 3 | 1 | 1 |
| 35 Barneys Joy, Dartmouth | 2 | 2 | 2 1 | Ö | - | _ |
| G/ Gooseberry Neck, Westport | 1 | 1 1 | 1 | 2 | , - | - . |
| 3 Horseneck Beach, Westport School Seach, Westport | 6 nd | 5 nd | 7 2 | 18 | - | - |
| Richmond Pd., Westport | nd | nd | 2 | 4 2 | - | _ |
| Bay Pt., Swansea | nd | nd | ō | Ö | - | - |
| UPPER CAPE GROUP | | | | • | | |
| 2060 Sippewisset, Falmouth | 0 | 0 | 0 | 0 | - | - |
| 68 South Cape Beach, Mashpee | 0 nd | 0 nd | • 0 nd | 0 nd | - | . - · |
| Popponesset, Mashpee | 2 | 1 | 2 | 2 | - 1* | 1 |
| 70. Town Neck Beach, Sand. | 1 | 0 | 1 | ō | _ | _ |
| o/ Springhill Beach, Sand. | 3 | 4 | 4 | nd | - | _ |
| <pre>7.2 E. Sandwich Beach, Sand. 7.5 Scorton Creek, Sandwich</pre> | 1 4 | 1 4 | 1 4 | 3 10 | 4 | 4 |
| Sandy Neck, Barnstable | 5 | 5 | 5 | 10 | 5 | 4 |
| 75 Dead NkSamp, Is., Barn. | 1 | 1 | 1 | Ō | 2 | i |
| 76 Long Beach, Barnstable | 1 | 1 | 1 | 1 | - | - |
| 77Kalmus Park, Barnstable 78Seagull Beach, Yarmouth | 2 . | 2 0 | 2 0 | 4 0 | 1* | 1 |
| ? Radio City, Yarmouth | ŏ | Ö | 0 | . 0 | _ _ | _ |
| C) West Dennis Beach, Dennis | 1 0 | 1 0 | 1 0 | 3 0 | 1* - | 1 - |
| LOWER CAPE GROUP | 4 | | | | | |
| 3082Hardings Beach, Chatham 3083 Hardings Bch. Pt., Chatham | 4 0 | 3 0 | 4 0 | 9 | - | - - |

Table 1. Mass. Piping Plovers, 1989 (cont.)

| | | | | | ** | |
|--|---|--|--|---|---|---|
| N. Monomoy Is., Chatham S. Monomoy Is., Chatham North Bch., Chatham North Bch. Is., Chatham Nauset Hts., Orleans New Is., Eastham/Orleans Coast Guard Bch., Eastham Marconi Bch., Wellfleet Jeremy Pt., Wellfleet Pamet R. mouth, Truro Pilgrim Bch., Provincetow High Head, Truro Race PtSouth, P'town Race PtNorth, P'town Wood End-Long Pt., P'town | 3 4 0 3 4 1 | nd 5 4 0 | 4 nd 5 4 0 | 0 7 nd 4 5 0 7 3 0 0 nd 2 6 3 0 | - 4 - 2 1* - 3 2 1 2 4 | -4 -2 1 -2 2 1 2 2 |
| OUTER ISLANDS GROUP Naushon Is., Gosnold Pasque Is., Gosnold 10 Nashawena Is., Gosnold 10 Cuttyhunk Is., Gosnold Martha's Vineyard Subgroup | 0 1 nd nd | 0 1 3 nd | 0 1 3 5 | 0 2 2 nd | - - - - | - - - - |
| Of Tashmoo Of Sylvia State Beach Of Harthaven Thong Beach Of Quansoo Of Norton's Pt. OloLobsterville Thattle Bch./Eel Pd. Cape Poge, Chappaquiddick Nantucket Subgroup: | 1 nd 0 0 0 nd 1 0 | 1 nd 0 1 1 1 0 0 | 1 0 2 2 1 1 0 1 | 2 nd 0 4 5 3 nd 0 2 | 1 - - 1 1 - - - 1 | 1 1 1 - 1 1 - 1 - 1 |
| Great Point The Galls Haulover Pd. Quidnet TSiasconset The Meach Tom Nevers Head Durfside Tuckernuck Is. Muskeget Is. | 3 0 0 0 0 0 0 1 1 2 1 | 6 2 1 0 0 0 0 0 1 2 nd | 6 2 1 0 0 0 0 1 1 2 | 8 1 0 0 0 0 0 1 4 nd | | |
| TOTALS: | 98 | 105 | 137 | 196 | 58 | 50 |
| _ | | | • | | | |

Data compiled by: BRADFORD G. BLODGET, State Ornithologist, Massachusetts Division of Fisheries and Wildlife, Field Headquarters, Rt. 135, Westboro, MA 01581.