

**U.S. FISH AND WILDLIFE SERVICE
SPECIES ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM**

SCIENTIFIC NAME: *Emballonura semicaudata semicaudata*

COMMON NAME: Pacific sheath-tailed bat (American Samoa Distinct Population Segment)

LEAD REGION: Region 1

INFORMATION CURRENT AS OF: March 2010

STATUS/ACTION

Species assessment - determined we do not have sufficient information on file to support a proposal to list the species and, therefore, it was not elevated to Candidate status

New candidate

Continuing candidate

Non-petitioned

Petitioned - Date petition received: March 4, 1986, and May 11, 2004

90-day positive - FR date: January 21, 1987

12-month warranted but precluded - FR date: May 11, 2005

Did the petition request a reclassification of a listed species?

FOR PETITIONED CANDIDATE SPECIES:

a. Is listing warranted (if yes, see summary of threats below)? Yes

b. To date, has publication of a proposal to list been precluded by other higher priority listing actions? Yes

c. If the answer to a. and b. is "yes", provide an explanation of why the action is precluded.

Higher priority listing actions, including court-approved settlements, court-ordered and statutory deadlines for petition findings and listing determinations, emergency listing determinations, and responses to litigation, continue to preclude the proposed and final listing rules for the species. We continue to monitor populations and will change its status or implement an emergency listing if necessary. The "Progress on Revising the Lists" section of the current CNOR (<http://endangered.fws.gov/>) provides information on listing actions taken during the last 12 months.

Listing priority change

Former LP:

New LP:

Date when the species first became a Candidate (as currently defined): January 21, 1987

Candidate removal: Former LPN:

A – Taxon is more abundant or widespread than previously believed or not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status.

U – Taxon not subject to the degree of threats sufficient to warrant issuance of a

proposed listing or continuance of candidate status due, in part or totally, to conservation efforts that remove or reduce the threats to the species.

- F – Range is no longer a U.S. territory.
- I – Insufficient information exists on biological vulnerability and threats to support listing.
- M – Taxon mistakenly included in past notice of review.
- N – Taxon does not meet the Act’s definition of “species.”
- X – Taxon believed to be extinct.

ANIMAL/PLANT GROUP AND FAMILY: Mammal: Family Emballonuridae (Sac-winged, Ghost, and Sheath-tailed bats)

HISTORICAL STATES/TERRITORIES/COUNTRIES OF OCCURRENCE: *Emballonura semicaudata semicaudata* occurred historically in American and Independent Samoa, Tonga, Fiji, and Vanuatu.

CURRENT STATES/COUNTIES/TERRITORIES/COUNTRIES OF OCCURRENCE: American and Independent Samoa, Tonga, Fiji, and Vanuatu.

LAND OWNERSHIP: The last known major roosting caves on Tutuila, American Samoa, are part of customary lands belonging to the village of Afono.

LEAD REGION CONTACT: Linda Belluomini, (503) 231-6283, linda_belluomini@fws.gov

LEAD FIELD OFFICE CONTACT: Pacific Islands Fish and Wildlife Office, Christa Russell, (808) 792-9400, christa_russell@fws.gov

BIOLOGICAL INFORMATION

Species Description

This small bat (forearm length c. 1.8 inches (45 millimeters), weight (.19 ounces) (5.5 grams)) is a member of the Emballonuridae, an Old World bat family that has an extensive distribution primarily in the tropics (Nowak 1994). The Pacific sheath-tailed bat was once common and widespread in Polynesia and Micronesia and is the only insectivorous bat recorded from a large part of this area (Hutson *et al.* 2001). Sheath-tailed bats are rich brown to dark brown above and paler below (Walker and Paradiso 1983). The common name “sheath-tailed bat” refers to the nature of the tail attachment; the tail pierces the tail membrane and its tip appears completely free on the upper surface of the membrane (Walker and Paradiso 1983). The biology of this species, including detailed information on reproduction, habitat use, and diet, is largely unknown (Hutson *et al.* 2001; Wiles and Worthington 2002).

Taxonomy

The classification of this species has received varied treatment, but the most thorough and recent taxonomic evaluation for this species was conducted by K. Koopman (Koopman 1997; Wiles and Worthington 2002). Koopman (1997) recognizes four subspecies: *E. s. rotensis*, endemic to the Mariana Islands (Guam and the Commonwealth of the Northern Mariana Islands (CNMI));

E. s. sulcata, occurring in Chuuk and Pohnpei; *E. s. palauensis*, found in Palau; and *E. s. semicaudata*, occurring in American and Independent Samoa, Tonga, Fiji, and Vanuatu. This assessment addresses the population of *E. s. semicaudata* that occurs in American Samoa. After review of the available taxonomic information, we conclude that *E. s. semicaudata* is a valid subspecies.

Habitat/Life History

The Pacific sheath-tailed bat is a small bat that appears to be cave-dependent, roosting during the day in a wide range of caves, including overhanging cliffs, crevices, and lava tubes (Grant 1993; Grant *et al.* 1994; Hutson *et al.* 2001; Palmeirim *et al.* 2005). Bats and cave swiftlets (*Aerodramus* spp.) may be found sharing caves (Lemke 1986; Hutson *et al.* 2001; Tarburton 2002; Wiles and Worthington 2002). Large roosting colonies appear fairly common in the Palau subspecies, but smaller aggregations may be more typical of at least the Mariana Island subspecies and perhaps other *Emballonura* (Nowak 1994; Flannery 1995; Wiles *et al.* 1997; Wiles and Worthington 2002). The subspecies that occurs on Aguiguan appears to prefer relatively large caves (Guam Division of Aquatic and Wildlife Resources (GDAWR) 1995). The Pacific sheath-tailed bat is nocturnal and typically emerges around dusk to forage on insects (Hutson *et al.* 2001). Survey work on Aguiguan in 2003 revealed that the Mariana Islands subspecies foraged almost entirely in forests (native and non-native) near their roosting caves (Esselstyn *et al.* 2004). Bruner and Pratt (1979) also observed sheath-tailed bats foraging in native forests on Pohnpei.

Historical Range/Distribution

American Samoa is made up of seven islands (Tau, Ofu, Olosega, Tutuila, Aunuu, Rose Atoll, and Swains Island). *Emballonura s. semicaudata* formerly occurred on Tau, Ofu, Olosega, collectively known as the Manua (Tau) Islands, Tutuila, and Aunuu (Amerson *et al.* 1982; Flannery 1995; Helgen and Flannery 2002; Department of Marine and Wildlife Resources (DMWR) 2006).

Current Range/Distribution

A precipitous decline of *Emballonura s. semicaudata* on the island of Tutuila has been documented since around 1990 (Amerson *et al.* 1982; Knowles 1988, Grant *et al.* 1994; Koopman and Steadman 1995; Helgen and Flannery 2002), and this subspecies may now be extirpated from American Samoa (Grant *et al.* 1994; R.C.B. Utzurrum, DMWR, pers. comm. 2005; DMWR 2006). In an attempt to ascertain whether the species is still extant, DMWR is in the midst of surveys, consisting of acoustic sweeps and cave checks, on all main islands (U.R. Tulafono, pers. comm. 2006). All of the main islands have been completed with the exception of the islands of Ofu and Olosega, and thus far the results have been negative (R. Utzurrum, pers. comm. 2007).

Population Estimates/Status

In American Samoa, Amerson *et al.* (1982) estimated a total population of approximately 11,000 Pacific sheath-tailed bats in 1975 and 1976. Since then, far fewer animals have been observed. Knowles (1988) recorded about 200 in 1988, and in 1993, observers caught one bat and saw only

three more (Grant *et al.* 1994). Ongoing systematic surveys by DMWR have failed to detect the presence of this species (DMWR 2006), and there is some concern that the subspecies may be extirpated in American Samoa (Grant *et al.* 1994; R. Utzurrum, pers. comm. 2005; DMWR 2006).

DISTINCT POPULATION SEGMENT (DPS)

The definition of “species” in section 3(15) of the Endangered Species Act (Act) includes any distinct population segment(s) of any species of vertebrate fish or wildlife that interbreed when mature. For a population to be listed under the Act as a distinct vertebrate population segment, three elements are considered: 1) the discreteness of the population segment in relation to the remainder of the species to which it belongs, 2) the significance of the population segment to the species to which it belongs, and 3) the population segment’s conservation status in relation to the Act’s standards for listing (i.e., is the population segment, when treated as if it were a species, endangered or threatened?) (61 FR 4722).

The available information indicates that distinct populations of *Emballonura s. semicaudata*, a cave-dwelling species whose populations are often highly localized, are definable. As expected in a species with broad geographic range and great discontinuities, the species as a whole shows strong geographic variation (Palmeirim *et al.* 2005). Koopman (1997) recognized four subspecies on morphological characters alone and it is possible that at least some of the subspecies will turn out to be sufficiently differentiated to be classified as distinct species (Palmeirim *et al.* 2005). The distinct population segment of Pacific sheath-tailed bat in American Samoa is discrete in relation to the remainder of the species as a whole. *Emballonura s. semicaudata* is known from American and Independent Samoa, Tonga, Fiji, and Vanuatu, but has shown drastic declines on the islands where it still occurs, or even to limited areas of single islands. In fact, it is believed to be extirpated from Independent Samoa (Grant *et al.* 1994; Koopman and Steadman 1995; Tarburton 2002) and may have never been present on Vanuatu (Flannery 1995; Koopman 1997; Palmeirim *et al.* 2005). The nearest possible population to the U.S. population is that of Independent Samoa, which is roughly 100 miles (161 kilometers) away and believed to be extirpated (Grant *et al.* 1994; Koopman and Steadman 1995; Tarburton 2002). The population segment of this species in American Samoa is therefore distinct based on geographic and distributional isolation from remaining Pacific sheath-tailed bats in Independent Samoa, Tonga, Fiji, and Vanuatu.

A population segment is considered “significant” if its loss would constitute a significant gap in the range of the taxon. The American Samoa population of the Pacific sheath-tailed bat represents the easternmost distribution of this species. The loss of this population would truncate the species’ range by approximately 100 miles (161 kilometers) and perhaps farther if the Independent Samoa population is also extirpated. Such a range reduction is significant when a species is declining range wide, as this one is, thereby reducing the possibility of dispersal and re-colonization from other areas. Furthermore, extremes of within-species variation in genotype, phenotype, and/or behavior are likely to occur at the periphery of species’ ranges (Erwin 1991); loss of the American Samoa population of the Pacific sheath-tailed bat therefore could constitute loss of important genetic or other variability within the species. The population structure of this species may be described as a metapopulation: although stochastic events may

result in extirpations on some islands, they may be re-colonized by dispersers coming from other populations (Palmeirim *et al.* 2005). Having such a structure, the species will only survive in an archipelago if the island colonization rate is sufficiently high to compensate for the rate of extirpation. Colonization rate is clearly proportional to the availability of source populations, therefore; if the rate of extirpation on individual islands due to stochastic events is higher than the rate of colonization, the long term result may be the permanent regional extinction of the species (Palmeirim *et al.* 2005). Based on the discreteness and significance of the American Samoa population of the Pacific sheath-tailed bat, the U.S. Fish and Wildlife Service (Service) considers this population to be a distinct vertebrate population segment which warrants review for listing under the Act. The American Samoa distinct population segment of the Pacific sheath-tailed bat faces severe and imminent threats. See SUMMARY OF THREATS and Rationale for Listing Priority Number, below.

THREATS

A. The present or threatened destruction, modification, or curtailment of its habitat or range.

The reasons for the decline of the sheath-tailed bat in American Samoa are unclear; however, the loss of roosting caves as a result of severe storms in American Samoa is believed to be a threat to this species (Craig *et al.* 1993; Grant *et al.* 1994). Two caves at Anapeapea Cove were reported as roosting sites for most of the bats estimated in 1976 and 1977 (Amerson *et al.* 1982). Severe storms, particularly Typhoons Ofa (1990) and Val (1991), removed the dense vegetation that had obscured the entrance to the larger cave at Anapeapea, inundated the cave with water, filled the cave with coral and fallen trees, and washed the cave walls clean (Grant *et al.* 1994). No bats were reported in either cave at Anapeapea during 1993 surveys (Grant 1993; Grant *et al.* 1994). Only small numbers of bats have been observed in other caves during those surveys, but there is no information on how many other caves there are, or how many bats they could support (Grant 1993; Grant *et al.* 1994).

In addition to the effects on the roost caves, storms are likely to have a negative impact on bats due to the deforestation (Palmeirim *et al.* 2005) that leads to loss of foraging habitat.

B. Overutilization for commercial, recreational, scientific, or educational purposes.

None known.

C. Disease or predation.

Rats (*Rattus* spp.) have been postulated as a problem for the Mariana Island subspecies of the Pacific sheath-tailed bat (Wiles and Worthington 2002) and are potential predators of Pacific sheath-tailed bats in American Samoa; however, the extent of this possibility is unknown. In addition, it is well known that domestic cats (*Felis catus*) can capture low flying bats, and it has been documented that they wait for bats as they emerge from caves and capture them in flight (Tuttle 1977; Ransome 1990; Woods *et al.* 2003). Consequently, even a few cats can have a major impact on a population of cave-dwelling bats (Palmeirim *et al.* 2005). Palmeirim *et al.* (2005) indicates that, of the predators introduced to Fiji, cats are the most likely to prey on bats. Feral cats are present on Tutuila and on Manua Islands in American Samoa (H. Freifeld, USFWS, pers. comm. 2007). The role of these introduced predators in the decline of the Pacific sheath-tailed bat is not proven but it is strongly supported as a factor (Palmeirim *et al.* 2005).

Likewise, the role of disease in the species' decline is not known as it has not been studied; however, disease could be a factor, especially for a communally roosting species such as the Pacific sheath-tailed bat (Wiles and Worthington 2002). While introduced pathogens are a plausible cause for the decline of this bat, there is currently no evidence for it (Palmeirim *et al.* 2005).

D. The inadequacy of existing regulatory mechanisms.

The inadequacy of existing regulatory mechanisms does not appear to be a threat to this species. The Pacific sheath-tailed bat has been afforded complete legal protection from “take” and “trapping” where “take” is defined as “...harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect or to attempt to engage in any such conduct,” in American Samoa since at least 1995 (Ch. 8, Title 24 of the American Samoa Administrative Code; U.R. Tulafono, pers. comm. 2006).

E. Other natural or manmade factors affecting its continued existence.

In addition to destroying suitable roosting caves, severe storms, such as Typhoons Ofa (1990) and Val (1991), likely killed the bats that were in the Anapeapea caves directly, and both storms may have led to the deaths of additional bats due to starvation (Grant 1993; Grant *et al.* 1994).

Other factors that may have led to the decline of the Pacific sheath-tailed bat in American Samoa are disturbance of roost caves by humans for recreational purposes (Wiles and Worthington 2002) and, more commonly, for guano mining (Grant *et al.* 1994; Tarburton 2002). Roost disturbance is a well known problem for many cave-dwelling species (Palmeirim *et al.* 2005). Disturbance at caves may cause bats to leave for alternate roost sites, in turn increasing their risk of predation and decreasing their roost time, the latter of which could increase stress.

Tarburton (2002) suggests that pesticides may have played an important role in the decline of bats on both American Samoa and Independent Samoa, particularly during the years in which taro was grown on large scales for export (1975-1985). However, pesticide use may not have been an important factor in bat declines on American Samoa (R. Utzurum, pers. comm. 2005).

The low numbers of individuals and populations of this subspecies place this bat at great risk of extinction from inbreeding and stochastic events such as storms (Wiles and Worthington 2002). The threat is significant for cave-dwelling species whose populations are often highly localized with few numbers of animals that can easily be lost in a severe storm, disease outbreak, or disturbance to the roost caves (Wiles and Worthington 2002).

CONSERVATION MEASURES PLANNED OR IMPLEMENTED

In an attempt to ascertain whether the species is still extant, DMWR is in the midst of surveys, consisting of acoustic sweeps and cave checks, on all main islands (U.R. Tulafono, pers. comm. 2006). All islands have been completed with the exception of the islands of Ofu and Olosega, and thus far the results have been negative (R. Utzurum, pers. comm. 2007). Upon completion of these surveys, DMWR intends to analyze cave sediments to assist in assessing the biological merits of reintroduction should the subspecies be deemed extirpated from the Territory (U.R. Tulafono, pers. comm. 2006).

SUMMARY OF THREATS:

In summary, current threats to this subspecies are its extremely small population size and limited distribution, habitat loss, and predation by introduced species. A potential threat is human disturbance to roosting caves (Grant *et al.* 1994; Hutson *et al.* 2001; Wiles and Worthington 2002). These threats indicate that the American Samoa DPS of the Pacific sheath-tailed bat continues to merit status as a candidate for listing. We find that this distinct population segment is warranted for listing throughout all of its range, and, therefore, find that it is unnecessary to analyze whether it is threatened or endangered in a significant portion of its range.

For species that are being removed from candidate status:

___ Is the removal based in whole or in part on one or more individual conservation efforts that you determined met the standards in the Policy for Evaluation of Conservation Efforts When Making Listing Decisions (PECE)?

RECOMMENDED CONSERVATION MEASURES:

- Complete surveys to determine whether the species is still extant on American Samoa.
- Conduct studies to determine causes of decline.
- Protect roost caves from disturbance.
- Control nonnative predators in and around roost caves.
- Conduct genetic analyses to determine if this subspecies warrants species status.
- Develop species augmentation or reintroduction plan, as appropriate.

LISTING PRIORITY

THREAT			
Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2
		Subspecies/population	3*
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/population	6
Moderate to Low	Imminent	Monotypic genus	7
		Species	8
		Subspecies/population	9
	Non-imminent	Monotypic genus	10
		Species	11
		Subspecies/population	12

Rationale for listing priority number:

Magnitude:

This species is highly threatened by its small size and extremely restricted distribution, which makes it highly susceptible to stochastic natural disturbances, such as typhoons and disease outbreaks; by loss of habitat (roosting caves and foraging habitat); and by predation from nonnative mammals. These threats occur throughout the Pacific sheath-tailed bat's range on American Samoa. Human disturbance of their roosting caves is a potential threat.

Imminence:

Threats to the Pacific sheath-tailed bat are imminent because they are ongoing.

Yes Have you promptly reviewed all of the information received regarding the species for the purpose of determining whether emergency listing is needed?

Is Emergency Listing Warranted?

No. The subspecies does not appear to be appropriate for emergency listing at this time because the immediacy of the threats is not so great as to imperil a significant proportion of the taxon within the timeframe of the routine listing process. The biggest threat at this time on American Samoa appears to be the vulnerability to stochastic events, such as typhoons, which cannot be addressed through the protections of listing the subspecies. If it becomes apparent that the routine listing process is insufficient to prevent significant losses that may result in this species' extinction, then the emergency rule process for this species will be initiated. We will continue to monitor the status of the Pacific sheath-tailed bat as new information becomes available. This review will determine if a change in status is warranted, including the need to make prompt use of emergency listing procedures.

DESCRIPTION OF MONITORING

The American Samoa DMWR is responsible for monitoring population status of *E. s. semicaudata*. This agency is funded through the Service's Federal Assistance program for wildlife restoration on an annual basis to monitor and manage the fish and wildlife resources of American Samoa, and the Service requests annual updates from DMWR on the status of candidate species. We also reviewed current scientific literature to seek new published information about the species in Samoa and elsewhere in its range, and we sent our most current information to regional and species experts for review. We received the latest update from DMWR on April 11, 2007; other queries and literature review yielded no new information. Surveys and monitoring of *E. s. semicaudata* in other areas throughout their range have been only sporadic and mostly undescribed. However, some more recent assessments of the Pacific sheath-tailed bat indicate a decline in numbers and distribution of Pacific sheath-tailed bats throughout its range in the tropical Pacific (Hutson *et al.* 2001; Helgen and Flannery 2002; Tarburton 2002; Palmeirim *et al.* 2005).

Although no regular surveys were previously conducted for the Pacific sheath-tailed bats in American Samoa, efforts are currently being made to ascertain whether the species is still extant. In 2004, researchers conducted searches at 80 percent of the roosting caves on Tutuila and made timed bat detector sweeps at three locations on Tutuila and two on Tau. Field assessments continued in 2005 and 2006, and are still ongoing (R. Utzurrum, pers. comms. 2005, 2007). All

islands have been completed with the exception of the islands of Ofu and Olosega, and thus far the results have been negative (R. Utzurrum, pers. comm. 2007).

This species is classified as endangered in the International Union for Conservation of Nature and Natural Resources (IUCN) Red Data List database (IUCN 2006), and is included in the Comprehensive Strategy for Wildlife Conservation in American Samoa (DMWR 2006).

COORDINATION WITH STATES

The latest assessment form was sent to Ray Tulafono, the Director of the American Samoa Department of Marine and Wildlife Resources on January, 29, 2010 . No additional information was provided.

LITERATURE CITED

- Amerson, A.B., Jr., W.A. Whistler, and T.D. Schwaner. 1982. Wildlife and wildlife habitat of American Samoa. II. Accounts of flora and fauna. U.S. Fish and Wildlife Service, Washington, D.C.
- Bruner, P.L. and H.D. Pratt. 1979. Notes on the status and natural history of Micronesian bats. *Elepaio* 40:1-4.
- Craig, P., P. Trail, G. Grant, J. Craig, and D. Itano. 1993. American Samoa: Natural history and conservation topics, Volume 1. Department of Marine and Wildlife Resources, Biological Report Series, Report No. 42.
- Department of Marine and Wildlife Resources. 2006. A comprehensive strategy for wildlife conservation in American Samoa (R.C.B. Utzurrum, J.O. Seamon, and K. Schletz Saili, authors). DMWR, Pago Pago. 109 pp.
- Erwin, T. L. 1991. An evolutionary basis for conservation strategies. *Science* 253:75-82.
- Esselstyn, J.A., G.J. Wiles, and A. Amar. 2004. Habitat use of the Pacific sheath-tailed bat (*Emballonura semicaudata*) on Aguiquan, Mariana Islands. *Acta Chiropterologica* 6: 303-308.
- Flannery, T. 1995. Mammals of the south-west Pacific and Moluccan Islands. Cornell University Press, Ithaca, New York.
- Grant, G.S. 1993. Sheath-tailed bats–Tutuila’s rarest mammal. Pp. 51-52 in American Samoa: Natural History and Conservation Topics, Vol. 1. Biological Report Series, Report No. 42, Department of Marine and Wildlife Resources, American Samoa.

- Grant, G.S., S.A. Banack, and P. Trail. 1994. Decline of the sheath-tailed bat Emballonura semicaudata (Chiroptera: Emballonuridae) on American Samoa. *Micronesica* 27:133-137.
- Guam Division of Aquatic and Wildlife Resources. 1995. Annual report fiscal year 1995, submitted to the U.S. Fish and Wildlife Service, Division of Federal Aid (Editors: R.D. Anderson, G.W. Davis, T.J. Pitlik, and G.J. Wiles).
- Helgen, K.M. and T.F. Flannery 2002. Distribution of the endangered Pacific sheath-tail bat (Emballonura semicaudata). *Australian Mammalogy* 24: 209-212.
- Hutson, A.M., S.P. Mickleburgh, and P.A. Racey (compilers). 2001. Microchiropteran bats: global status survey and conservation action plan. IUCN/SSC Chiroptera Specialist Group, International Union for the Conservation of Nature and Natural Resources, Gland, Switzerland and Cambridge, United Kingdom.
- IUCN 2006. 2006 IUCN red list of threatened species. <www.iucnredlist.org>. Downloaded on 12 April 2007.
- Knowles, W.C. 1988. The status of the wildlife and wildlife habitats in American Samoa. Annual report, Department of Marine and Wildlife Resources, American Samoa.
- Koopman, K.F. 1997. The subspecies of Emballonura semicaudata (Chiroptera: Emballonuridae). *J. Mamm.* 78(2):358-360.
- Koopman, K.F. and D.W. Steadman. 1995. Extinction and biogeography of bats on Eua, Kingdom of Tonga. *American Museum Novitates* 3125:1-13.
- Lemke, T.O. 1986. Distribution and status of the sheath-tailed bat (*Emballonura semicaudata*) in the Mariana Islands. *Journal of Mammalogy* 67:743-746.
- Nowak, R.M. 1994. Walker's bats of the world. The Johns Hopkins University Press, Baltimore, Maryland.
- Palmeirim, J.M., A. Champion, A. Naikatini, J. Niukula, M. Tuiwawa, M. Fisher, M. Yabaki-Gounder, S. Qalovaki, and T. Dunn. 2005. Distribution, status, and conservation of bats in the Fiji islands. Unpublished report to Flora and Fauna International. 68 pp.
- Ransome R. 1990. The natural history of hibernating bats. Christopher Helm, London. 235pp.
- Tarburton, M.K. 2002. Demise of the Polynesian sheath-tailed bat Emballonura semicaudata in Samoa. *Micronesica* 34: 105-108.
- Tuttle M.D. 1977. Gating as a means of protecting cave dwelling bats, pp. 77-82
In: T. Aley and D. Rhodes (eds.) National cave management symposium proceedings – 1976. Speleobooks, Albuquerque, NM. 106pp.

Walker, R.M. and J.L. Paradiso. 1983. Walker's mammals of the world, 4th edition. The Johns Hopkins University Press, Baltimore, Maryland.

Wiles, Gary J. and David J. Worthington. 2002. A population assessment of Pacific sheath-tailed bats (*Emballonura semicaudata*) on Aguiguan, Mariana Islands. Unpublished report to the U.S. Fish and Wildlife Service, Honolulu, HI. 31 pp.

Wiles, G. J., J. Engbring, and D. Otobed. 1997. Abundance, biology, and human exploitation of bats in the Palau Islands. *J. Zool. (London)* 241: 203-227.

Woods, M., R. A. McDonald, and S. Harris. 2003. Predation of wildlife by domestic cats *Felis catus* in Great Britain. *Mammal Rev.* 33(2):174-188.

Personal Communications and *in litt.*:

Brooke, A.P. U.S. Navy. 2005. Electronic message to Annie Marshall, U.S. Fish and Wildlife Service, regarding *Emballonura* listing, dated August 23, 2005.

Freifeld, H. U.S. Fish and Wildlife Service. 2007. Electronic message to Karen Rosa, U.S. Fish and Wildlife Service, regarding feral cats in American Samoa, dated April 12, 2007.

Tulafono, U.R. Department of Marine and Wildlife Resources. 2006. Letter to Patrick Leonard, U.S. Fish and Wildlife Service in response to request for review of the draft candidate assessment forms, dated August 31, 2006.

Utzurum, R.C.B. Department of Marine and Wildlife Resources. 2005. Electronic message to Annie Marshall, U.S. Fish and Wildlife Response, regarding the status of *E. semicaudata*, dated September 16, 2005.

Utzurum, R.C.B. Department of Marine and Wildlife Resources. 2007. Electronic message to Karen Rosa, U.S. Fish and Wildlife Response, regarding the sheath-tailed bat, dated April 11, 2007.

APPROVAL/CONCURRENCE: Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes, including elevations or removals from candidate status and listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all resubmitted 12-month petition findings, additions or removal of species from candidate status, and listing priority changes.

Approve:

Acting Carolyn D. Bohan 5/18/10
Regional Director, Region 1, Fish and Wildlife Service Date

Rowan W. Gould
ACTING :
Director, Fish and Wildlife Service October 22, 2010

Concur:

Do not concur: _____
Director, Fish and Wildlife Service Date

Director's Remarks:

Date of annual review: April 2, 2010
Conducted by: Lorena Wada, Pacific Islands FWO
Biologist, Prelisting and Listing Program

Comments:
PIFWO Review

Reviewed by: Christa Russell Date: April 5, 2010
Prelisting and Listing Program Coordinator

Marilet Zablan Date: April 26, 2010
Assistant Field Supervisor, Endangered Species Division

Gina Shultz Date: April 30, 2010
Acting Field Supervisor