

U.S. FISH AND WILDLIFE SERVICE SPECIES ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM

Scientific Name:

Porzana tabuensis

Common Name:

Spotless Crake

Lead region:

Region 1 (Pacific Region)

Information current as of:

06/19/2014

Status/Action

☐ Funding provided for a proposed rule. Assessment not updated.

☐ Species Assessment - determined species did not meet the definition of the endangered or threatened under the Act and, therefore, was not elevated to the Candidate status.

☐ New Candidate

☒ Continuing Candidate

☐ Candidate Removal

☐ Taxon is more abundant or widespread than previously believed or not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status

☐ Taxon not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status due, in part or totally, to conservation efforts that remove or reduce the threats to the species

☐ Range is no longer a U.S. territory

☐ Insufficient information exists on biological vulnerability and threats to support listing

☐ Taxon mistakenly included in past notice of review

☐ Taxon does not meet the definition of "species"

☐ Taxon believed to be extinct

☐ Conservation efforts have removed or reduced threats

___ More abundant than believed, diminished threats, or threats eliminated.

Petition Information

___ Non-Petitioned

X Petitioned - Date petition received: 05/11/2004

90-Day Positive:05/11/2005

12 Month Positive:05/11/2005

Did the Petition request a reclassification? **No**

For Petitioned Candidate species:

Is the listing warranted(if yes, see summary threats below) **Yes**

To Date, has publication of the proposal to list been precluded by other higher priority listing?
Yes

Explanation of why precluded:

Higher priority listing actions, including court-approved settlements, court-ordered and statutory deadlines for petition findings and listing determinations, emergency listing determinations, and responses to litigation, continue to preclude the proposed and final listing rules for this species. We continue to monitor populations and will change its status or implement an emergency listing if necessary. The Progress on Revising the Lists section of the current CNOR (<http://endangered.fws.gov/>) provides information on listing actions taken during the last 12 months.

Historical States/Territories/Countries of Occurrence:

- **States/US Territories:** American Samoa
- **US Counties:** Manu'a, AS
- **Countries:** Australia, Fiji, French Polynesia, Philippines, Tonga, Western Samoa

Current States/Counties/Territories/Countries of Occurrence:

- **States/US Territories:** American Samoa
- **US Counties:** Manu'a, AS
- **Countries:** Australia, Fiji, French Polynesia, Philippines, Tonga, Western Samoa

Land Ownership:

All of the land on which the spotless crane occurs in American Samoa is privately owned, but is under long-term lease (50 years) to the U.S. National Park Service (NPS) as part of the National Park of American Samoa.

Lead Region Contact:

ARD-ECOL SVCS, Jesse D'Elia, 5032312349, jesse_delia@fws.gov

Lead Field Office Contact:

PACIFIC ISLANDS FISH AND WILDL OFC, Kristi Young, 808-792-9419, kristi_young@fws.gov

Biological Information

Species Description:

The following description is from Watling (2001). The spotless crake is a very small rail, approximately 15 centimeters (6 inches) in length, and appears entirely black at first glance. On closer observation the grey wash on the neck, head and underparts, and its dark brown mantle and wings, are noticeable. Also conspicuous are the red iris and legs. The bill of this species is black. Immature birds have white marks on the chest and abdomen.

Taxonomy:

The genus *Porzana* is widespread in the Pacific, where it is represented by numerous island-endemic and flightless species (many of which are extinct as a result of anthropogenic disturbances) as well as several more cosmopolitan species, including the spotless crake. No subspecies of the spotless crake are recognized (Mayr 1945; Pratt et al. 1987; del Hoyo et al. 1996). We have carefully reviewed the available taxonomic information (Mayr 1945; Pratt et al. 1987; del Hoyo et al. 1996) and have concluded the species is a valid taxon.

Habitat/Life History:

Until 2001, all sightings of this species in American Samoa were from the island of Tau in marshy habitat or tall, moist, grassy swales, and this is typical of its habitat elsewhere (Watling 1982, 2001). In 2001, however, the spotless crake was discovered in dense cloud forest (rank vegetation) on the summit of Tau on Mt. Lata (Rauzon and Fialua 2003), the highest point in American Samoa. The presence of the spotless crake on Mt. Lata was reconfirmed in January 2011 (Tulafono 2011, pers. comm.). This is the only known site where this species is found in American Samoa.

The following is summarized from the account of this species in the Handbook of the Birds of the World (del Hoyo et al. 1996.) This species typically nests once annually and builds a cup nest of grasses on or near the ground in dense vegetation. Island populations probably produce fewer than the average of three to four eggs recorded in continental populations. The spotless crake is territorial and monogamous and may form lifetime pair bonds.

Historical Range/Distribution:

In American Samoa, the spotless crake was known from a series of 10 specimens that were collected from Tau in 1923 during the Whitney South Sea Expedition (Murphy 1924; Banks 1984).

Current Range Distribution:

This species was not found during surveys in 1976 (Amerson et al. 1982), and was reported as extirpated on Tau (Muse and Muse 1982), but three individuals were detected on Tau from 1985 to 1987 (Engbring and Engilis 1988; Engbring and Ramsey 1989). The spotless crake was rediscovered on the summit of Tau (Mt. Lata) in 2001, at roughly 945 meters (m) (3,100 feet (ft)) (Rauzon and Fialua 2003). In July 2001, two birds were heard and seen in dense vegetation near the summit field camp, and a similar small number of individuals were heard on the summit in December 2001 and December 2002 (Rauzon and Fialua 2003). One

bird was heard within 1,000 m (3,281 ft) of the same area in November 2003 (Seamon, in litt. 2004 and 2007). The spotless crane may occur in other remote areas of Tau, but difficult access and the extremely cryptic nature of this species complicate survey efforts.

Population Estimates/Status:

The population of the spotless crane in American Samoa is presumed to be very small and restricted to the summit of Tau Island. No current population estimate is available; the secretive habits of this species and difficult access to the summit of Tau make monitoring extremely difficult. In 2001, two cranes were observed during seabird surveys on the Tau summit (Rauzon and Fiaula 2003), and at least one bird was heard calling in the same area in 2003 (Seamon, in litt. 2004 and 2007). The presence of the spotless crane on Mt. Lata was reconfirmed in January 2011 (Tulafono 2011, pers.comm.). The presence of the species in other areas of Tau or American Samoa cannot be ruled out, but to date surveys have yielded no additional observations.

Distinct Population Segment(DPS):

The definition of species in section 3(15) of the Endangered Species Act (Act) includes any distinct population segment(s) (DPS) of any species of vertebrate fish or wildlife that interbreed when mature. For a population to be listed under the Act as a distinct vertebrate population segment, three elements are considered: 1) the discreteness of the population segment in relation to the remainder of the species to which it belongs, 2) the significance of the population segment to the species to which it belongs, and 3) the populations segments conservation status in relation to the Acts standards for listing (i.e., is the population segment, when treated as if it were a species, endangered or threatened?) (61 FR 4722).

The available information indicates that distinct populations of the spotless crane, a species not noted for long-distance dispersal, are definable. The DPS of spotless cranes in American Samoa is discrete in relation to the remainder of the species as a whole. Although the spotless crane (and other rails) have dispersed widely in the Pacific, island rails have tended to reduce or lose their power of flight over evolutionary time and so become isolated (Watling 2001). The population segment of this species in American Samoa is therefore distinct based on geographic and distributional isolation from spotless crane populations on other islands in the oceanic Pacific, the Philippines, and Australia.

A population segment is considered significant if its loss would constitute a significant gap in the range of the taxon. The American Samoa population of the spotless crane represents an important link in the Pacific range of the species; it is the link between the Central and Eastern Pacific portions of the species range. The loss of this population could cause an increase of roughly 805 km (500 mi) in the disjunction between the Central and Eastern Polynesian portions of the spotless cranes range in the Pacific, and could result in the increased isolation of the Marquesas and Society islands populations by further limiting the potential for genetic exchange.

Based on the discreteness and significance of the American Samoa population of the spotless crane, the U.S. Fish and Wildlife Service (FWS) considers this population to be a distinct vertebrate population segment which warrants review for listing under the Act. The DPS of the spotless crane faces imminent, severe threats. See Summary of Threats and Rationale for Listing Priority Number sections, below.

Threats

A. The present or threatened destruction, modification, or curtailment of its habitat or range:

The loss of wetland habitat probably contributed to the species likely extirpation from lowland habitats where it was recorded formerly. Almost all wetlands in American Samoa have been lost or converted to agricultural

fields. The non-wetland habitat where the species is known to occur today on the summit of Tau Island is afforded some protection from human disturbance by its remote location and its status as part of the National Park of American Samoa. The continuing loss of wetlands will further limit opportunities for restoration and conservation of the taxon.

B. Overutilization for commercial, recreational, scientific, or educational purposes:

None known.

C. Disease or predation:

Trapping on the summit of Tau, the site of the only known population of spotless crakes in American Samoa, revealed the presence of at least one species of rat, the Norway rat (*Rattus norvegicus*) (OConnor and Rauzon 2004). Rats are known to prey on the eggs, nestlings, and sometimes on adults of island bird species (Atkinson 1977, 1985) and ground-nesting species are especially vulnerable (Bertram and Nagorsen 1995; Flint 1999; Zino et al. 2001). Rail species all over the Pacific have been decimated by predation (Steadman 1995). Predation by rats likely poses a significant threat for this small, ground-nesting bird. In addition, native predators include the Pacific boa (*Candoia bibroni*) and the purple swamphen (*Porphyrio porphyrio*). Both species have the potential to predate the eggs and young of the spotless crane (Tulafono, in litt. 2011).

D. The inadequacy of existing regulatory mechanisms:

Wetlands, which may be an important habitat for this species and a refugium from rats and other predators (Watling 2001), are protected under American Samoa Government and U.S. Federal laws, but enforcement of these laws historically has been weak (Seamon, in litt. 2004). The NPS has sporadically indicated interest in controlling predators on the summit of Tau Island, but logistics and costs have so far proven prohibitive (OConnor and Rauzon 2004).

E. Other natural or manmade factors affecting its continued existence:

The probable low number of individuals and the extremely restricted distribution of this population place this species at great risk of extinction in American Samoa from stochastic events such as hurricanes and from inbreeding depression (Meffe and Carroll 1997; Thevenon and Couvet 2002; Frankham 2003). Hurricanes may cause the direct and indirect mortality of some birds, as well as modify the already limited habitat. This species has coexisted with hurricanes in Samoa for millennia. If the spotless crane were widely distributed in American Samoa, had ample habitat and sufficient numbers, and were not under chronic pressure from anthropogenic threats such as introduced predators, it might recover from hurricane-related mortality and the temporary loss or redistribution of resources in the wake of severe storms. However, this species current status in American Samoa makes it highly vulnerable to a single, chance event. Two hurricanes (Heta in 2004 and Olaf in 2005) have hit Samoa since the last visit to the Tau summit in 2003, and the impact of these storms to the crane and its habitat are unknown. The remote location of the one known population and the concomitant logistical and fiscal challenges make conservation actions such as monitoring surveys and rat control difficult to accomplish.

Conservation Measures Planned or Implemented :

The Department of Marine and Wildlife Resources (DMWR) has conducted surveys on Tau in habitats and locations other than the summit that potentially harbor spotless crakes, but these surveys have not yielded additional observations (Seamon 2004, in litt.; Miles 2014, in litt.). One spotless crane was captured in a Tomahawk trap set for rats on the Summit of Mt. Lata in February 2011 (Miles 2014, in litt.). Surveys using playbacks were conducted during the breeding season on the island of Ta'u from May until September 2013. Numerous vocalizations in response to playbacks were documented, however, only one visual sighting was

reported (Badia 2013, in litt.). Conducting additional surveys in areas on Ta'u not covered by the 2013 effort is recommended.

Summary of Threats :

The population in American Samoa is threatened by small population size, limited distribution, and predation by nonnative mammals. The loss of wetlands is a continuing threat, as well. The co-occurrence of the only known population of the spotless crane under U.S. jurisdiction with a known predator of ground-nesting birds, the Norway rat, along with the extremely restricted observed distribution and low numbers of this taxon indicate that the American Samoa DPS of the spotless crane continues to merit status as a candidate for listing. We find that this distinct population segment is warranted for listing throughout all its range, and, therefore, find that it is unnecessary to analyze whether it is threatened or endangered in a significant portion of its range.

For species that are being removed from candidate status:

_____ Is the removal based in whole or in part on one or more individual conservation efforts that you determined met the standards in the Policy for Evaluation of Conservation Efforts When Making Listing Decisions(PECE)?

Recommended Conservation Measures :

- Conduct surveys in suitable habitat for additional individuals/populations and determine the size and status of each population
- Conduct studies to determine habitat requirements and basic life history of the species
- Control predators (rats) at known locations
- Determine other limiting factors and implement control measures

Priority Table

Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2
		Subspecies/Population	3
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/Population	6
Moderate to Low	Imminent	Monotype genus	7
		Species	8
		Subspecies/Population	9
	Non-Imminent	Monotype genus	10
		Species	11
		Subspecies/Population	12

Rationale for Change in Listing Priority Number:

Magnitude:

The magnitude of the threats facing the spotless crane is high because of its observed small population, limited distribution, and the especially serious threat posed by introduced predators. Populations of other ground-nesting island birds, including populations of this species on other islands and numerous other rail species, have been limited or extirpated by such predators. Single, small populations are vulnerable to inbreeding depression, and are significantly more vulnerable than widespread, abundant species to substantial reductions or extinction resulting from direct mortality or habitat destruction caused by chance natural occurrences such as hurricanes. The entire American Samoa population of this taxon is likely to be affected by these threats.

Imminence :

Threats to the spotless crane (predation and small population size) are imminent because they are ongoing.

Yes Have you promptly reviewed all of the information received regarding the species for the purpose of determination whether emergency listing is needed?

Emergency Listing Review

No Is Emergency Listing Warranted?

The species does not appear to be appropriate for emergency listing at this time because the immediacy of the threats is not so great as to imperil a significant proportion of the taxon within the timeframe of the routine listing process. In addition, the population on American Samoa is afforded some protection and management because it occurs within the National Park of American Samoa. We will continue to monitor the status of the spotted crane in American Samoa as new information becomes available. This review will determine if a change in status is warranted, including the need to make prompt use of emergency listing procedures.

Description of Monitoring:

DMWR is responsible for monitoring the spotless crane and is the sole source of current, on-the-ground information about wildlife in the Territory. This agency is funded through the U.S. Fish and Wildlife Services Federal Assistance program for wildlife restoration on an annual basis to monitor and manage the fish and wildlife resources of the Territory of American Samoa, and the FWS requests annual updates from DMWR on the status of candidate species. We also requested information from biologists with the National Park of American Samoa, reviewed current scientific literature to seek new published information about the species in Samoa and elsewhere in its range, and sent our most current information to regional and species experts for review. We received the latest update from DMWR on August 31, 2006; other queries and literature review yielded no new information. In addition to the surveys described under the Conservation Measures Planned or Implemented Section above, irregular visits are made to the Tau summit, where the agency conducts surveys using night-vision and videography to verify that the spotless crane still is present. The last visit was made in 2003, and two hurricanes have struck Samoa since then. The current rate of detection of this species is insufficient to accurately estimate total population size. The spotless crane is a cryptic species and its only known occurrence in American Samoa is in an extremely remote location, therefore it is difficult to monitor. While more intensive population monitoring would certainly be desirable given these challenges, our assessment uses the best available scientific and commercial data and is therefore adequate to conduct our annual assessment.

This species is classified as a taxon of Least Concern (LC) in the International Union for Conservation of Nature and Natural Resources (IUCN) Red Data List database (IUCN 2006), and is included as a priority species for new studies in the Comprehensive Strategy for Wildlife Conservation in American Samoa (DMWR 2006).

Indicate which State(s) (within the range of the species) provided information or comments on the species or latest species assessment:

none

Indicate which State(s) did not provide any information or comment:

American Samoa

State Coordination:

On February 22, 2013, we sent a letter to the American Samoa Department of Marine and Wildlife Resources requesting their review and comment on our most recent candidate assessment of this species. No additional information or comments were received. We met with American Samoa biologists in November, 2013, and discussed candidate species. Therefore, we believe this assessment contains the most recent available information about the species.

Literature Cited:

Amerson, A.B., Jr., W.A. Whistler, and T.D. Schwaner. 1982. Wildlife and wildlife habitat of American Samoa. II. Accounts of flora and Fauna. U.S. Fish and Wildlife Service. 151 pp.

Atkinson, I.A.E. 1977. A reassessment of factors, particularly *Rattus rattus* L., that influenced the decline of endemic forest birds in the Hawaiian Islands. *Pacific Science* 31:109-133.

Atkinson, I.A.E. 1985. The spread of commensal species of *Rattus* to oceanic islands and their effects on island avifaunas. Pp. 35-81. In Moors, P.J. (ed.), *Conservation of island birds*. Tech. Publ. No. 3, Int. Coun. Bird Preserv., Cambridge, England.

Banks, R.C. 1984. Bird specimens from American Samoa. *Pacific Science* 38:150-169.

Bertram, D., and D. Nagorsen. 1995. Introduced rats *Rattus* spp. on the Queen Charlotte Islands: implications for seabird conservation. *Canadian Naturalist* 10:6-10.

del Hoyo, J., A. Elliott, and J. Sargatal. 1996. *Handbook of the Birds of the World*. Volume 3: Hoatzin to Auks. Lynx Ediciones. 752 pp.

(DMWR) Department of Marine and Wildlife Resources. 2006. A comprehensive strategy for wildlife conservation in American Samoa (R.C.B. Utzurrum, J.O. Seamon, and K. Schletz Saili, authors). DMWR, Pago Pago. 109 pp.

Engbring, J., and A.E. Engilis. 1988. Rediscovery of the sooty rail (*Porzana tabuensis*) in American Samoa. *Auk* 105:391.

Engbring, J., and F.L. Ramsey. 1989. A 1986 survey of the forest birds of American Samoa. U.S. Fish and Wildlife Service. 145 pp.

Flint, E. 1999. Status of seabird populations and conservation in the tropical island Pacific. In Eldredge, L.P., P. Holtus, and J. Maragos (eds.), *Marine and coastal biodiversity in the tropical island Pacific region: population, development, and conservation priorities*, vol. 2. East-West Center, Honolulu, Hawaii. 32 pp. plus appendices.

- Frankham, R. 2003. Genetics and conservation biology. C.R. Biologies, 326 Suppl 1:S22-9.
- (IUCN) International Union for Conservation of Nature and Natural Resources. 2006 IUCN Red List of Threatened Species. www.iucnredlist.org, accessed on April 10, 2007.
- Mayr, E. 1945. Birds of the Southwest Pacific. The Macmillan Company, New York, NY. 316 pp.
- Meffe, G.K., and C.R. Carroll. 1997. Principles of Conservation Biology, 2nd ed. Sinauer Associates, Sunderland, MA. 729 pp.
- Murphy, R.C. 1924. Birds collected during the Whitney South Sea Expedition. II. American Museum Novitates 124:1-13.
- Muse, C., and S. Muse. 1982. The birds and birdlore of Samoa. Pioneer Press, Walla Walla, Washington. 156 pp.
- OConnor, P.J., and M.J. Rauzon. 2004. Inventory and monitoring of seabirds in national park of American Samoa. National Park of American Samoa Technical Report 136. 140 pp.
- Pratt, H.D., P.L. Bruner, and D.G. Berrett. 1987. A field guide to the birds of Hawaii and the tropical Pacific. Princeton University Press, Princeton, NJ. 409 pp. plus plates.
- Rauzon, M., and M. Fialua. 2003. Status of the spotless crane (*Porzana tabuensis*) in American Samoa. Wilson Bulletin 115:489-491.
- Steadman, D.W. 1995. Prehistoric extinctions of Pacific island birds: biodiversity meets zooarchaeology. Science 267:1123-1131.
- Thivenon, S., and D. Couvet. 2002. The impact of inbreeding depression on population survival depending on demographic parameters. Animal Conservation 5: 53-60.
- Watling, D. 1982. Birds of Fiji, Tonga, and Samoa. Millwood Press, Wellington, New Zealand. 176 pp.
- Watling, D. 2001. Birds of Fiji and Western Polynesia. South Pacific Regional Environmental Programme, Apia, Samoa. 272 pp.
- Zino, F., P. Oliviera, S. King, A. Buckle, M. Biscoito, H. Costa Neves, and A. Vasconcelos. 2001. Conservation of Zinos petrel *Pterodroma madeira* in the archipelago of Madeira. Oryx 35:128-135.

Personal Communications and In Litteris

- Badia, R., Contractor, DMWR. 2013. Field Notes regarding audio and visual bird surveys conducted on the island of Ta'u from May through July 2013. 13 pp.
- Miles, A., Chief Biologist , American Samoa DMWR. 2014. Electronic mail message to Carrie Harrington, FWS, regarding the Spotless Crane and Friendly Ground Dove, dated August 14, 2014.
- Seamon, J.O., DMWR. 2004. Electronic mail message to Holly Freifeld, FWS, regarding candidate status updates for American Samoa, dated May 31, 2004.
- Seamon, J.O., DMWR. 2007. Electronic mail message to Karen Rosa, FWS, regarding spotless crane candidate assessment update, dated April 12, 2007.

Tulafono, U.R., Director, American Samoa DMWR. Emailed letter dated March 31, 2011, regarding the Departments response to candidate assessment forms, received April 4, 2011.

Approval/Concurrence:

Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes, including elevations or removals from candidate status and listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all resubmitted 12-month petition findings, additions or removal of species from candidate status, and listing priority changes.

Approve:



06/18/2014

Date

Concur:



11/18/2014

Date

Did not concur:

Date

Director's Remarks: