

Recovery Plan for Four Subspecies of Island Fox *(Urocyon littoralis)*




Photo courtesy of Dan Richards

Recovery Plan for Four Subspecies of Island Fox (*Urocyon littoralis*)

(2015)

**Region 8
U.S. Fish and Wildlife Service
Sacramento, California**

Approved: _____

A handwritten signature in black ink, appearing to read "Dan L. Johnson", written over a horizontal line.

Regional Director, Pacific Southwest Region, Region 8
U.S. Fish and Wildlife Service

Date: February 27th, 2015

Recovery Plan for Four Subspecies of Island Fox

DISCLAIMER

Recovery plans delineate reasonable actions that are believed to be required to recover and/or protect federally listed species. We, the U.S. Fish and Wildlife Service, publish recovery plans, sometimes preparing them with the assistance of recovery teams, contractors, State agencies, and others. Objectives will be attained and any necessary funds made available subject to budgetary and other constraints affecting the parties involved, as well as the need to address other priorities. Recovery plans do not necessarily represent the views nor the official positions or approval of any individuals or agencies involved in the plan formulation, other than our own. They represent our official position only after they have been signed by the Regional Director or Director as approved. Approved recovery plans are subject to modification as dictated by new findings, changes in species status, and the completion of recovery tasks.

Notice of Copyrighted Material

Permission to use copyrighted illustrations and images in this recovery plan has been granted by the copyright holders. These illustrations **are not** placed in the public domain by their appearance herein. They cannot be copied or otherwise reproduced, except in their printed context within this document, without the written consent of the copyright holder.

Literature citation should read as follows:

U.S. Fish and Wildlife Service. 2015. Recovery Plan for Four Subspecies of Island Fox (*Urocyon littoralis*). U.S. Fish and Wildlife Service, Sacramento, California. xiv + 180 pp

An electronic copy of this recovery plan will be made available at <http://www.pacific.fws.gov/ecoservices/endangered/recovery/plans.html> and <http://endangered.fws.gov/recovery/index.html#plans>

ACKNOWLEDGEMENTS

There were many contributors to the development of this recovery plan. We would like to acknowledge the staff of the Ventura Fish and Wildlife Office, the Island Fox Recovery Coordination Group, the technical experts to the recovery team, and the researchers and field biologists for their contributions to this recovery plan. We appreciate the long-term commitment of the recovery team and their assistance in preparing this recovery plan over the period of development. Affiliations of the following individuals can be found in Appendix 1 of this document.

Primary Authors

This recovery plan was developed by the Island Fox Recovery Coordination Group: Brian Cypher, Carlos de la Rosa, David Graber, Devra Kleiman, Robert McMorrان, Eric Morrissette, Peter Schuyler, Rebecca Shaw, Dale Steele, and Rosie Woodroffe. Peter Schuyler served as the Chair of the Recovery Team, and Eric Morrissette served as the Recovery Team Coordinator.

Additional Contributors

The following people provided valuable assistance in the development of this recovery plan:

Vickie Bakker, Christina Boser, Paul Collins, Tim Coonan, Dan Doak, Julie King, Colleen Lynch, Linda Munson, Kathy Ralls, Esther Rubin, Winston Vickers, Kirk Waln, and Rachel Wolstenholme.

We thank the land management agencies/organizations and members of their staff for their continued support of island fox recovery efforts: the National Park Service - Channel Islands National Park, The Nature Conservancy, and the Santa Catalina Island Conservancy. We also recognize the contributions from the Integrated Island Fox Recovery Team and the Institute for Wildlife Studies in their efforts to aid in the development of this plan and promote island fox conservation and recovery.

We further recognize the Friends of the Island Fox, Inc. for their efforts to promote island fox conservation and recovery through public outreach and education.

DEDICATION

This recovery plan is dedicated to the memory of Devra Kleiman and Linda Munson, who passed away in 2010. They were both instrumental in developing this plan and in constructing effective recovery actions for island foxes.

Devra served on the U.S. Fish and Wildlife Service's Island Fox Recovery Coordination Group, and as such was an author of this document. She was truly a giant in conservation biology and generously lent her considerable talents to island fox recovery. She was a passionate advocate of implementing recovery actions in a scientific manner, and of appropriate environmental education, and she successfully integrated the mainland zoo community into the island fox conservation effort. Her forthright, direct style of communication enlivened many recovery team meetings.

Linda Munson was one of the world's leading carnivore pathologists, and led studies of island fox disease and pathology since 1999. She conducted hundreds of island fox necropsies, and it is not hyperbole to say that all of our current insight into island fox health issues is due to her tireless work. Linda led the Island Fox Health Group since 1999, and under her guidance they were always the most disciplined, effective and productive of the island fox recovery team working groups. Among the issues she discovered or brought to the forefront were the unique, island-specific parasite fauna of island foxes, the nature of the canine distemper strain on Santa Catalina Island and the endemic distemper-like virus found naturally in island fox populations, chronic amyloidosis, ear tumors on Santa Catalina, and neonatal deaths from maternal neglect.

Recovery Plan for Four Subspecies of Island Fox

EXECUTIVE SUMMARY

In 2004, the U.S. Fish and Wildlife Service (FWS) listed four of the six subspecies of island fox endemic to the California Channel Islands as endangered under the Endangered Species Act of 1973, as amended (Act), following catastrophic population declines (69 FR 10335). The San Miguel Island fox (*Urocyon littoralis littoralis*) went from an estimated 450 individuals to 15; the Santa Rosa Island fox (*U. l. santarosae*) declined from over 1,750 individuals to 15; the number of Santa Cruz Island foxes (*U. l. santacruzae*) went from approximately 1,450 individuals to approximately 55; and the Santa Catalina Island fox (*U. l. catalinae*) declined from over 1,300 to 103. The San Clemente Island fox (*U. l. clementae*) and the San Nicolas Island fox (*U. l. dickeyi*) were not federally listed, as their population numbers had not experienced similar declines. However, both non-federally listed subspecies could experience the same type of population decline as those subspecies that are federally listed. Additionally, all six subspecies are listed as threatened by the State of California. Therefore, the San Clemente and San Nicolas Island fox subspecies are included for discussion in this plan where appropriate.

The Channel Islands inhabited by island foxes are owned by four major landowners: the National Park Service (NPS), the U.S. Navy (Navy), The Nature Conservancy (TNC), and the Santa Catalina Island Conservancy (CIC). Although San Miguel Island is owned by the Navy, it is within Channel Islands National Park and is managed by the NPS. The NPS, TNC, and CIC manage the islands where the listed subspecies occur. The FWS guides the recovery planning process for the four federally-listed island fox subspecies under the Act. In addition, the State of California has regulatory authority over the island fox on non-Federal lands because the species is listed as threatened under the California Endangered Species Act.

The two primary known threats that resulted in the listing of the four subspecies of island fox as federally endangered were: 1) predation by golden eagles (*Aquila chrysaetos*) (San Miguel Island fox, Santa Rosa Island fox, and Santa Cruz Island fox) and 2) the transmission of canine distemper virus (Santa Catalina Island fox). Additionally, because the size of each island fox population was small, they were highly vulnerable to **stochastic events** and the effects of low genetic diversity. Other threats that either contributed to the decline of island foxes or continue to affect the island fox subspecies and/or their habitat include habitat degradation from grazing; other diseases and parasites; competition with feral cats, deer, and pigs; and mortality from vehicle strikes. We note however, many of these threats are not present on all islands, such as vehicle strikes, feral cats, pigs, and deer.

The current knowledge regarding the evolution, ecology, behavior, and population biology of island foxes has been amassed by numerous researchers from around the

Recovery Plan for Four Subspecies of Island Fox

country. Collaboration with researchers has been, and will continue to be, critically important in understanding island fox natural history and recovery challenges.

Recovery of each subspecies of island fox will be achieved by removing, or substantially reducing, known threats and increasing populations to viable levels for long-term survival of each subspecies. The strategy of this recovery plan is to continue the current recovery efforts and to improve and expand recovery actions as necessary. Recent and ongoing island fox recovery efforts include: removing golden eagles from the northern Channel Islands; reducing the threat of disease; breeding island foxes in captivity and reintroducing them to the wild; monitoring wild island fox populations; removal of non-native species (e.g., golden eagle prey); and while not part of island fox recovery efforts, reintroducing bald eagles (*Haliaeetus leucocephalus*) with the goal of deterring golden eagles from establishing territories on the Channel Islands. Additionally, ongoing activities that contribute to a long-term conservation strategy include: conducting research on behavioral ecology and reproductive biology; increasing island fox education and outreach activities to reduce anthropogenic impacts; restoring island habitat; and assessing the **demographic** impact of other threats such as mortality from vehicles, competition with feral cats, and emerging disease issues (e.g., ear cancer).

Since 1999, island fox recovery efforts by the land management agencies (NPS, TNC, and CIC) have included efforts to reduce the two major threats to island foxes that caused the precipitous declines. Mortality due to golden eagle predation on the three island fox subspecies from the northern Channel Islands (San Miguel, Santa Rosa, and Santa Cruz Islands) has been reduced. The threat posed by disease to Santa Catalina Island foxes continues to be ameliorated with disease mitigation efforts. All land management efforts have included bringing wild island foxes into captivity to serve as a temporary sanctuary from threats, increasing populations of each subspecies through captive breeding, and releasing captive individuals back into the wild. For a period of time, the entire San Miguel Island and Santa Rosa Island fox populations were held in captivity. Reintroduction was eminently successful. Released foxes had high survival and reproductive success, and recovering fox populations grew rapidly. This allowed management agencies to cease all captive breeding and reintroduction by 2008.

Recovery efforts to date have increased the numbers of foxes on all islands and reduced the risk of extinction. Wild populations of island foxes have been re-established on San Miguel and Santa Rosa Islands. Predation has been a negligible mortality factor on San Miguel Island and Santa Cruz Island, where annual survival of island foxes has remained greater than 80 percent since 2004 for San Miguel Island and since 2008 for Santa Cruz Island. Annual survival has been greater than 80 percent on Santa Rosa Island since 2008, though eagle predation occurred again in 2010.

Recovery Plan for Four Subspecies of Island Fox

Following disease mitigation efforts, the Santa Catalina Island fox population is increasing. The threat of disease, such as that posed by ear tumors, is of continued concern for the Santa Catalina Island fox. Other potential threats to the Santa Catalina Island foxes include competition with feral cats and mortality from vehicle strikes.

As of 2013, island fox populations increased to >1,000 individuals on Santa Catalina and Santa Cruz Islands, almost 900 on Santa Rosa Island, and close to 600 on San Miguel Island. Additionally, all island fox subspecies currently have annual survival estimates greater than 80 percent.

Recovery Goal: The goal of this recovery plan is to recover the San Miguel Island fox, the Santa Rosa Island fox, the Santa Cruz Island fox, and the Santa Catalina Island fox so they can be delisted (removed from listing under the Act) when existing threats to each respective subspecies have been ameliorated, thereby stabilizing and augmenting their populations. Each listed subspecies may be considered for downlisting or delisting independently of the other subspecies.

Recovery Objectives: Recovery objectives identify mechanisms for measuring progress toward and achieving the recovery goal for each subspecies. Achieving the recovery goal requires: 1) increasing the population size to levels and demographic rates that are self-sustaining, and 2) reducing or eliminating the current threats to the survival of each subspecies.

Recovery Objective 1:

Each federally listed subspecies of island fox exhibits demographic characteristics consistent with long-term viability.

Recovery Objective 2:

Land managers are able to respond in a timely fashion to predation by nesting golden eagles or significant predation rates by transient golden eagles, to potential or incipient disease outbreaks, and to other identified threats using the best available technology.

In order for any one of the four listed subspecies of island fox to be considered for downlisting from endangered to threatened status, recovery objective 1 should be met.

In order for any one of the four listed subspecies of island fox to be considered for delisting from endangered or threatened to delisted status, recovery objective 1 and recovery objective 2 should be met.

Recovery Criteria

Recovery criteria are measurable standards for determining that a species has achieved its recovery objectives and may be considered for downlisting or delisting. The

Recovery Plan for Four Subspecies of Island Fox

recovery criteria presented in this recovery plan represent our best assessment of the conditions that would most likely result in a determination that downlisting and/or delisting of the San Miguel Island fox, Santa Rosa Island fox, Santa Cruz Island fox, or Santa Catalina Island fox is warranted.

Population Risk-based Recovery Criteria

Recovery criterion E/1 was developed to address recovery objective 1.

Recovery Criterion E/1:

An island fox subspecies has no more than 5 percent risk of quasi-extinction over a 50-year period. This risk level is based upon the following:

- Quasi-extinction is defined as a population size of ≤ 30 individuals.
- The risk of extinction is calculated based on the combined lower 80 percent confidence interval for a 3-year running average of population size estimates, and the upper 80 percent confidence interval for a 3-year running average of mortality rate estimates.
- This 5 percent (or less) risk level is sustained for at least 5 years, during which time the population trend is not declining. A declining trend is defined as the 3-year risk-level being greater in year 5 than year 1.

This risk-based recovery criterion is based on models developed separately for each listed subspecies.

Threat-based Recovery Criteria

To meet recovery objective 2, recovery criteria C/1 and C/2 are achieved.

Recovery Criterion C/1 – Golden Eagle Predation:

To reduce the threat of extinction to the San Miguel Island fox, the Santa Rosa Island fox, and the Santa Cruz Island fox from golden eagle predation:

1. A golden eagle management strategy is developed and approved by the land manager(s) in collaboration with the FWS, including review by the appropriate Integrated Island Fox Recovery Team (IRT) Technical Expertise Group (TEG) or the equivalent. This strategy must include:
 - Response tactics (including the use of helicopters and net-guns) to capture nesting golden eagles and any transient golden eagle responsible for significant island fox predation per the golden eagle response strategy;

Recovery Plan for Four Subspecies of Island Fox

- Tactics to minimize the establishment of successful nesting golden eagles;
 - An established island fox monitoring program for each subspecies that is able to detect an annual island fox predation rate caused by golden eagles of 2.5 percent or greater, averaged over 3 years; and
 - An established mortality rate or population size threshold for each subspecies of island fox that, if reached due to golden eagle predation, would require the land manager(s) to bring island foxes into captivity for safety.
2. The golden eagle prey base of deer and elk is removed from Santa Rosa Island. At present, golden eagles are not known to prey upon Santa Catalina Island foxes. If mortality as a result of golden eagle predation becomes a threat to the Santa Catalina Island fox, the golden eagle management strategy will be implemented.

Recovery Criterion C/2 – Disease:

To reduce the threat of extinction to the San Miguel Island fox, the Santa Rosa Island fox, the Santa Cruz Island fox, and the Santa Catalina Island fox from disease:

1. A disease management strategy is developed, approved, and implemented by the land manager(s) in collaboration with the FWS, including review by the appropriate IRT TEG or the equivalent. This strategy must include:
 - Identification of a portion of each subspecies that will be vaccinated against canine distemper virus and rabies, the diseases posing the greatest risk and for which vaccines are safe, effective, and available. Vaccinations to be provided and numbers vaccinated will be developed in consultation with appropriate subject-matter experts;
 - Identification of actual and potential pathogens of island foxes, and the means by which these can be prevented from decimating fox populations;
 - Measures to prevent diseases in island foxes;
 - A monitoring program that provides for timely detection of a disease outbreak, and an associated emergency response strategy as recommended by the appropriate subject-matter experts; and
 - A process for updating the disease management strategy as new information arises.

Recovery Actions

The actions identified below are those that, in our opinion, are necessary to bring about the recovery of island foxes. These actions are subject to modification as dictated by new findings, changes in species status, and the completion of recovery actions.

1. Reduce mortality for each subspecies of island fox to ensure populations persist at sustainable levels.
2. Conduct captive breeding and reintroduction of island foxes to increase population size.
3. Establish island fox monitoring strategies.

Estimated Total Cost of Recovery

San Miguel Island fox: \$1,076,000

Santa Rosa Island fox: \$1,076,000

Santa Cruz Island fox: \$1,076,000

Santa Catalina Island fox: \$5,257,500

Grand Total: \$8,485,500, plus costs that are unable to be determined at this time.

Date of Recovery:

If recovery criteria are met, we estimate the northern Channel Islands fox subspecies (San Miguel Island fox, Santa Rosa Island fox, and Santa Cruz Island fox) could be recovered by 2020 (4 years) and the Santa Catalina Island fox by 2024 (8 years).

Long-term Conservation Strategy

The long-term conservation strategy included herein identifies actions that would further the conservation of the island fox. Long-term conservation may be benefitted by conducting research on behavioral ecology and reproductive biology; increasing island fox education and outreach activities to reduce anthropogenic impacts; restoring island habitat; and assessing the **demographic** impact of other threats such as mortality from vehicles, competition with feral cats, and emerging disease issues (e.g., ear cancer). At this time, these activities are not essential for preventing extinction and are not required for downlisting or delisting a particular island fox subspecies; however, these activities could substantially enhance the long-term conservation of the species and may also increase our scientific understanding of the island fox. In the event that an island fox subspecies is recovered and delisted, completion of these actions may further reduce the potential for any of the subspecies to be relisted in the future.

TABLE of CONTENTS

I.	Background	1
	A. Introduction.....	1
	1. Legal Status.....	1
	2. Affected Agencies, Landowners, and Partners	3
	3. Integration of Conservation and Recovery Efforts	6
	B. Biological Information.....	7
	1. Description and Taxonomy	7
	2. Distribution, Evolution, and Genetics.....	7
	3. Habitat Use and Food Habits	9
	4. Social Organization and Reproduction	10
	5. Mortality Sources and Population Dynamics.....	11
	C. Historical Population Status and Observed Declines of Island Fox Populations	15
	1. San Miguel Island	16
	2. Santa Rosa Island.....	17
	3. Santa Cruz Island	18
	4. Santa Catalina Island.....	18
	5. San Clemente Island.....	19
	6. San Nicolas Island.....	20
	D. Threats to the Species	21

Recovery Plan for Four Subspecies of Island Fox

1.	Summary of Listing Factors A through E	29
E.	Recovery and Conservation Efforts	29
1.	Recovery Actions for Listed Island Fox Subspecies.....	29
2.	Conservation Efforts for non-Listed Island Fox Subspecies.....	37
3.	Monitoring Efforts	39
F.	Current Status and Trend.....	40
1.	San Miguel Island fox	40
2.	Santa Rosa Island fox.....	41
3.	Santa Cruz Island fox	42
4.	Santa Catalina Island fox	43
5.	San Clemente Island fox	44
6.	San Nicolas Island fox	44
7.	Summary	45
II.	Recovery Strategy.....	47
III.	Recovery Goals, Objectives, and Criteria.....	49
A.	Recovery Goal	49
B.	Recovery Objectives.....	49
C.	Recovery Criteria.....	50
IV.	Recovery Program.....	55
A.	Recovery Action Narrative.....	55
1.	Reduce mortality and maintain productivity for each subspecies of island fox to sustainable levels.....	55

	Recovery Plan for Four Subspecies of Island Fox	
	2. Manage captive island fox populations for recovery.	69
	3. Establish island fox monitoring strategies.	73
V.	Implementation Schedule	77
VI.	Long-term Conservation Strategy	101
	1. Establish a mainland captive island fox population on which to conduct research to better understand fox behavior, ecology and reproduction, and disease and vaccine efficacy.....	101
	2. Establish, expand and continue island fox education and outreach programs.	103
	3. Assessing the demographic impact of other threats such as mortality from vehicle strikes, competition with feral cats, and emerging disease issues (e.g., ear cancer).	104
	4. Restoring island habitat.....	104
	5. Establishment of Conservation Agreements.	106
VII.	Literature Cited	107
VIII.	Appendices	125
	A. Appendix 1: Current and Former Members of the Island Fox Recovery Team and Primary Authors of this Recovery Plan.....	127
	B. Appendix 2: Discussion of the Risk-based Recovery Criterion ...	131
	C. Appendix 3: Guidelines and Recommendations on Vaccination Protocols, Collection of Health Data, and Future Research Needs Related to Fox Health.	137
	D. Appendix 4: Guidelines for Vaccination and Sampling of Captive and Wild Populations.....	143
	E. Appendix 5: Technical Analysis Request 2.1 - Development of Population Monitoring Plans for Free-Ranging Island Foxes	147

F. Appendix 6: Recommendations for Dog Entry and Exit from the Northern Channel Islands (Santa Cruz, Santa Rosa, and San Miguel) and the Navy Islands (San Clemente and San Nicolas) ... 151

G. Appendix 7: Guidelines for establishing a mainland population of island foxes with tasks, responsible parties, and time frames..... 157

H. Appendix 8: Identified research needs using a mainland captive island fox population..... 159

I. Appendix 9: List of Technical Analysis Requests (TARs)..... 161

J. Appendix 10: Glossary of Terms in the recovery plan for four subspecies of island fox 165

K. Appendix 11: Acronyms and Abbreviations in the Recovery Plan for Four Subspecies of Island Fox 171

L. Appendix 12: Summary of Comments 173

LIST OF TABLES

Table 1 Estimated number of wild adult and juvenile island foxes for each subspecies.....2

Table 2 Ownership and management of the California Channel Islands inhabited by island fox5

Table 3 Implementation schedule for the recovery plan for four subspecies of island fox.....78

LIST OF FIGURES

Figure 1 Channel Islands Land Ownership.....4

NOTE: Words and terms found with bold font throughout the Plan are defined in Appendix 10.

I. Background

A. INTRODUCTION

The island fox (*Urocyon littoralis*), a diminutive relative of the gray fox (*U. cinereoargenteus*), is endemic to the California Channel Islands. Island foxes inhabit six of the eight Channel Islands (San Miguel Island, Santa Rosa Island, Santa Cruz Island, Santa Catalina Island, San Nicolas Island, and San Clemente Island) and are recognized as a distinct subspecies on each of the six islands (Figure 1). Both morphologic and genetic distinctions support the classification of separate subspecies for each island (Collins 1993; Gilbert et al. 1990; Goldstein et al. 1999; Wayne et al. 1991a).

1. Legal Status

Four of the six island fox subspecies experienced catastrophic declines in the late 1990s, primarily due to golden eagle predation on the northern Channel Islands (San Miguel Island, Santa Rosa Island, and Santa Cruz Island) and canine distemper virus (CDV) outbreak (Table 1) on Santa Catalina Island (Timm et al. 2009). In June 2001, the Center for Biological Diversity petitioned the U.S. Fish and Wildlife Service (FWS) to list the four subspecies in catastrophic decline as endangered as defined by the Endangered Species Act of 1973, as amended (Act). In 2004, the FWS listed the San Miguel Island fox, Santa Rosa Island fox, Santa Cruz Island fox, and Santa Catalina Island fox as endangered (U.S. Fish and Wildlife Service 2004) pursuant to the Act. The remaining two subspecies, the San Nicolas Island fox (*U. l. dickeyi*) and San Clemente Island fox (*U. l. clementae*), did not experience the same type of population declines and thus, were not federally listed. However, concerns about the status of the San Clemente Island fox population prompted the Navy to enter into a Conservation Agreement with the U.S. Fish and Wildlife Service (FWS-LA-3287.1) and to undertake proactive measures to understand and mitigate potential impacts.

The California Fish and Game Commission listed the island fox as a rare species in 1971. All animals that had been determined to be rare on or before January 1, 1985 were designated as “threatened species” at that time. The IUCN (World Conservation Union) listed the entire species as Critically Endangered in 2001 (Sillero-Zubiri and Macdonald 2004).

The San Clemente Island fox (*U. l. clementae*) and the San Nicolas Island fox (*U. l. dickeyi*) are not federally listed, as their population numbers did not experience similar declines. However, all six subspecies are listed as threatened by the State of California. Therefore, the San Clemente Island fox (*U. l. clementae*) and the San Nicolas Island fox (*U. l. dickeyi*) subspecies are included for discussion in this plan.

Following the Federal listing of the island fox in 2004, the FWS considered designating critical habitat for the four listed subspecies. However, in its final

Recovery Plan for Four Subspecies of Island Fox

determination concerning critical habitat for the island fox (U.S. Fish and Wildlife Service 2005), the FWS concluded that no habitat met the definition of critical habitat in the Act and therefore, did not designate any critical habitat for any of the four subspecies. Critical habitat is defined in section 3(5)(A) of the Act in part as: the specific areas within the geographical area occupied by a species, at the time it is listed in accordance with the Act, on which are found those physical or biological features essential to the conservation of the species and that may require special management considerations or protection. The FWS did not designate any critical habitat for the island fox because: 1) the island fox is a habitat generalist and an opportunistic omnivore; 2) the primary reasons for the listing of the fox were predation and disease; and 3) prior to predation by golden eagles and the outbreak of disease, habitat did not appear to be a limiting factor despite human-induced habitat changes that have occurred. The FWS concluded that there are no specific areas where physical or biological features are essential to the conservation of the species and that may require special management considerations or protection; therefore, designating critical habitat would not be beneficial.

Table 1. Estimated number of wild adult and juvenile island foxes for each subspecies.

Island/ Subspecies	1994 Estimate ¹	1999/2000 Estimate ²	2012/2013 Estimate ³
San Miguel*	450	15	577
Santa Rosa*	1,780	15	894
Santa Cruz*	1,465	55	1354
Santa Catalina*	1,342	103	1852
San Clemente	1,003	535	1000
San Nicolas	520	452	341

* federally listed endangered subspecies.

¹source: Roemer et al. (1994).

²sources: Coonan et al. (2005a); Timm et al. (2002); Roemer et al. (2002)

³source: Coonan (2014); C. Boser, The Nature Conservancy, pers. comm. (2014); King and Duncan (2014); M. Booker, U.S. Navy, pers. comm. (2014); F. Ferrara, U.S. Navy, pers. comm. (2014); Hudgens and Garcelon (2014).


2. Affected Agencies, Landowners, and Partners

Islands inhabited by island foxes are owned by four major landowners: the National Park Service (NPS), the U.S. Navy (Navy), The Nature Conservancy (TNC), and the Santa Catalina Island Conservancy (CIC), all of whom have management authority for wildlife on their lands (Figure 1, Table 2). The NPS, TNC, and CIC manage the islands where the listed subspecies occur. The FWS guides the recovery planning process for the four listed island fox subspecies under the Act. Additionally, the State of California has regulatory authority over the island fox on non-Federal lands.

The bulk of the current knowledge regarding the evolution, ecology, behavior, and population biology of island foxes has been amassed by researchers from California institutions, including the University of California (Los Angeles, Davis, Santa Barbara, and Santa Cruz), California State University (Los Angeles), the Santa Barbara Museum of Natural History, and the non-profit Institute for Wildlife Studies (IWS).

In addition, researchers from a number of other U.S. institutions and organizations, including the Association of Zoos and Aquariums (AZA), Conservation Breeding Specialist Group (SSC/IUCN), the Honolulu Zoo, the Lincoln Park Zoo, New Mexico State University, the Santa Barbara Zoo, the Saint Louis Zoo, and the U.S. Geological Survey – Biological Resource Discipline (USGS-BRD) have contributed to the understanding of island fox natural history and recovery challenges. Collaboration with researchers has been, and will continue to be, critically important for island fox recovery efforts.

Figure 1. Channel Islands Land Ownership



Recovery Plan for Four Subspecies of Island Fox

Table 2. Ownership and management of the California Channel Islands inhabited by island fox

Island	Island size (km ²)	Island Management/Ownership ¹					
		Federal Agency		The Nature Conservancy	Santa Catalina Island Conservancy	Other Private Property	
		National Park Service	U.S. Navy			Santa Catalina Island Company ⁵	Other Private Landowners ⁶
San Miguel ^{2,3}	39	100% (manager)	100% (owner)				
Santa Rosa ^{2,3}	216	100%					
Santa Cruz ^{2,3,4}	243	24%		76%			
Santa Catalina ³	194				88%	11%	1%
San Clemente	149		100%				
San Nicolas	58		100%				

¹Both land owner and manager except as noted.

²Entire island within Channel Islands National Park boundary.

³Federally-listed endangered subspecies of island fox present.

⁴The Park and TNC cooperate fully on resources management and research issues via a cooperative agreement.

⁵The majority of this land is developed.

⁶These include home owners in the town of Avalon, the Wrigley Marine Science Institute run by University of Southern California, and Southern California Edison – the utility company that provides power, water, and gas for Catalina Island.

3. Integration of Conservation and Recovery Efforts

By 1999, island fox populations on the northern Channel Islands were considered to be in need of immediate conservation action (Coonan et al. 1998; Roemer 1999). The NPS convened a multi-disciplinary group of experts in 1999 (Island Fox Conservation Working Group) to evaluate available island fox status information and develop strategies to recover the island fox populations to viable levels. This was a loose affiliation of public agency representatives, landowners, conservancies, zoological institutions, non-profits, and academics concerned about conservation efforts for the island fox.

This group met annually from 1999 through 2004 and broadened its focus to include concerns about all six island fox subspecies. The working group served as a forum for information exchange and evaluation of recovery efforts. To address most issues, the group further divided into subject matter groups, such as management of wild populations, management of captive populations, island fox husbandry, veterinary issues, policy issues, and educational outreach needs. The group reported annually on the status of island foxes on all islands and listed findings regarding threats to the species and appropriate mitigation actions (Coonan and Rutz 2001, 2002, 2003; Coonan et al. 2004, 2005b).

In 2004, after four of the six subspecies were federally listed, the island fox Integrated Recovery Team (IRT; see Appendix 1) was established and incorporated the expertise of all 70+ individuals from the former working groups. At the same time, the island fox Recovery Coordination Group (RCG; see Appendix 1) was established with representatives from each of the land management agencies as well as additional canid experts. The FWS' goal for the RCG was both to draft the recovery plan using the knowledge and expertise of the IRT and to advise the FWS on immediate conservation needs. Tasks regarding management and recovery of island foxes were developed by the RCG and submitted to a task force for analysis; each task was formally referred to as a Technical Analysis Request (TAR). Each task force group was comprised of individuals from the larger IRT. In addition to incorporating the information into this recovery plan, the RCG forwarded the resulting analyses to the FWS, with recommendations for recovery actions needed immediately. The FWS then transmitted the analyses and recommendations to the land management agencies so that ongoing recovery efforts could take advantage of the best available information prior to the formal completion of an approved recovery plan. Since 2004, the land management agencies have received recommendations regarding releases from captivity and post-release monitoring, management and husbandry for captive populations, management of golden eagles, and establishment of a mainland captive population. The RCG also organized two population viability analysis meetings for the four federally listed subspecies, a monitoring workshop, and coordinated the annual island fox meetings in 2005 and 2006. In 2007, the

meeting marked a return to the format of the working group, focusing on status updates for all subspecies and results of recent research, as well as consideration of issues by small workgroups. The Island Fox Working Group has met annually through 2014.

B. BIOLOGICAL INFORMATION

1. Description and Taxonomy

A diminutive relative of the mainland gray fox, the island fox weighs approximately 1.8 to 3.0 kilograms (kg) (3 to 6 pounds (lb)) and stands approximately 30 centimeters (cm) (12 inches (in)) tall. The island fox is distinguished from the gray fox by its darker **pelage** and its smaller size (Collins 1982); most linear measurements of island foxes are 25 percent smaller than those of the gray fox. The dorsal coloration is grayish-white and black, and the base of the ears and sides of the neck and limbs are cinnamon-rufous in color (Moore and Collins 1995). The underbelly is a dull white, and the tail is conspicuously short. Island foxes display sexual size dimorphism, with males being larger and heavier than females (Collins 1982, 1993).

The island fox was first described as *Vulpes littoralis* by Baird in 1857 from the type locality on San Miguel Island, Santa Barbara County, California (Baird 1857). Merriam (1888, in Hall and Kelson 1959) reclassified the island fox into the genus *Urocyon* and later described island foxes from Santa Catalina, San Clemente, and Santa Cruz Islands as three separate species (*U. catalinae*, *U. clementae*, and *U. littoralis santacruzae*) (Merriam 1903). Grinnell et al. (1937) revised Merriam's classification, placing foxes from all islands under the species *U. littoralis* and assigning each island population a subspecific designation (*U. l. catalinae* on Santa Catalina Island, *U. l. clementae* on San Clemente Island, *U. l. dickeyi* on San Nicolas Island, *U. l. littoralis* on San Miguel Island, *U. l. santacruzae* on Santa Cruz Island, and *U. l. santarosae* on Santa Rosa Island). Recent morphological and genetic studies support this division of the *U. littoralis* complex into six subspecies, each restricted in range to a single island (Collins 1991a, 1993; Gilbert et al. 1990; Goldstein et al. 1999; Wayne et al. 1991a, 1991b).

2. Distribution, Evolution, and Genetics

Island foxes inhabit the six largest California Channel islands off the coast of southern California (San Miguel Island, Santa Rosa Island, Santa Cruz Island, San Nicolas Island, Santa Catalina Island, and San Clemente Island). Until recently, colonization of the islands by foxes was thought to have occurred sometime during the Pleistocene, before human presence on the islands. This was supported by genetic evidence that suggested all island foxes are descended from one colonization event (George and Wayne 1991), possibly from chance, over-water

Recovery Plan for Four Subspecies of Island Fox

dispersal by rafting on floating debris (Moore and Collins 1995). However, recent re-dating of island fox fossils indicates that the earliest known island fox remains are no more than 6,000 years old (Rick et al. 2009). This raises the possibility that Native Americans transported foxes from the mainland to the northern islands sometime after human contact (10,000-13,000 years ago).

Island foxes may have reached the southern Channel Islands (San Nicolas, San Clemente, and Santa Catalina Islands) much more recently (2,200 to 3,800 years ago), and were most likely introduced to these islands by Native Americans as pets or semi-domesticates (Collins 1991a, 1991b). Island fox remains recently recovered from San Nicolas Island extend this time period to approximately 5,200 years before present (Vellanoweth 1998).

Morphologically, the species exhibits inter-island variability in size, nasal shape and projection, and the number of tail vertebrae (Collins 1982). Genetic evidence supports the separation of the species into six distinct subspecies, and confirms the pattern of dispersal, though perhaps not the timing, suggested by archeology and geology. A study of genetic variability in **DNA restriction fragments** in island foxes (Gilbert et al. 1990) revealed that inter-island variability was greater than intra-island variability. **Phylogeny** based upon restriction fragment variability supports the geological evidence for the sequence of isolation of each island, and each population, as rising sea levels separated *Santarosae* into the northern Channel Islands. Santa Cruz Island separated from the other northern islands first, about 11,500 years ago, followed by the separation of San Miguel Island and Santa Rosa Island about 9,500 years ago. Together with the fossil record, restriction fragment evidence indicates that San Clemente Island was the first southern Channel Island colonized, probably by immigrants from San Miguel Island. Dispersal then occurred from San Clemente Island to San Nicolas and Santa Catalina Islands.

Island forms generally have less genetic variability than their mainland counterparts. Mainland gray foxes were found to be more variable in morphology, **allozymes**, mitochondrial DNA, and hypervariable nuclear DNA than island foxes (Goldstein et al. 1999; Wayne et al. 1991a). The island fox populations with the fewest numbers of individuals, San Miguel Island and San Nicolas Island, showed the least genetic variability, and the San Nicolas Island population was actually monomorphic (showing no variation) in allozyme, hypervariable minisatellite and microsatellite DNA, and mitochondrial DNA, which is highly unusual among mammals. This lack of variability could be attributed either to extensive inbreeding or to **bottlenecking** resulting from low population densities (George and Wayne 1991). On San Miguel and San Nicolas Islands, the species has apparently existed for thousands of years at low **effective population sizes** (150 to 1000), with low genetic variability (Wayne et al. 1991a,

1991b). The Santa Rosa Island and San Miguel Island populations have been shown to be closely related (Wayne et al. 1991b).

Recently, Aguilar et al. (2004) found considerable variation at the major histocompatibility complex (MHC) in San Nicolas Island foxes, which contain genes that code for disease resistance and kin recognition. Modeling by the authors suggests that the pattern of MHC and neutral marker variation in San Nicolas Island foxes was caused by an extreme bottleneck (a decline to fewer than 10 animals) in the past 10 to 20 generations.

Recently, genetic relatedness among individuals was determined for the San Miguel Island and Santa Rosa Island captive populations (Gray et al. 2001; Gray 2002). Analysis of island fox blood samples from 1988 and from the captive population in 2001 indicated that the level of variation in island foxes on the two islands had declined since 1988. During that time period, there was a reduction in the number of **alleles** at some **loci** and, at some loci, a complete loss of **polymorphism**. However, there are no apparent deleterious effects of inbreeding in island foxes (Coonan et al. 2010).

3. Habitat Use and Food Habits

The island fox is a habitat generalist, occurring in all natural habitats on the Channel Islands, although it prefers areas of diverse topography and vegetation (von Bloeker 1967; Laughrin 1977; Moore and Collins 1995). Island foxes occur in valley and foothill grasslands, southern coastal dune, coastal bluff, coastal sage scrub, maritime cactus scrub, island chaparral, southern coastal oak woodland, southern riparian woodland, Bishop and Torrey pine forests, and coastal marsh habitat types. Crooks and Van Vuren (1996) found island foxes to prefer fennel (*Foeniculum vulgare*) and to avoid ravines and scrub oak patches on Santa Cruz Island. Island foxes may use non-native grasslands less than other habitats, even though insect prey is abundant in grasslands, because grasslands are denser and may be more difficult to forage in (Roemer and Wayne 2003). Also, low vegetation types such as grasslands may render island foxes more vulnerable to aerial predators (Roemer 1999).

Island foxes are omnivores and forage opportunistically, eating a wide variety of seasonally available plants and animals (Collins 1980; Collins and Laughrin 1979; Crooks and Van Vuren 1995; Kovach and Dow 1981; Laughrin 1973, 1977; Moore and Collins 1995). Island foxes feed on a wide variety of insect prey, such as grasshoppers, crickets, and katydids (Crooks and Van Vuren 1995; Moore and Collins 1995) and Jerusalem crickets (*Stenopelmatus fuscus*) when seasonally available (Moore and Collins 1995).

Island foxes prey on native deer mice (*Peromyscus maniculatus*) on all islands and also likely prey upon introduced house mice (*Mus musculus*) on Santa

Catalina Island and introduced rats (*Rattus rattus*) on Santa Catalina, San Miguel, and San Clemente Islands. Deer mice may be especially important prey during the breeding season, because they are large, energy-rich food items that adult foxes can bring back to their growing pups (Garcelon et al. 1999). In addition to small mammals, island foxes prey on ground-nesting birds such as horned larks (*Eremophila alpestris*) and western meadowlarks (*Sturnella neglecta*). Less common in the diet are amphibians, reptiles, and the carrion of marine mammals (Collins and Laughrin 1979). Island foxes feed on a wide variety of native plants, including the fruits of *Arctostaphylos*, *Comarostaphylis*, *Heteromeles*, *Opuntia*, *Prunus*, *Rhus*, *Rosa*, *Solanum*, and *Vaccinium* (Moore and Collins 1995). San Miguel Island foxes rely more on the fruits of sea-fig, *Carpobrotus chilensis* (Collins 1980; Crowell 2001). A comprehensive treatment of island fox diet is found in Moore and Collins (1995).

The island fox is a **docile canid**, exhibiting little fear of humans in many instances. Although primarily nocturnal, the island fox is more **diurnal** than the mainland gray fox (Collins and Laughrin 1979; Crooks and Van Vuren 1995; Fausett 1993), possibly a result of historical absence of large predators and freedom from human harassment on the islands (Laughrin 1977).

4. Social Organization and Reproduction

Island foxes generally have smaller territories, exist at higher densities, and have shorter dispersal distances than mainland fox species, characteristics typical of vertebrate populations on islands (Roemer 1999; Roemer et al. 2001a). Island fox home range size and configuration are dependent on landscape features, resource distribution, fox population density, habitat type, season, and sex of the animal (Fausett 1982; Laughrin 1977; Crooks and Van Vuren 1996; Thompson et al. 1998). Recorded home-range estimates range from 0.24 square kilometer (km²) (0.09 square mile (mi²)) in mixed habitat (Crooks and Van Vuren 1996) and 0.87 km² (0.34 mi²) in grassland habitat (Roemer 1999) on Santa Cruz Island, to 0.77 km² (0.3 mi²) in canyons on San Clemente Island (Thompson et al. 1998). Island fox territory size on Santa Cruz Island varied from 0.15 to 0.87 km² (0.06 to 0.34 mi²) and averaged 0.55 km² (0.21 mi²) during a period of moderate to high fox density (7 island foxes per km² [18 per mi²]) (Roemer et al. 2001a).

Research on Santa Cruz Island found that island foxes, like most foxes, exist as socially monogamous pairs occupying discrete territories (Roemer et al. 2001a). Territory configuration changed after the death and replacement of paired male foxes, but not after the death and replacement of paired females or juveniles, indicating that adult males are involved in territory formation and maintenance. Despite being socially monogamous and territorial, island foxes are not necessarily genetically monogamous. On Santa Cruz Island, 4 of 16 offspring whose parents were identified by paternity analysis were a result of extra-pair

fertilizations (Roemer et al. 2001a). All extra-pair fertilizations occurred between foxes from adjoining territories.

Island fox courtship activities occur from late January to early March (Moore and Collins 1995). In the island fox captive breeding facility on San Miguel Island, copulations were observed during the first 2 weeks of March 2000, and copulation for the successful pairs likely occurred between mid-February and early March (Coonan and Rutz 2001, 2002). Young are born from early to late April after a gestation period of approximately 50 to 53 days. Births occurred in the NPS' island fox captive breeding facilities from April 1 to April 25 (Coonan et al. 2010).

Island foxes give birth to their young in simple dens, under shrubs, or in the sides of ravines (Laughrin 1973). Litter size ranges from one to five (Moore and Collins 1995); mean litter size for 24 dens on Santa Cruz Island was 2.17 (Laughlin 1977). The average number of foxes produced in 51 litters in captivity from 1999 to 2004 was 2.4 (Coonan et al. 2005b). Like other fox species, island foxes exhibit biparental care (care by both parents), evidenced by the capture of adult male foxes in the same traps as pups and observations of adults and known offspring foraging together (Garcelon et al. 1999; Roemer 1999). By 2 months of age, young foxes spend most of the day outside the den and will remain with their parents throughout the summer. Some pups disperse away from their natal territories by winter, although others may stay on their natal territories into their second year.

Although island foxes are physiologically capable of breeding at the end of their first year (Laughrin 1977), most breeding involves older animals. Coonan et al. (2000) found that only 16 percent of 1 to 2 year old females bred over a 5-year period on San Miguel Island, in contrast to 60 percent of older females. Roemer (1999) found yearling females to have lower fertility than older females on Santa Cruz Island. However, females reintroduced from captive facilities on San Miguel Island have produced litters at 1 year of age (Coonan et al. 2010).

Prior to the catastrophic population declines of the 1990s, adult island foxes were reported to live an average of 4 to 6 years (Moore and Collins 1995); Coonan et al. (1998) recorded eight individuals on San Miguel Island that lived 7 to 10 years in the wild.

5. Mortality Sources and Population Dynamics

In an effort to describe the basic biology and life history characteristics of the island fox, the following section describes current and historic sources of mortality to the island fox, as well as the population dynamics of the species. In many cases, there is an overlap between the sources of mortality described here and the current threats to the species. However, the specific threats to the species

are considered more fully in the section of this document entitled, “Threats to the Species.”

(a) Golden Eagles. Predation by golden eagles drove the island fox subspecies on San Miguel, Santa Cruz, and Santa Rosa Islands to near extinction in the late 1990s (Roemer 1999; Roemer et al. 2001b; Coonan et al. 2005c). Golden eagle predation has continued to be the primary mortality factor for foxes on the northern Channel Islands.

The extirpation of bald eagles (*Haliaeetus leucocephalus*) from the Channel Islands as a result of dichlorodiphenyltrichloroethane (DDT) may have facilitated golden eagle colonization. Bald eagles historically bred on the islands and aggression by breeding bald eagles may have discouraged foraging golden eagles from establishing residence. Bald eagles are represented in the prehistoric fossil record of the northern Channel Islands (Guthrie 1993) and bred there until 1960, when nest failures, as a result of DDT contamination, extirpated them from the northern Channel Islands (Kiff 1980). The northern Channel Islands (Anacapa, Santa Cruz, Santa Rosa, and San Miguel Islands) likely supported more than 14 pairs of bald eagles before their decline (Kiff 1980).

Bald eagles normally rely on marine resources as a food resource base (Newsome et al. 2010); while golden eagles traditionally focus on terrestrial species (Collins and Latta 2009). Additionally, on much of the northern Channel Islands, historic sheep grazing changed the predominant vegetation from shrub to non-native grasslands, which offered foxes much less cover from aerial predators.

Except for golden eagles, red-tailed hawks (*Buteo jamaicensis*) are the only other confirmed avian predator of island foxes (Laughrin 1980; Moore and Collins 1995), and likely only prey on island fox pups, not on adult island fox. There have been unconfirmed historical reports of predation by bald eagles, but there is no current or recent evidence to suggest that foxes are a dominant prey item of bald eagles. Island fox remains have been found in bald eagle nests; however, it is not known whether the individuals were depredated or scavenged (Collins et al. 2005).

(b) Canine Distemper Virus. A CDV outbreak was the cause of decline for the island fox subspecies on Santa Catalina Island (Timm et al. 2009). This disease remains a potential mortality factor for island foxes and is capable of causing a catastrophic decline (Timm et al. 2000, 2002; Kohlmann et al. 2005). Recent **serological** surveys recorded the presence

of antibodies reactive against CDV in wild foxes on all islands (Clifford et al. 2006, Coonan et al. 2010), suggesting that exposure to CDV or a similar **morbillivirus** has occurred in all island fox subspecies, with survival of many infected individuals. A natural CDV-like morbillivirus is thus likely circulating within island fox populations, and confers some immunity to CDV. The endemic CDV-like strain is apparently less virulent than other strains of CDV, and has no apparent effect on fox populations (Coonan et al. 2010).

(c) Other Factors. Additional mortality factors for island foxes include, but are not limited to, vehicle strikes on roads, other diseases, and parasites. At least one case of island fox mortality due to shooting by an unknown person(s) was confirmed in 2007 on Santa Catalina Island (King and Duncan 2008). Collision with motor vehicles remains a threat to island foxes on San Nicolas and San Clemente Islands (Moore and Collins 1995) and on Santa Catalina Island (Munson 2010). On Santa Catalina Island, annual averages of four foxes per year were killed by vehicles from 2002 to 2007. More than 30 foxes are killed by vehicles annually on San Clemente Island (Garcelon et al. 2008). On San Nicolas Island, an average of 17 foxes was killed by vehicles annually between 1993-2013, with 22 foxes killed by vehicles in 2013 (F. Ferrara, U.S. Navy, pers. comm. 2014). This average includes only foxes that were killed instantly. We believe it is likely that some foxes were hit and later succumbed to their injuries, or that there are juveniles who did not survive following the death of the mother. Because we do not have a method to document this type of mortality, the actual annual mortality due to vehicles is likely higher. However, vehicle speed limits have been lowered, speed limits are enforced, road shoulders are mowed, and outreach and awareness have increased in an attempt to reduce vehicle related mortality on San Nicolas Island.

Island foxes have shown previous exposure to infectious agents such as canine parvovirus, canine adenovirus, canine corona virus, canine herpes virus, and toxoplasmosis (Garcelon et al. 1992; Roemer et al. 2001b; Clifford et al 2006), but disease resulting from these infectious agents was not found to be a mortality factor until CDV and toxoplasmosis was confirmed in a dead fox on Santa Catalina Island in 1999 (Munson 2010). Different island fox subspecies have been exposed to multiple **serovars** of **Leptospira** in the past, but leptospirosis was not a mortality cause until 2010, when a *Leptospira* outbreak on Santa Rosa Island was associated with the mortalities of two radio-collared foxes (Coonan and Guglielmino 2012).

Recovery Plan for Four Subspecies of Island Fox

The recent finding of ear tumors in Santa Catalina Island foxes, confirmed to be a source of mortality in wild foxes, is of high enough frequency to be considered a concern (Coonan et al. 2010). The first case of this ceruminous gland carcinoma, a rare but aggressive malignant tumor, was diagnosed in 2001. The tumors are primarily confined to the ears of the animals, but in some cases spread to the head and neck region and eventually may **metastasize** (Munson 2010). The disease has been found in all Santa Catalina Island fox age groups, except pups. In 2004, veterinarians found that a high proportion of the adults either had these tumors or showed signs of tissue changes that are possible precursors to tumor development (Munson 2010). The tumors are associated with severe otitis and infections of *Otodectes* (ear mites). *Otodectes* are present in island fox populations on other islands; however, the tumors only occur in Santa Catalina Island foxes.

Parasites have not been confirmed as a mortality source, except for rare cases of complications from *Spirocerca* (nematode) infection (Munson 2010). In a species-wide survey, *Spirocerca* was found in a high prevalence of **necropsied** island foxes, but in most cases appeared to have little effect on individual health (Munson 2010). Preliminary genetic analysis and the location of lesions suggest that the *Spirocerca* found in island foxes may be a different species than *S. lupi*, which occurs in domestic dogs and other North American carnivores on the mainland. Currently, *Spirocerca* is not a major health concern for most island foxes. However, if island foxes are to be brought to the mainland, efforts should be made to prevent transmission of *Spirocerca* from island foxes to mainland carnivores and vice versa.

Heavy parasite infections by hookworms (*Uncinaria stenocephala*) and a lungworm (*Angiocaulus gubernaculatus*) may have contributed to two mortalities in the San Miguel Island fox subspecies (Coonan et al. 2005c). *Angiocaulus* is not found in other island fox subspecies (Faulkner et al. 2001). Unusual infection by acanthocephalans (spiny-headed worms) was detected in San Miguel Island foxes in 2011-2012, and may have contributed to the death of at least one radio-collared fox (Coonan 2013).

(d) Population Dynamics. Even in the absence of catastrophic mortality sources, island fox populations may have fluctuated markedly over time (Laughrin 1980). Residents of Santa Cruz Island occasionally noted periods of island fox scarcity and abundance. Santa Catalina Island fox population levels were low in 1972 and again at low density in 1977 (Laughrin 1980). However, by 1994 the adult Santa Catalina Island fox population was estimated at over 1,300 individuals (Roemer et al. 1994). The San Nicolas Island fox population was considered to be at very low

densities in the early 1970s (Laughrin 1980), and may have reached approximately 500 individuals by 1984 (Kovach and Dow 1985, as cited by Wayne et al. 1991b).

Demographic analysis indicated that island fox survival was positively related to the previous year's El Niño Southern Oscillations (ENSO) events in the drier southern islands and negatively related to current and previous year's ENSO events in the wetter northern islands (Bakker et al. 2009; see Appendix 2). Thus, indirect evidence suggests an effect of climate on island fox survival.

C. HISTORICAL POPULATION STATUS AND OBSERVED DECLINES OF ISLAND FOX POPULATIONS

The four federally listed island fox subspecies (San Miguel Island fox, Santa Rosa Island fox, Santa Cruz Island fox, and Santa Catalina Island fox) all experienced precipitous population declines in the latter half of the 1990s (see Table 1) (Coonan et al. 2000, 2005c; Roemer 1999; Roemer et al. 2001b; Timm et al. 2000). Island fox populations on San Miguel, Santa Rosa, and Santa Cruz Islands declined by 90 to 95 percent and, prior to removal of foxes from the wild for captive breeding, were estimated to have a 50 percent chance of extinction over 5 to 10 years (Roemer 1999; Roemer et al. 2001b). Thus, by 1999 researchers considered island fox subspecies on the northern Channel Islands to be critically endangered (Coonan et al. 1998; Roemer 1999), as was the Santa Catalina Island subspecies by 2000 (Timm et al. 2000).

The decline of island foxes in the northern Channel Islands is considered a consequence of **hyperpredation** (Roemer et al. 2001b). The presence of non-native species (feral pigs on Santa Cruz Island and mule deer and elk on Santa Rosa Island) and the absence of bald eagles enabled golden eagles to colonize the islands successfully and prey heavily on island foxes, which evolved in the absence of predators. Fox carcass characteristics that indicated golden eagle predation included **evisceration**, **degloving**, spinal cord separation, and the presence of golden eagle feathers at carcass sites (Roemer et al. 2001b, Coonan et al. 2005c). Additional evidence of eagle predation included an increase in golden eagle sightings on the northern Channel Islands, discovery of nesting golden eagles (previously unknown from the Channel Islands), and the presence of pig and island fox remains in golden eagle nests (Collins and Latta 2006). A mathematical model of hyperpredation showed that pigs would have been a necessary food source to support a large, resident golden eagle population (Roemer 1999; Roemer et al. 2001b, 2002) and that as few as six golden eagles could have driven the island fox populations to the lows recorded during the 1990s. In 1999, prior to golden eagle removal efforts, there were estimated to be as many as 27 golden eagles on the northern Channel Islands (Latta et al. 2005).

Based on an analysis of extinction likelihood, Roemer (1999) concluded that if mortality and reproduction continued at rates similar to those observed just prior to intervention, both the San Miguel Island fox and the Santa Cruz Island fox, and likely the Santa Rosa Island fox, would decline to extinction. Successful long-term suppression of golden eagles would likely require removal of the non-native prey base (feral pigs removed from Santa Cruz Island and deer and elk removed from Santa Rosa Island), as well as the successful restoration of bald eagles to the northern Channel Islands (Coonan 2003; Coonan et al. 2005a).

1. San Miguel Island

Laughrin surveyed San Miguel Island foxes in the early 1970s (Laughrin 1973). Trap success (number of fox captures per available trap) was high (43 percent) and Laughrin concluded that island fox populations were stable at 2.7 island foxes per km² (7 per mi²). In the late 1970s, the San Miguel Island fox density averaged 4.6 island foxes per km² (11.9 per mi²) and the island-wide population was estimated to be 151 to 498 individuals (Collins and Laughrin 1979). In 1993, the NPS began a long-term monitoring program for San Miguel Island foxes, using standardized mark-recapture methods (Roemer et al. 1994). Adult density on two grids was 7.8 island foxes per km² (20.2 per mi²) and 8.0 island foxes per km² (20.7 per mi²) in 1993, and the island-wide estimate was about 300 foxes (Coonan et al. 1998). A third grid was added the following year. That grid, the Dry Lakebed grid, recorded the highest density then known for island foxes in 1994 (15.9 island foxes per km² [41.2 per mi²]) and the island-wide estimate rose to 450 adult foxes.

Annual monitoring documented a substantial decline in the San Miguel Island fox population between 1994 and 1999 (Coonan et al. 1998; Coonan et al. 2000; Coonan et al. 2005c), when the estimated island-wide population steadily and sharply declined, falling to only 15 adults in 1999. In 1999, the NPS brought 14 San Miguel Island foxes into captivity (4 males and 10 females) to initiate a captive breeding program. The only known individual left in the wild at that time, a previously radio-tagged female (Coonan et al. 2005c), was brought into captivity in 2003, but died in December of that year. A necropsy indicated the fox had healed scars on the intercostal muscles between her ribs, suggesting she had survived a predation attempt (Coonan et al. 2004).

The cause of the San Miguel Island fox population decline was almost certainly predation by golden eagles (Roemer 1999; Roemer et al. 2001b; Coonan et al. 2005c). During a radio-telemetry study in 1998 and 1999, six of eight collared foxes died within 4 months, four of which were preyed upon by golden eagles (Coonan et al. 2005c).

2. Santa Rosa Island

Laughrin (1980) surveyed the Santa Rosa Island fox population in 1972, reporting a trap success rate of 50.0 percent and a density of 4.2 island foxes per km² (10.9 per mi²), which coincides with an island-wide population estimate of 898 individuals. No other previous data are available for the Santa Rosa Island fox population except for surveys conducted from 1998 to 2000. Based on island size, Roemer et al. (1994) estimated the island-wide population to be 1,780 adult foxes. More recent trapping data as well as anecdotal evidence suggest that the Santa Rosa Island fox population experienced a decline similar to that of the Santa Cruz Island fox and San Miguel Island fox (Roemer et al. 2001b; Coonan et al. 2005a). Roemer (1999) reported that during 132 trap nights in 1998, only 9 individuals were captured (10 total fox captures), for a trap success rate of 7.5 percent. In 2000 and 2001, the NPS brought the remaining 15 wild Santa Rosa Island foxes into captivity for captive breeding (Coonan and Rutz 2002). No further fox sign was seen on Santa Rosa Island after May 2001 (Coonan et al. 2005a).

Given the proximity of Santa Rosa Island to Santa Cruz and San Miguel islands, the concurrent timing of the population decline, and the presence of golden eagle nests, golden eagle predation was the likely cause of the decline of the Santa Rosa Island fox (Roemer 1999; Roemer et al. 2001b). Golden eagle breeding was confirmed on the island in 2003 (Latta et al. 2005). Both currently and formerly active golden eagle nests were found in two eagle breeding territories, Trap Canyon and Trancion Canyon. Some nests were used in successive years. Layering of prey remains in the nests indicated that golden eagles had been successfully breeding (fledging young) on Santa Rosa Island since as early as 1997, and island fox remains in the lower layers confirmed predation of eagles upon island foxes (Latta et al. 2005; Collins and Latta 2006). The examined nests on Santa Rosa Island did not contain feral pig remains, indicating that the examined nests were established after pigs were eradicated from the island (post 1992). Examination of golden eagle nests on Santa Rosa Island found remains of island foxes as well as mule deer fawns, island spotted skunks (*Spilogale gracilis amphialus*), and many birds including ravens (*Corvus corax*), mallards (*Anas platyrhynchos*), barn owls (*Tyto alba*), and California quail (*Callipepla californica*) (Latta 2001; Collins and Latta 2006). The prevalence of mule deer fawns in the prey remains underscored their importance for golden eagle breeding on Santa Rosa Island. Golden eagles are also known to eat carrion and carcasses from the annual cull of deer and elk that occurred in November and December, supporting wintering golden eagles. In addition, fawn availability in the spring allowed nesting eagles to successfully fledge young. The non-native deer and elk (*Cervus elaphus*) were managed by the former owners of Santa Rosa Island for a sport-hunting operation. In 2011, large scale efforts to remove the non-native mule deer and elk on Santa Rosa Island were implemented as part of a court

settlement (National Parks and Conservation Association v. Kennedy, United States District Court for the Central District of California, No. CV 96-7412-WJR (RNBx) to remove the deer and elk by the end of 2011) (NPS 1998). Monitoring in 2013 verified the success of these efforts and all elk and all but a few deer remained at the end of 2013. The NPS and a cooperator plan to remove any remnant deer.

3. Santa Cruz Island

Santa Cruz Island is the largest of the Channel Islands and historically supported high densities of island foxes (Laughrin 1973). An early population estimate for the Santa Cruz Island fox was believed to be no more than 3,000 individuals (Laughrin 1971). Between 1973 and 1977, Laughrin (1980) estimated the Santa Cruz Island fox population to be 1,968 individuals based on an average density of 7.9 island foxes per km² (20.5 per mi²). However, island-wide population estimates extrapolated from annual Santa Cruz Island fox densities on two grids in 1993 suggest the population decreased from a high of approximately 1,000 to 1,300 foxes (which is believed to be a more accurate estimate than previous population estimates; Roemer et al. 1994) to an estimated 55 adults in 2001 (Dennis et al. 2001, 2002), while trapping efficiency was 2.9 percent in 1998 (Roemer 1999).

All available evidence indicates the decline of the Santa Cruz Island fox was caused by golden eagle predation (Roemer 1999; Roemer et al. 2001b). From August 1993 to September 1995, golden eagles were linked to 19 of 21 fox mortalities on the western end of Santa Cruz Island. Examination of golden eagle nests on Santa Cruz Island found remains of island foxes as well as island spotted skunks (*Spilogale gracilis amphialus*), feral pigs (*Sus scrofa*), and many birds including ravens (*Corvus corax*), mallards (*Anas platyrhynchos*), barn owls (*Tyto alba*), and California quail (*Callipepla californica*) (Latta 2001; Collins and Latta 2006). Santa Cruz Island foxes were brought into captivity for breeding in 2002 to provide a “safety net” against extinction and offspring to supplement the wild population.

4. Santa Catalina Island

Santa Catalina Island fox numbers appear to have fluctuated widely over the past 30 years. During surveys from 1972 to 1977, Laughrin (1980) caught only 2 individuals, and trap success was 3.0 percent, although Propst (1975) caught 55 individuals with a trap success rate of 11 percent. Between 1988 and 1991, average density increased, ranging from 2.6 island foxes per km² (6.7 per mi²) to 12.7 island foxes per km² (32.9 per mi²) (Garcelon et al. 1991). The Santa Catalina Island fox population increased to an estimated 1,342 foxes by 1994 (Roemer et al. 1994).

Recovery Plan for Four Subspecies of Island Fox

The Santa Catalina Island fox population experienced a catastrophic decline of more than 90 percent from 1999 to 2000. Sightings of dead and dying foxes, retrieval of a fox carcass infected with CDV, and confirmation of antibodies against CDV in live foxes suggest this decline was likely due to the introduction of canine distemper to the Santa Catalina Island fox population (Timm et al. 2000). The outbreak occurred principally on the large, eastern portion of the island, which is separated by a narrow isthmus from the smaller western end. Trap success on the eastern side of the island dropped from 26.0 percent in 1998 to 1.0 percent in 1999 and 2000, while remaining stable at approximately 36.0 percent on the western portion. The Santa Catalina Island fox population was reduced to perhaps 100 foxes by 2000, mostly on the west end (Timm et al. 2002).

Currently, there is considerable concern about the high rate of ceruminous gland carcinoma (ear tumors) in Santa Catalina Island foxes and how it might affect the recovery and long-term viability of the population (Coonan et al. 2010; W. Vickers, Institute for Wildlife Studies. pers. comm. 2014).

Santa Catalina Island has a human population of approximately 4,000, a large population of domestic dogs (*Canis familiaris*), and a considerable number of domestic and feral cats (*Felis catus*). Santa Catalina Island also has the highest degree of human activity and accessibility of any of the Channel Islands (over 1,000,000 visitors per year).

5. San Clemente Island

The earliest density estimate on San Clemente Island was 4.2 island foxes per km² (10.9 per mi²) (Laughlin 1973). Wilson (1976) recorded fox density to be 5.7 island foxes per km² (14.8 per mi²) and island-wide population size to be 2,000 foxes. The San Clemente Island fox population has been monitored annually since 1988 (except for the years 1998 and 2006). Population sampling between 1988 and 1991 found densities of 4.8 island foxes per km² (12.4 per mi²) to 9.1 island foxes per km² (23.6 per mi²) (Garcelon et al. 1991). Roemer et al. (1994) found similar densities and estimated an island-wide population of 1,003 foxes. However, Garcelon (1999) estimated that the San Clemente Island fox population ranged between 506 and 875 individuals from 1989 to 1999.

Data from grid trapping indicate that from 1990 to 2000 the San Clemente Island fox population experienced a gradual decline from over 800 foxes to fewer than 600, but the population stabilized, if not increased thereafter, and as of 2004, the population estimate was over 750 foxes (Garcelon 1999, Schmidt et al. 2005a). Densities in 2004 ranged from 2.4 island foxes per km² (6.2 per mi²) in grassland to 12.6 island foxes per km² (32.6 per mi²) in scrub/dune habitats (Schmidt et al. 2005a). Concerns about the status of the San Clemente Island population prompted the Navy and the FWS to enter into a Conservation Agreement (2003)

and undertake proactive measures to understand and mitigate potential threats. Since 2004, the population has increased, and in 2013, the adult population estimate reached 1002 (Bridges et al. 2014).

The causes of population fluctuations of foxes on San Clemente Island are unknown; however, predator management activities to protect the federally endangered San Clemente loggerhead shrike might have been a contributing factor. As part of this program, the Navy initially focused on non-native predators (cats and rats), but in 1999 implemented control measures for native predators as well, including the San Clemente Island fox (Department of the Navy 1999). In 1999, the Navy euthanized 13 foxes and relocated 15 to zoos (Garcelon 1999). After 1999, San Clemente Island foxes in San Clemente Island loggerhead shrike breeding territories were shock-collared or captured and held in captivity for the duration of the San Clemente Island loggerhead shrike breeding season. Shock collaring and removal of San Clemente Island foxes to captivity were suspended in 2003. Accidental poisoning from rodenticides used for pest management has also caused San Clemente Island fox mortalities (Munson 2010); however, there are no records of island foxes having been poisoned on other islands. Feral cats exist on the island in high densities (Phillips and Schmidt 1997) and could be competing with San Clemente Island foxes for prey and may expose them to pathogens.

6. San Nicolas Island

Laughrin (1980) reported a density of 0.12 San Nicolas Island fox per km² (0.3 per mi²) in 1977, which suggested an island-wide estimate of only 7 animals. Laughrin's reported low trap success rate (4.7 percent) is comparable to the low trap success rates on San Miguel and Santa Rosa Islands at the latter end of the population declines recorded there in the late 1990s. The San Nicolas Island fox population declined to fewer than 30 individuals in the mid-1970s, coincident with the termination of a supplemental feeding program (Laughrin 1980) and an increase in the feral cat population on the island (Kovach and Dow 1982). Using genetic data, Aguilar et al. (2004) estimated that the population had declined to fewer than 10 individuals during the bottleneck. Following the initiation of a feral cat eradication program in 1980, San Nicolas Island fox numbers increased from approximately 120 to 600 foxes in 4 years (Kovach and Dow 1985). Grid densities in 2004 ranged from 8.4 island foxes per km² (21.8 per mi²) to 20.1 island foxes per km² (52 per mi²), and the island-wide population was estimated to be 548 foxes (Garcelon and Schmidt 2005). The island-wide population estimate in 2009 was 619 individuals; in 2012 it declined to 460 individuals; and in 2013, declined again to 341 individuals (F. Ferrara, U.S. Navy, pers. comm. 2014; Hudgens and Garcelon 2014).

D. THREATS TO THE SPECIES

Section 4(a)(1) of the Act identifies five major categories of threats, which are considered when a species is listed. These are (a) the present destruction, modification, or curtailment of its range, (b) overutilization for commercial, recreational, scientific, or educational purposes, (c) disease or predation, (d) the inadequacy of existing regulatory mechanisms, and (e) other natural or manmade factors affecting its continued existence. Each of these potential categories of threats is analyzed below.

Factor A: Present or Threatened Destruction, Modification, or Curtailment of the Species' Habitat or Range

Although it is difficult to quantify the effects of past habitat loss and/or alteration on the status of island foxes, habitat on all islands occupied by island foxes has been heavily affected by livestock grazing, cultivation, and other disturbances. A century and a half of overgrazing by non-native herbivores (e.g., sheep, goats, deer, elk, cattle, pigs, and horses) has resulted in substantial impacts to the soils, topography, and vegetation of the islands (Johnson 1980; Coblenz 1980; Clark et al. 1990; Peart et al. 1994; O'Malley 1994). Much of the native coastal sage scrub, chaparral, and oak woodland habitats have been replaced by other vegetation, especially non-native annual grasses (Brumbaugh 1980; Clark et al. 1990; Klinger et al. 1994). Annual grasslands constitute less preferred habitat for island foxes (Laughrin 1977; Roemer and Wayne 2003) and do not provide cover from predators such as golden eagles (Roemer 1999; Roemer et al. 2001b; Coonan et al. 2005c). In 1987, the California Department of Fish and Game (CDFG), now recognized as California Department of Fish and Wildlife (CDFW), recommended that the island fox retain its classification as threatened under State law because of continued habitat degradation from herbivorous mammals on Santa Rosa, Santa Cruz, Santa Catalina, and San Clemente Islands (CDFG 1987). Since that time, non-native species removal programs have eradicated or reduced the introduced herbivore populations on many of the Channel Islands, including a recent complete removal of over 5,000 feral pigs from Santa Cruz Island (Parkes et al. 2010), and the removal of all elk and all but a few deer from Santa Rosa Island, resulting in an island that is essentially ungulate free (Coonan, pers. comm. 2014). On Catalina Island, all but one pig and three female goats have been removed resulting in an island that is essentially pig and goat free (Garcelon et al. 2005, King, Santa Catalina Island Conservancy, pers. comm. 2014). On San Nicholas Island, feral cats were recently removed (Hanson 2012).

Although some plant species have increased in number following the removal of non-native herbivores and omnivores from the islands, other aspects of recovery of the native habitats can be slow (Hochberg et al. 1979). In particular, community composition can be altered by the spread of non-native plants that

Recovery Plan for Four Subspecies of Island Fox

were able to gain a foothold during the period of disturbance. These non-native species continue to invade and modify island fox habitat resulting in lower diversity of vegetation, less diverse habitat structure, and reduced food availability. At present, habitat degradation by herbivores continues only on Santa Catalina Island, primarily by bison. However, effects from past grazing activity, such as loss of topsoil or spread of non-native species, continue to occur on all islands. Habitat modification also occurs as a result of facilities or recreational development.

Although it is possible that these habitat changes may have affected island foxes at some point in the past, populations remained relatively stable prior to the commencement of golden eagle predation in the mid-1990s and disease in 1999. Also, habitat alteration has not been a hindrance to the rapid recovery of the fox that has taken place.

Factor B: Overutilization for Commercial, Recreational, Scientific, or Educational Purposes

Island foxes were used in the past for pelts and ceremonial uses by Native Americans (Collins 1991b); however, this is no longer occurring. Therefore, island foxes are not currently exploited for commercial, recreational, scientific, or educational purposes. However, scientists are continually performing recovery efforts through FWS-issued 10(a)(1)(A) recovery permits. These research activities are not known to be a threat to island foxes.

Factor C: Disease or Predation

Disease

Island foxes are vulnerable to canine and other diseases. The catastrophic population decline of Santa Catalina Island foxes during 1999 to 2000 was caused by CDV; probably vectored to the island by a raccoon (*Procyon lotor*) (Timm et al. 2009). Analysis of CDV isolated from a Santa Catalina Island fox indicated it to be most closely related to the CDV strain found in mainland raccoons (Coonan et al. 2010; Timm et al. 2009) and a number of stowaway raccoons have been recently removed from Santa Catalina Island (Coonan 2013). Disease remains a concern for Santa Catalina Island foxes, since the island has high accessibility and a sizable human population.

All island fox populations have been surveyed for CDV, canine parvovirus, canine adenovirus, canine herpes virus, canine corona virus, leptospirosis, and toxoplasmosis (Garcelon et al. 1992; Coonan et al. 2000; Roemer 1999; Roemer et al. 2001b; Clifford et al. 2006). Antibodies against canine parvovirus and canine adenovirus are highly prevalent in most island fox populations, with the prevalence differing between islands and years (Garcelon et al. 1992; Coonan et

Recovery Plan for Four Subspecies of Island Fox

al. 2000; Roemer et al. 2001b; Clifford et al. 2006). Differences may be explained in part by differences in test sensitivities in the labs used for these surveys; the most recent survey used a lab with the most sensitive tests (Clifford et al. 2006). This recent survey indicated that Santa Catalina Island fox subspecies apparently has no protection against canine adenovirus (Clifford et al. 2006).

The recent finding of ear tumors in Santa Catalina Island foxes, confirmed to be a source of mortality in wild foxes, is of high enough frequency to be considered a concern (Coonan et al. 2010). The ear inflammation associated with cancer in Santa Catalina Island foxes may be due in part to *Otodectes* mite infections (Schwemm 2008). Treatment with acaricide may reduce the incidence of ear mite infection and thus inflammation (Coonan 2011).

The disease risk that domestic cats pose to island foxes is unclear. Pathogen sharing between island foxes and cats is minimal, but not absent (Clifford et al. 2006). A Santa Catalina Island fox mortality was known to be infected with *Toxoplasma* sp., an infectious agent that may have been acquired from cats (Timm et al. 2009). Infection with CDV in cats has been previously reported (Appel et al. 1974; Ikeda et al. 2001), and infected cats are capable of shedding CDV into the environment (Munson 2010). Two cats from Santa Catalina Island with CDV antibodies were also seropositive for feline immunodeficiency virus (FIV) and feline leukemia virus (FeLV), which make cats more vulnerable to other diseases and increases shedding of disease organisms, including toxoplasmosis (*Toxoplasma gondii*) (Hoover and Mullins 1991; Pedersen and Barlough 1991; Lin et al. 1992). Island foxes may be exposed to *T. gondii* oocysts shed in cat feces, in addition to the tissue cysts in prey items (Tenter et al. 2000). *Toxoplasma* has been documented to cause mortality in dogs (Dubey et al. 1989) and a Santa Catalina Island fox (Munson 2010). Concurrent distemper and *T. gondii* infection is associated with a high level of mortality in gray foxes (Davidson et al. 1992; Kelly and Sleeman 2003) and domestic dogs (Brito et al. 2002; Moretti et al. 2006).

Although caliciviruses have been shown to infect a variety of hosts and could possibly be passed between cats and foxes (Smith et al. 1998), calicivirus exposure is not correlated among foxes and cats, and presence of calicivirus antibodies in foxes on islands without cats suggests this interaction is not necessary for fox infection (Clifford et al. 2006). Although competition with cats is likely a more pressing threat to the island fox, the presence of cats on San Clemente and Santa Catalina Islands may initiate or help propagate an infectious disease epidemic, as pathogens such as distemper and rabies could circulate among these **sympatric** carnivores (Clifford et al. 2006).

Other mammals are a potential source of pathogens for island foxes, such as bats infected with rabies. Although rabies has never been found in island wildlife, if it were to occur, it would be difficult to effectively mitigate an outbreak. Hence the only defenses are a vaccination program and a subset of each fox subspecies are vaccinated against rabies.

Predation

On the northern Channel Islands, golden eagle predation was the primary threat to island foxes from the mid-1990s until 2007 (Coonan et al. 2005a; Coonan et al. 2010, Coonan 2013). Golden eagle predation was the cause of 13 of 15 mortalities of wild-born and released island foxes on San Miguel and Santa Rosa Islands from 2003 to 2005 (Coonan and Schwemm 2009). Golden eagle predation accounted for 69 of 92 Santa Cruz Island fox mortalities from December 2000 through June 2007 (Schmidt et al. 2007; R. Wolstenholme, The Nature Conservancy, pers. comm. 2007).

The onset of golden eagle predation resulted in the population decline of Santa Cruz Island foxes as demonstrated by the decrease in annual survivorship from 83 percent in 1994 to 39 percent in 1995 (Roemer et al. 2001b). San Miguel Island fox survivorship was 12 percent from 1998 to 1999, the tail end of the decline (Coonan et al. 2005c). As golden eagles were removed from the northern Channel Islands, annual Santa Cruz Island fox survivorship increased to 80 percent by 2003, a level that was previously estimated by demographic modeling to be the minimum necessary for recovery (Miller et al. 2001; Coonan et al. 2005a). Survival rates on Santa Cruz Island have remained above 90 percent since 2011 (C. Boser, The Nature Conservancy, pers. comm. 2014; Coonan 2014).

There are a number of resident domestic dogs in the interior of Santa Catalina Island and in the leeward coves and camps, many within active island fox territories as well as in the city of Avalon and the town of Two Harbors. In 2005, two deadly interactions occurred between Santa Catalina Island foxes and domestic dogs (IWS 2006; King, Santa Catalina Island Conservancy, pers. comm. 2014).

Factor D: Inadequacy of Existing Regulatory Mechanisms

As identified above, the primary causes of the island fox population declines were attributed to the unprecedented and unnatural levels of predation by golden eagles, the spread of canine distemper through the Santa Catalina Island fox subspecies, and the degradation of habitat by introduced herbivores. Federal, State, and local laws have not been sufficient to prevent island fox declines from these causes.

Recovery Plan for Four Subspecies of Island Fox

In 1971, the State of California listed the island fox as rare (a designation later changed to threatened), which means that either an incidental take permit is required under the California Endangered Species Act (Section 2081(b)) for otherwise lawful projects or a scientific collecting permit/research memorandum of understanding (Section 2081(a)) is required to take, collect, capture, mark, or salvage for scientific, educational, and non-commercial propagation purposes. State law does not require Federal agencies to avoid or compensate for impacts to the island fox and its habitat.

No regulatory mechanisms have been specifically designed for the protection of the four listed island fox subspecies, except for prohibitions against bringing pets ashore within Channel Islands National Park. Section 2.15 of the superintendent's compendium prohibits pets from all NPS islands, except for guide dogs for visually impaired persons. However, dogs have been used to eradicate pigs from Santa Rosa and Santa Cruz Islands, albeit with implementation of stringent quarantine procedures. Prohibitions against bringing dogs ashore are difficult to enforce (e.g., boaters have been observed bringing pets onshore to all three northern Channel Islands with island fox populations) (P. Schuyler, Independent Biologist, pers. comm. 2006). There is no prohibition of mainland animal visitation to Santa Catalina Island, and there is no requirement for health certification for dogs coming to the island, where the most tourist traffic and private residents occur, thus increasing the risk of exposing Santa Catalina Island foxes to disease. On Santa Rosa Island, the special use permit for the commercial hunting operation allowed for island-resident employees of the permittee to have "ranch dogs" on the island, though that practice ended in 2011 with the expiration of the permit and cessation of hunting activities (Coonan, pers. comm. 2014).

Several Federal laws apply to the management of NPS and Navy lands. These laws and guidelines include the National Environmental Policy Act (NEPA) and the Endangered Species Act. The NPS management is further dictated by Department of the Interior policies and NPS policies and guidelines, including NPS guidelines for natural resources management (NPS 1991), the Channel Islands National Park General Management Plan (NPS 1985), and the NPS Organic Act (16 U.S.C. 1, 2, 3, and 4). Both the NPS and the Navy have adequate authority to manage the land and activities under their administration for conservation of the island fox (e.g., feral animal removal). In addition to removing golden eagles, their prey base must be removed to prevent recolonization. Feral pigs have been removed from Santa Cruz Island, and all the elk and all but a few deer have been removed from Santa Rosa Island, with complete removal of the few remaining deer likely by early 2014.

San Miguel Island is owned by the Navy, but the NPS has responsibility for management of the natural, historic, and scientific resources of San Miguel Island

Recovery Plan for Four Subspecies of Island Fox

through a Memorandum of Agreement (MOA) originally signed in 1963, an amendment signed in 1976, and a supplemental Interagency Agreement (IA) signed in 1985. The MOA states that the “paramount use of the islands and their environs shall be for the purpose of a missile test range, and all activities conducted by, or in behalf of, the Department of the Interior on such islands, shall recognize the priority of such use” (Department of the Navy 1963). In addition to San Miguel Island, Santa Cruz Island and Santa Rosa Island lie wholly within the Navy’s Pacific Missile Test Center (PMTTC) Sea Test Range. The 1985 IA provides for PMTTC to have access to and use of portions of those islands, for expeditious processing of any necessary permits by the NPS, and for mitigation of damage to NPS resources from any such activity (Department of the Navy 1985). To date, conflicts concerning protection of sensitive resources on San Miguel Island have not occurred.

Federal protection of golden eagles by the Bald and Golden Eagle Protection Act of 1962, as amended, has increased the golden eagle population in mainland California (B. Walton, Santa Cruz Predatory Bird Research Group, pers. comm. 2004), and golden eagles have expanded their range. The protections extended to golden eagles limit management alternatives; removal of golden eagles requires a depredation permit from the FWS and lethal removal has not been authorized. Such a permit would allow golden eagles to be taken by firearms, traps, or other suitable means except by poison or from aircraft (50 CFR 22.23). A California State law (California Fish and Wildlife Code, section 3511) passed in 2003 allows the take of golden eagles and several other “fully protected” species, after a 30-day public notice period, for the purpose of recovering endangered species.

Regulatory mechanisms relevant to control of feral cats are discussed in the section on feral cats below.

Factor E: Other Natural or Manmade Factors Affecting its Continued Existence

Several other factors, including climate change, competition from introduced species, stochastic environmental factors and road mortalities may have negative effects on island foxes and their habitats.

Climate change

Climate change was not included as a threat in the listing rule for the island foxes (U.S. Fish and Wildlife Service 2004). Current climate change predictions for terrestrial areas in the Northern Hemisphere indicate warmer air temperatures, more intense precipitation events, and increased summer continental drying (Field et al. 1999, Cayan et al. 2005, IPCC 2007). It is unknown at this time if climate change in California will result in a warmer trend with localized drying, higher

precipitation events, or other effects, and predictions of climatic conditions for smaller sub-regions, let alone small offshore islands, remains uncertain.

Competition with feral cats

The CDFG, in recommending the retention of the threatened classification of the island fox under State law, cited the presence of competition with feral cats on Santa Catalina Island (CDFG 1987). Feral cats weigh on average twice as much as island foxes and may negatively affect foxes through direct aggression, predation on young, competition for food resources, and disease transmission (Laughrin 1978).

Direct aggression between foxes and cats has been documented in the wild, primarily near leased coves and campgrounds that provide food and shelter (D. Guttilla, Santa Catalina Island Conservancy, pers. comm. 2007). On Santa Catalina Island, frequent capture of cats in canyon bottoms and island foxes higher on slopes (Propst 1975) was attributed to competition and displacement of foxes by cats. On San Nicolas Island, where feral cat and island fox diets overlapped by 80 percent, foxes were absent from areas with cat densities exceeding 4 cats per km² (10 per mi²) (Kovach and Dow 1982; Phillips et al. 2007). After a large number of feral cats were removed from San Nicolas Island, foxes moved into areas previously occupied by cats (Laughrin 1978; Kovach and Dow 1982).

California State law (Food and Agricultural Code 31752.5) prohibits lethal control of feral cats unless cats are held for a minimum of 3 days. On Santa Catalina Island, this law could prevent the CIC from controlling or managing feral cats on the interior of the island, as it does not have adequate facilities to hold cats. The multiple ownership of the island further complicates the application of regulations and other strategies to address the resident feral cat population on the island. A Feral Animal Task Force convened by the City of Avalon, with representatives of the CIC and other island stakeholders, is working to address feral and free-ranging cats in the city and on the rest of the island. San Clemente Island, the other island with feral cats, is under federal jurisdiction, and thus is not bound by this State law. Feral cats were recently removed from San Nicolas Island (Hanson 2012).

Lack of genetic variation and stochastic environmental factors

As a population becomes genetically homogeneous, its susceptibility to disease, parasites, and extinction increases (O'Brien and Evermann 1988) as its ability to evolve and adapt to environmental change is diminished (Templeton 1994). The four listed island fox subspecies have all suffered large declines and are at risk of having reduced or low genetic diversity due to the population bottlenecks they have experienced (San Miguel Island fox: Gray 2002; Gray et al. 2001; San

Recovery Plan for Four Subspecies of Island Fox

Nicolas Island fox: Gilbert et al. 1990; Wayne et al. 1991a; Goldstein et al. 1999) (see Biological Information section for more complete discussion, p. 7). However, it should be noted that, although island foxes have little genetic variability and lost more during the recent decline, they are probably tolerant of low genetic variation, occasional bottlenecks, and higher inbreeding (Coonan et al. 2010).

The extremely small population sizes of the San Miguel Island fox and Santa Rosa Island fox made them vulnerable to extinction. Island endemics have a high extinction risk due to isolation and small population sizes (MacArthur and Wilson 1967), both of which make them more vulnerable to stochastic events such as drought or wildfires (Miller et al. 2001; Kohlman et al. 2005). In addition, lack of genetic variation may make a population less capable of overcoming stochastic events and the relationship between stochastic events and low genetic diversity can become **synergistic**. Therefore, the interrelationship between demographic risk (stochasticity) and genetic risk (low genetic diversity) can increase the risk of extinction.

Road mortalities

The lack of fear of human activities in wild island foxes coupled with relatively high levels of vehicle traffic on the southern Channel Islands result in a number of vehicle collisions each year. Death from vehicle collision on roads is the largest known source of mortality on San Nicolas and San Clemente Islands, accounting for approximately 17 island fox mortalities annually on San Nicolas Island (F. Ferrara, U.S. Navy. pers. comm. 2014) and a minimum of 26 foxes per year between the years 1991 and 1995 on San Clemente Island (Garcelon 1999). In 2013, 43 foxes died from vehicular trauma (Bridges et al. 2014). On Santa Catalina Island, annual averages of four foxes per year were killed by vehicles from 2003 to 2007 (Schmidt et al. 2004; Schmidt et al. 2005b; IWS 2006; IWS 2007; King and Duncan 2008), but the number of foxes killed has increased in the past several years as the fox numbers have increased: in 2011, 16 foxes were killed by vehicles, and in 2013, 12 foxes were killed by vehicles (King and Duncan 2013). Vehicle collisions on the northern Channel Islands are less common due to low traffic volume and rough dirt roads, which reduce vehicle speed.

Competition with deer and pigs for food items

Deer and elk consume fruits that are also preferred by island foxes. For example, mule deer and elk on Santa Rosa Island have been known to heavily browse the federally endangered Santa Rosa Island manzanita (*Arctostaphylos confertiflora*), the fruits of which have been found in island fox feces (Coonan, pers. comm. 2011). Similarly, pigs consume a variety of plant and animal items that are also

used by island foxes. Recent feral ungulate removal programs on both Santa Cruz and Santa Rosa Islands have removed this source of competition, with the exception of a few remaining deer on Santa Rosa Island.

1. Summary of Listing Factors A through E

Listing Factor A, the present or threatened destruction, modification, or curtailment of habitat or range, was not considered a substantial threat at listing. However, habitat modification, in the form of conversion of shrublands to alien annual grasslands by grazing, facilitated predation by golden eagles by decreasing cover available to island foxes. Additionally, habitat alteration has and continues to occur from vegetation type conversion, development, and/or fire. Thus, introducing measures and practices to maintain habitat integrity is recommended for attaining the long-term conservation of the island fox (see LONG-TERM CONSERVATION STRATEGY). Listing Factor B, overutilization for commercial, recreational, scientific, or educational purposes, was not considered a threat at the time of listing and is not considered a threat at this time. The primary threats to the island fox are encompassed within Listing Factor C, pertaining to disease or predation. Predation by golden eagles was one of the primary threats to the island fox at the time of listing and although still considered to be a threat to island fox populations on the northern Channel Islands, the degree of this threat has been decreased as a result of ongoing management practices. At the time of listing it was noted that a disease outbreak is believed to be the cause of the Santa Catalina Island fox population decline. Island fox populations will always be at risk of a disease outbreak, especially on Santa Catalina Island; however, the risk potential for disease outbreak can be and has been reduced. Listing Factor D, inadequacy of existing regulatory mechanisms, were not considered to be reasons for island fox decline at time of listing but were identified to have impeded or precluded the implementation of island fox recovery efforts, including The Bald and Golden Eagle Protection Act of 1962, as amended, California Fish and Game Code, section 3511, and California Food and Agricultural Code 31752.5. However, successful recovery strategies have been developed within the constraints of these regulatory controls. Listing Factor E, including climate change, competition from introduced species, stochastic environmental factors, and road mortality may have negative effects on island foxes and their habitats.

E. RECOVERY AND CONSERVATION EFFORTS

1. Recovery Actions for Listed Island Fox Subspecies

Recent island fox recovery efforts to date have included efforts to ameliorate the impact of golden eagle predation on the three island fox subspecies that occur on the northern Channel Islands and disease on Santa Catalina Island foxes. All of

these efforts have included captive breeding of island foxes to increase each of the four subspecies' populations to viable levels.

Northern Channel Islands

In April 1999, the Island Fox Working Group concluded that:

- Predation by golden eagles was the primary mortality factor acting on the island fox populations;
- Disease or parasites may have compounded the effects of predation; and
- The size of each of these three island fox populations was critically small and natural reproductive potential and recruitment were low.

At the time, the group agreed that establishing an island fox sanctuary and captive breeding program was necessary to safeguard individuals and to augment natural recruitment into the population.

The NPS began initiating emergency actions in 1999. The objectives were to remove the primary mortality factor affecting island foxes (golden eagle predation), and to recover island fox populations to viable levels via captive breeding. The NPS' island fox recovery strategy (Coonan 2003) utilized demographic modeling (Miller et al. 2001) to set the program size and determine the augmentation schedule for captive breeding. To achieve desired annual augmentation rates, the model estimated that an on-island captive population of 20 breeding pairs would be required.

Removal of Golden Eagles

Golden eagle translocation from the northern Channel Islands commenced in summer 1999. Golden eagles were trapped and subsequently released in northeastern California. Satellite telemetry affixed to the first 12 translocated golden eagles confirmed that none of the relocated eagles attempted to return to the islands for the 1.5 year life of the transmitter.

Between November 1999 and July 2006, 44 golden eagles, including 22 adults or near adults, were removed from Santa Cruz and Santa Rosa Islands (Latta et al. 2005; Coonan et al. 2010). Most adult and subadult eagles were trapped using a radio-controlled bow net set over dead or live bait (Jackman et al. 1994). Two helicopter net-gunning operations (O'Gara and Getz 1986) on Santa Cruz Island in June and October 2002 failed to capture any golden eagles, due to the difficulty in forcing eagles to ground in the rugged topography and dense vegetation. Ten nestlings were removed by hand from seven different nests (five from Santa Cruz Island and two from Santa Rosa Island) and fostered into mainland golden eagle nests or released via **hacking**. By mid-2005, seven golden eagles were estimated to remain on the northern Channel Islands, and the removal efforts were yielding

Recovery Plan for Four Subspecies of Island Fox

diminishing returns. In June 2006, a pair of nesting golden eagles was successfully captured via a net-gun from a helicopter using improved equipment and methods. This pair was removed from Santa Cruz Island and their single chick was removed from the nest by hand (Coonan et al. 2010). These represent the last eagles captured and removed from the islands.

Since 2006, two golden eagle predation events occurred where multiple individuals were preyed upon. In early 2007, 10 radio-collared foxes died from eagle predation on Santa Cruz Island, though helicopter and ground surveys failed to locate any golden eagles. In winter-spring 2010, 7 radio-collared foxes died from predation on Santa Rosa Island; two golden eagle sightings were recorded on Santa Rosa Island in that time period, and feathers collected at three fox mortality sites were identified as golden eagle feathers (Coonan and Guglielmino 2012). As on Santa Cruz, helicopter surveys failed to locate any golden eagles. Genetic analysis of the feathers confirmed their golden eagle origin, and further identified them as coming from three highly-related individuals, likely juvenile sibling eagles (Talbot et al., in prep.).

With the exception of the two scenarios described above, eagle predation on northern Channel Island foxes has been almost negligible since 2006. Island fox annual survival has remained above 90 percent for most of that period, and populations have steadily increased. Any predation has been short-term in nature and has likely been the result of dispersing, juvenile or sub-adult eagles; no adult golden eagles have attempted nesting on the northern Channel Islands (Coonan et al. 2010). Nesting adult golden eagles present a higher threat to foxes, because eagle energetic requirements during breeding are high and are satisfied by the delivery of small prey items to the nest (as opposed to carrion) (Coonan et al. 2010).

Other golden eagle genetic work supports the long-term success of eagle translocation efforts. Sonsthagen et al. (2012) investigated the genetics of mainland golden eagles and those translocated from the islands, finding that the island population was likely the result of one colonization event. The likelihood of another successful golden eagle colonization is low, given changes in non-native prey availability (see below) and monitoring/mitigation by land management agencies.

Island fox recovery may ultimately depend on promoting ecological conditions that dissuade golden eagle use of the Channel Islands, including maintaining the islands free of non-native herbivores and restoring bald eagles to the northern Channel Islands.

In 2005-2006, 5,036 feral pigs were removed from Santa Cruz Island, with no individuals known to be remaining on the island (Macdonald and Walker 2007;

Morrison et al. 2007). In 2011, removal of the non-native mule deer and elk on Santa Rosa Island was implemented as part of a court settlement (NPS 1998). Elk were quickly removed and as of 2014, all but a few mule deer had been removed from Santa Rosa Island. Thus, in 2014, the northern islands were effectively ungulate-free for the first time since the mid-19th Century.

The recent, successful restoration of bald eagles to the Channel Islands may also provide a deterrent to future golden eagle colonization of the islands. Sixty-one bald eagles were released on Santa Cruz Island as the result of annual experimental reintroductions of juvenile bald eagles from 2002 to 2006 (Coonan et al. 2010). By 2013, there were estimated to be at least 40 bald eagles occupying Santa Cruz, Santa Rosa, and Anacapa Islands. In spring 2006, two bald eagle pairs established nests on Santa Cruz Island, and each successfully fledged a single chick (Coonan et al. 2010). These breeding pairs represented the first active and successful bald eagle nests on the northern Channel Islands since the late 1950s (Kiff 1980). In 2013 there were 8 breeding pairs of bald eagles on the northern Channel Islands, and 6 young were fledged from their nests (Sharpe 2014).

Captive Breeding

The critically low island fox populations on the northern Channel Islands and on Santa Catalina Island in 1999/2000, prompted management agencies to begin captive breeding on each of the islands where federally listed subspecies occurred. Captive breeding was conducted on each island, rather than on the mainland, due to the potential for vectoring pathogens to the islands by animals raised in mainland facilities. Island foxes had never been bred in captivity before, and so island managers and others worked with the zoo community and drew from other captive canid programs to develop all aspects of island fox husbandry: pen size and shape, diet, breeding strategies, veterinary care, and release methods (Coonan et al. 2010). Island foxes bred successfully in captivity on all islands (see below), though the programs encountered obstacles such as stress, male aggression, female abandonment and an outbreak of mastitis, all of which reduced reproductive success and prompted research and adjustment in captive breeding methods. Cessation of captive breeding and reintroduction 10 years after the programs began speaks to the program's success; by the program's end, reproduction in the reintroduced wild populations was outpacing that in captivity.

Upon receiving recommendations from a panel of experts in 1999, the NPS began captive breeding on San Miguel Island in the summer of 1999, with construction of pens and capture of wild island foxes for captive propagation. By January 2000, 14 island foxes had been captured and placed in the pens; the remaining wild fox was captured in 2001. Four of the captured foxes were males and were paired with four females for breeding, and 8 of the 15 potential founders

Recovery Plan for Four Subspecies of Island Fox

eventually bred in captivity. In 2004, after five years of breeding, the San Miguel Island fox captive population had increased to 50 animals, exceeding the target captive population size of 40 animals and allowing initial releases back to the wild in fall 2004. The San Miguel Island fox captive breeding and reintroduction program ended in 2007, due to high reproductive success and survival in the wild. During 9 years of captive breeding, 53 pups were born in captivity, and 62 foxes released to the wild. The recovering wild population steadily increased since releases began in 2004 (Coonan and Schwemm 2009; Coonan et al. 2010).

A captive breeding program was initiated for Santa Rosa Island in 2000. The initial captive population on Santa Rosa Island was 15 animals, which proved to be the island's remaining fox population. Some females were pregnant when captured, and three litters were born in captivity in 2000. With an increase to 56 foxes in 2003, the captive population on Santa Rosa Island exceeded the target captive population size of 40 foxes, and initial releases began in winter 2003/2004. Annual releases continued through 2008, after which captive breeding was ceased on Santa Rosa Island. In 9 years of captive breeding, 87 pups were born in captivity, and 93 foxes (including some of the foxes originally brought into captivity) were released to the wild (Coonan et al. 2010).

Captive breeding was also conducted on Santa Cruz Island as a joint venture by the NPS and TNC. The status of eagles and foxes on Santa Cruz Island was assessed at the 2001 meeting of the Island Fox Conservation Working Group, and consensus was that captive breeding was warranted. In February 2002, a 10-pen captive breeding facility was built on Santa Cruz Island by the NPS and TNC. Between 2001 and 2003 this facility was stocked with 18 adult island foxes caught as known pairs or individuals from separate areas of the island. Captive-born pups of breeding pairs were released in 2002 and 2003, but experienced high mortality and were pulled back into captivity in 2004. A second facility with 27 pens was added in 2004 to accommodate captive born individuals (Schmidt et al. 2007). No releases occurred in 2004 or 2005, and the captive population grew to 62 animals in 2005. After capturing the last breeding golden eagle pair in June 2006, releases of captive individuals began in July 2006 and concluded in July 2008 (Hudgens and Sanchez 2009).

On Santa Catalina Island, captive breeding was developed by IWS, funded by the CIC (see below).

Parasites

Parasites have not specifically been considered a disease threat to wild island foxes; however, parasite burdens in captive individuals were a cause of concern. For this reason, fecal parasite surveys were conducted for captive foxes on San Miguel, Santa Rosa, and Santa Cruz Islands as part of a risk assessment for treating **endoparasites** in captive island foxes (Coonan, pers. comm. 2011). A

Recovery Plan for Four Subspecies of Island Fox

panel convened by USGS-BRD for a risk assessment (Sohn and Thomas 2005) determined that there was little clinical justification for the widespread use of **anthelmintics** in island foxes, given that non-target parasites might be killed by these drugs, with dire consequences for treated foxes. The panel compiled a list of preferred anthelmintics, recommended dosages, and **contraindications**, should treatment for internal parasites be required.

Canine heartworm (*Dirofilaria immitis*) was suspected to be a threat to island foxes because positive *Dirofilaria* antigen tests were documented in four of the six island fox subspecies (San Miguel Island Fox, Santa Cruz Island fox, Santa Rosa Island fox, and San Nicolas Island fox) (Roemer et al. 2000). However, necropsies of over 400 island foxes from all islands have found no evidence of heartworm or heartworm disease (Munson 2010; Coonan et al. 2010). In fact, these results suggest tests for *Dirofilaria* antigen are not specific and possibly cross-react with another parasite antigen, and given the geographic distribution of *Dirofilaria* antigen-positive foxes with *Spirocerca* infections, cross-reaction with *Spirocerca* antigens by these tests is very likely (Coonan et al. 2010).

In 2013, necropsies of five radio-collared San Miguel Island foxes revealed substantial, and in several cases massive, parasitism by an unidentified Acanthocephalan (spiny-headed) parasite in the intestines (Coonan 2014). The parasite burdens were associated with colitis, enteritis and emaciation, and likely contributed to mortality of the individuals. Acanthocephalans, which occasionally cause mortality in sea otters (*Enhydra lutris*), have not been previously recorded in island foxes.

Disease

Island foxes are thought to be vulnerable to canine and other diseases. The catastrophic decline on Santa Catalina was caused by CDV (Timm et al. 2009). Therefore, proactive and ongoing mitigation for disease is being implemented for island foxes. Island foxes on all islands are vaccinated against CDV and rabies, the two diseases for which active mitigation measures could not be implemented in a timely manner once an outbreak was detected. The number of foxes vaccinated on each island is generally the number required to start a captive breeding program (75-100), were the population to be affected by an epidemic (see Appendix A in Coonan 2011, and Appendix 3 and 4 of this plan).

Epidemic response plans, which are required for delisting (see Recovery Criteria) are being developed for each island fox subspecies and represent a critical tool in disease mitigation. The plans provide guidelines to ensure diseases, even novel ones, are appropriately detected and mitigated. Detection is primarily by maintaining a sample of radio-collared, unvaccinated sentinel animals and monitoring them for mortality signals at least once a week. When a mortality

Recovery Plan for Four Subspecies of Island Fox

signal is detected, carcasses are retrieved and sent for necropsy to determine mortality cause and possible presence of disease. If a disease is detected, decision trees and the use of the incident command system guide selection and implementation of appropriate responses. An epidemic response plan has been developed for San Clemente Island (Hudgens et al. 2011), and plans are currently being developed for the northern Channel Islands and Santa Catalina Island (Hudgens et al. 2013; Hudgens et al., in prep.). Monitoring of radio-collared foxes for survival rate and mortality cause is currently being conducted for five of the six island fox subspecies (Coonan 2013). Serology is conducted at least once every 5 years on each fox subspecies to detect presence of antibodies to various diseases. The results indicate wide variation among the six subspecies in seroprevalence to CDV, canine adenovirus, canine parvovirus, canine herpes virus and canine coronavirus (Coonan et al. 2010). None of the seroprevalence values were associated with population decline. The relatively high seroprevalence of canine parvovirus and canine adenovirus in some island fox populations without accompanying mortality suggests that these viruses are endemic and of low pathogenicity in island foxes (Munson 2010).

Radio-telemetry monitoring as well as information from annual trapping efforts have revealed the occurrence of other diseases in island foxes. These include toxoplasma and ceruminous gland tumors (in Santa Catalina Island foxes), leptospirosis (in Santa Rosa Island foxes), amyloidosis, systemic mineralization, and thyroid disease (Munson 2010; Coonan 2013; Guglielmino 2012). Ceruminous gland tumors occur in the ear canals of one third of adult foxes on Santa Catalina Island, and are associated with severe inflammation and the presence of ear mites. Experimental treatment of mites with ivermectin has reduced inflammation in fox ears on the island (Coonan 2011).

Guidelines for vaccination (Appendix 4), preparation of epidemic response plans, regular serosurveys and collection of other biomaterials, and quarantine protocols for dogs or foxes moved between Santa Cruz, Santa Rosa, San Miguel, San Clemente, and San Nicolas Islands (Appendix 6) have all been developed by the Fox Health Group of the Island Fox Conservation Working Group (Munson 2010; Coonan 2010, 2011, 2012).

Santa Catalina Island

In response to the catastrophic Santa Catalina Island fox decline that was due to CDV, the CIC, which owns and manages 88 percent of the island, contracted with the IWS to develop and implement island fox recovery actions between 1999 and 2005. Beginning in 2006, CIC assumed full responsibility for Santa Catalina Island fox conservation efforts.

CIC and IWS implemented the following four recovery actions (Kohlmann et al. 2005):

Recovery Plan for Four Subspecies of Island Fox

1. Intensive mark-recapture sampling to estimate Santa Catalina Island fox population size after the decline was detected;
2. Translocation of juvenile Santa Catalina Island foxes from the dense population on the western end of the island to the eastern end, where foxes had been essentially extirpated;
3. Vaccination of nearly the entire Santa Catalina Island fox population against CDV following trials of vaccine safety and efficacy using captive individuals; and
4. A captive breeding program to augment the Santa Catalina Island fox population.

In 2001 and 2002, 22 juvenile Santa Catalina Island foxes were translocated from the west end of the island to the east end. Survival of these individuals was very high; in 2004, at least 77 percent (n=17) of the translocated foxes were known to be alive, with at least 6 individuals reproducing in their new locations (Coonan et al. 2010). Nearly 50 percent of the translocated foxes started reproducing in their new locations within a year of being moved.

The gray fox, a close relative of the island fox is known to be highly susceptible to CDV and modified live CDV vaccines (Hallbrooks et al. 1981). As a result of this susceptibility, trials were conducted on captive Santa Catalina Island foxes and demonstrated that a new recombinant vaccine (Merial Purevax Ferret®, Merial, Inc., Athens, GA) was both safe and induced antibody production. Following these trials, vaccination of wild Santa Catalina Island foxes began in 2000 (Timm et al. 2000), and currently all six island fox subspecies (including two non-federally listed subspecies) are vaccinated against CDV (Coonan et al. 2010).

In 2000, the CIC in conjunction with IWS established a captive breeding program for Santa Catalina Island foxes. Between 2000 and 2002, 27 Santa Catalina Island foxes were brought into captivity. From 2001 to 2004, 79 individuals were released from captivity, including 37 captive-born pups and 20 of the original wild-captured adults. Survival of captive-born pups was very high (Schmidt et al. 2005b). In 2003, the first wild pup was born to a released captive-born individual (Clifford 2006). Reproduction by released individuals has continued and both translocated and captive-bred foxes have formed pairs with each other and with resident wild foxes. Based on the high survival (75 percent) of foxes released from 2001 to 2003, and the natural productivity of foxes in the wild, the captive breeding effort on Santa Catalina Island was terminated after the 2004 breeding season.

Recovery Plan for Four Subspecies of Island Fox

Although wildlife biologists and conservationists have recommended removal of feral cats from Santa Catalina Island for decades (Anon 1931; Propst 1975; Collins and Martin 1985; Menke and Miller 1985; S. Sillett, Smithsonian Migratory Bird Center, Washington, D.C., pers. comm. 2004; Backlin et al. 2005; Clifford et al. 2006), there is still no long-term, island-wide feral cat management program on Santa Catalina Island. For the last 20 years, the local humane society has practiced trap-neuter-release in Avalon and Two Harbors, where cats are maintained in unconfined feeding colonies ranging from 5 to 75 cats each. These colonies attract reproductively intact cats from surrounding wildland areas and serve as disposal sites for unwanted pets (Guttilla 2007).

During the annual Santa Catalina Island fox trapping efforts from 2004 to 2007, feral cats that were captured incidentally were tested for feline leukemia virus (FeLV) and feline immunodeficiency virus (FIV). Diseased cats were euthanized and healthy cats were sterilized, pit-tagged, and vaccinated for rabies (Guttilla and Stapp 2010). The CIC has continued to collect data on disease prevalence, diet, and feral cat distribution across the island; however, the low trapping-success-rate and difficulty in detecting feral cats has precluded the ability to accurately calculate feral cat population estimates (Guttilla 2007). Additionally, the introduction of animals, domestic or exotic, to the island has not been regulated and municipal and county regulations are outdated and not enforced. Furthermore, public opposition to lethal control hinders efforts to fundraise for the development and maintenance of a feral cat control program.

Staff from CIC now pursue and remove stowaway raccoons and opossums (*Didelphis virginiana*) reported from the island (King and Duncan 2014). Raccoons are the likely source of CDV that decimated the Santa Catalina Island fox population in 1999-2000 (Munson 2010).

2. Conservation Efforts for non-Listed Island Fox Subspecies

Conservation efforts are currently implemented by the Navy for San Nicolas and San Clemente Island foxes, the two island fox subspecies that are not federally listed. In 2003, the FWS and the Navy signed a Conservation Agreement to facilitate implementation of conservation measures for the island fox on San Clemente Island. Implementation of the Conservation Agreement has supported ongoing research, improved population monitoring, and impact minimization measures to reduce the potential of threats to the island fox on San Clemente Island.

The FWS currently coordinates with the Navy regarding conservation measures to benefit island foxes on San Nicolas and San Clemente Islands. The effort includes an evaluation of population status, identification of directions for future

Recovery Plan for Four Subspecies of Island Fox

research, and recommendations on continuing the following conservation measures:

- Including effects on island foxes in all NEPA documents and mechanisms to minimize effects to island foxes;
- Continuing measures to minimize mortality from vehicle strikes;
- Continuing public awareness campaigns concerning island fox biology and status;
- Reducing potential adverse effects from pest management on the island fox;
- Prohibiting dogs on San Clemente Island or San Nicolas Island;
- Continuing feral cat control on San Clemente Island and;
- Maintaining refuse bin modifications on San Nicolas Island and implementing bin modifications on San Clemente Island.

To reduce the impact of vehicles as a mortality source, speed limits have been established and education programs have been developed targeting island personnel. The Navy has modified refuse bins, and discourages hand-feeding of island foxes.

Feral cats were recently removed from San Nicolas Island in an effort funded by the Montrose Settlements Restoration Program and implemented by Island Conservation and IWS. A total of 59 feral cats were removed by live trapping and hunting in 2009-2010 (Hanson 2012).

Efforts to control feral cats on San Clemente Island began in 1986 (Phillips and Schmidt 1997; U.S. Fish and Wildlife Service 2004). When cat eradication efforts were interrupted for a 6-month interval, cat populations rebounded to pre-control levels and, in some instances, doubled in size (Phillips and Schmidt 1997). However, in areas where control was maintained for three consecutive seasons, cat numbers were reduced by 20 to 50 percent (Phillips and Schmidt 1997). Feral cat removal continues on San Clemente Island (D. Garcelon, Institute for Wildlife Studies, pers. comm. 2014). The Institute for Wildlife Studies removed 328 cats from San Clemente Island in 2009, 276 cats in 2011, and 166 cats in 2012 (Biteman et al 2010; 2012; 2013).

3. Monitoring Efforts

Monitoring island fox populations has been, and will continue to be, a necessary activity. Given the success of recent recovery efforts, risk to all subspecies has decreased substantially. However, some risk will remain even after recovery and de-listing because of the inherently small subspecies population sizes, lack of genetic diversity as a result of bottlenecks, and isolation from other potential population sources. Island fox monitoring has been conducted for a number of years on each of the Channel Islands. Although methods have varied among islands and through the years, island fox monitoring is practically standardized across the islands, thanks to work by the IRT and the Island Fox Working Group. Island fox population monitoring became standardized in the 1990s, when density estimation from grid trapping was used on four of the six islands (Roemer et al. 1994). Recent efforts have used demographic modeling with >25 years of fox monitoring data to refine monitoring so that it is threat-based and able to gauge recovery (Rubin et al. 2007, Bakker and Doak 2009, Bakker et al. 2009, Coonan et al. 2010).

Monitoring Plans

A multi-year and highly collaborative monitoring planning process has been completed by the IRT and the RCG. The process included:

- Issuance of the Technical Analysis Request (TAR) 2.1 “Development of Population Monitoring Plans for Free-Ranging Island Foxes” to identify island fox monitoring needs (see Appendix 5).
- Development of estimates of demographic parameters by V. Bakker and colleagues through the compilation and robust analysis of island fox population data (Bakker et al. 2009).
- Development of a population viability analysis (PVA) by D. Doak and V. Bakker (see Appendix 2) to provide the conceptual framework for understanding island fox demographics and threats to island fox viability (Bakker and Doak 2009).
- Development of a set of guidelines by the Island Fox Health Technical Expertise Group (TEG) (see Appendix 3), which outlines recommendations for monitoring the health of wild island foxes.
- Development of a monitoring plan for San Clemente Island foxes (Spencer et al. 2006) to serve as a framework for TAR 2.1.

The Conservation Biology Institute (CBI) coordinated the development of island-specific monitoring protocols with land managers, TEGs, and statisticians to

identify monitoring needs and to develop the most robust and efficient monitoring protocols for each island. Rubin et al. (2007) developed specific monitoring recommendations for each of the four listed subspecies, as well as for the San Nicolas Island fox. Recommendations considered managers' goals, ecological and physical characteristics of the islands as they relate to monitoring needs and constraints, population modeling, evaluation of statistical robustness, and assessment of island representation (see Table 1 in Appendix 5). The monitoring plan for each island includes a scenario for monitoring survival and cause-specific mortality rates and two alternative scenarios for trapping to collect demographic data, such as population size and density (see Appendix 5).

F. CURRENT STATUS AND TREND

1. San Miguel Island fox

On San Miguel Island, no foxes existed in the wild during the period of captive breeding. Reintroduction to the wild began in 2004 and ended in 2007, by which time 62 foxes had been released. Survival and reproduction was high for released foxes and the population grew rapidly; the annual rate of increase averaged 76 percent from 2004-2013. The estimated adult population increased from less than 100 in 2006 to approximately 550 in 2013 (Table 1; Coonan et al. 2010, Coonan 2014).

Relatively few San Miguel Island foxes have died from predation since re-introduction. Although occasionally eagles have taken foxes, it has not affected fox annual survival, which has stayed above 90 percent for most of period from 2004-2013. In 2013, however, island fox survival declined to about 80 percent, and five of the 11 mortalities that occurred in radio-collared foxes had evidence of a parasite never before recorded in island foxes (Coonan 2014). Necropsy of those foxes revealed acanthocephalans (spiny-headed worms) in massive amounts in the lower intestine, associated with enteritis (inflammation of the lower intestine), colitis, and emaciation. Acanthocephalans likely contributed to the death of four of the foxes. Specimens of the parasite are currently being studied to identify it to species, which will suggest likely intermediate hosts (sand crabs or lizards, depending on whether the parasite is *Prolificollis* or *Oncicola*). Continued monitoring of mortality causes will determine whether the parasite is a significant mortality source for San Miguel foxes, and requires management.

Tracking population estimates for the total population (both adults and juveniles) reveals that it has hovered around 550 foxes since 2010, and this may very well represent carrying capacity for the island (Coonan 2014). This is supported by the general decline in reproductive effort as the population has increased. In 2013, only three pups were caught on the San Miguel monitoring grids, compared to 32 the previous year. The low reproductive output is likely due both to high fox

density and extended drought. The combination of low mortality and robust population growth puts the San Miguel Island fox subspecies at acceptably low risk of extinction, according to the recovery planning tool (see Recovery Criteria).

2. Santa Rosa Island fox

Santa Rosa Island foxes were brought into captivity in 2000, a year after San Miguel Island foxes. As on San Miguel Island, there were no foxes in the wild during the period of captivity. Releases to the wild occurred in 2003-2008, during which time 93 foxes were released. As on San Miguel Island, there was considerable reproductive success in the wild, along with variable mortality (see below), and the recovering population grew rapidly. The estimated adult population grew from about 40 animals in 2005 to 732 in 2013 (Coonan et al. 2010, Coonan 2014), with an average annual rate of increase (λ , lambda) of 60 percent. The high reproductive success and survival in the wild allowed the Santa Rosa Island captive breeding and reintroduction program to cease operations in 2008. When juveniles were included, the island-wide population estimate was 894 in 2013 (Table 1). This is likely not close to carrying capacity, since the rate of increase for both the adult and total population continues to be high (Coonan 2014). The Santa Rosa Island population may ultimately approach 1,200 - 1,500 adults, similar to neighboring Santa Cruz Island.

The released and recovering fox population on Santa Rosa Island incurred greater mortality than did the neighboring populations on San Miguel and Santa Cruz Islands (Coonan et al. 2010; Coonan and Guglielmino 2012; Coonan 2014). Predation by golden eagles was significant in 2004-2005, and again in 2010. Golden eagles bred on the island, at two territories, through 2005.

Morphological and molecular analysis of feathers from fox kill sites in 2010 suggested that predation was the work of three closely related, juvenile golden eagles which likely had dispersed to the island from the mainland (P. Trail, U. S. Fish and Wildlife Service – National Fish and Wildlife Forensics Laboratory, Ashland, Oregon, pers. comm. 2014; S. Talbot, U.S. Geological Survey, pers. comm. 2014). Although occasional eagle predation may occur on Santa Rosa Island and the other northern islands, it has not contributed significantly to annual mortality since 2010. Also, golden eagles are unlikely to colonize the islands again, because the alien ungulate prey base has been removed, and predation mortality – and eagle presence – is tracked via monitoring of radio-collared foxes.

Overall predation accounted for 33 of 73 island fox mortalities from 2003-2013 on Santa Rosa Island (Coonan, pers. comm. 2014). Mortality cause was not determined in 29 cases, due to advanced decay of the carcass. Other mortality causes included entrapment in drain pipes (3 cases); intestinal intussusception (collapse of one section of the intestine into another, 1 case); cholecystitis

Recovery Plan for Four Subspecies of Island Fox

(inflammation of the gall bladder caused by gallstones, 2 cases); leptospirosis (2 cases); and emaciation (3 cases, following wounds). Leptospirosis on Santa Rosa Island, as well as acanthocephalans on San Miguel Island, both illustrate the dynamic nature of pathogens in island foxes. *Leptospira* had been observed rarely in island foxes until 2010, when 19 of 31 Santa Rosa Island fox blood samples had antibodies for *Leptospira*, many at very high titers (Coonan and Guglielmino 2012). This apparent outbreak was confirmed by culturing urine samples in early 2011, which showed that both island foxes and island spotted skunks were shedding the pathogen. Lower titers and seroprevalence in late 2010 suggested the outbreak had run its course. These two cases, discovered by mortality monitoring of radio-collared foxes, underscore the need for disease monitoring and mitigation plans (e.g., epidemic response plans).

Significant mortality during the early phase of reintroduction and again in 2010 prevented the Santa Rosa subspecies from attaining the level of biological recovery that the San Miguel and Santa Cruz Islands subspecies had attained by 2013. By that year, the Santa Rosa Island fox subspecies had one three-year average of population and mortality estimates, below the 5 percent extinction isocline (see Appendix 2). Five such three-year estimates are required to meet Recovery Criterion 1, and so the Santa Rosa subspecies is likely to meet this recovery criterion by 2017.

3. Santa Cruz Island fox

Between 2001 and 2003, the NPS and TNC brought 18 Santa Cruz Island foxes into captivity, leaving no more than 100 individuals in the wild (Coonan and Rutz 2002; Coonan et al. 2004). Sixteen of the 18 founders, and a number of captiveborn foxes bred in captivity produced a total of 85 pups over 6 breeding seasons. By summer 2006, the captive population had increased to 81 individuals.

Due to the impact of golden eagle predation on released Santa Cruz Island foxes, few were released from captivity prior to 2006. Seven of 12 captive-born Santa Cruz Island foxes released to the wild in 2002 to 2003 died from golden eagle predation within 5 weeks of release (Coonan et al. 2005a). The remaining captive-born animals were brought back into captivity by January 2004, and no Santa Cruz Island foxes were released in 2004 or 2005.

A high survival rate (approximately 80-percent survival in 2003) coupled with excellent reproduction in the wild, increased the wild Santa Cruz Island fox population to at least 156 known individuals by early 2006, with an island-wide estimate of 207.

With the removal of the last known nesting golden eagle pair from Santa Cruz Island in June 2006, releases of captive-born pups resumed in July 2006. Survival

rates of captive foxes were much higher than in previous release attempts (survival of captive born foxes released in 2006 was 67 percent in 2007). Therefore, the remaining foxes were released in 2007 and 2008 and the captive breeding facility was closed. Since that time golden eagle predation on foxes has decreased. Only 10 of 37 recorded mortalities in 2008-2013 were ascribed to golden eagle predation. As of 2013 the total estimated population of Santa Cruz Island foxes was 1,354 and the survival rate has been greater than 90 percent since 2011 (Table 1; C. Boser, The Nature Conservancy, pers. comm. 2014). As on San Miguel and Santa Rosa Islands, there have been occasional golden eagle sightings on Santa Cruz Island, and perhaps one predation mortality per year, but it has had a negligible effect on fox survival and population growth.

Pathogens such as acanthocephalans or *Leptospira* have not been found on Santa Cruz Island. Recent serological testing showed no exposure of Santa Cruz Island foxes to any of five pathogens (canine adenovirus, CDV, canine parvovirus, canine coronavirus and canine herpes virus) (Coonan et al. 2010).

As on San Miguel and Santa Catalina Islands, Santa Cruz Island foxes have reached the threshold of density and mortality combinations that denote biological recovery. Estimates for 2013 demonstrate five three-year averages of mortality and population safely below the 5 percent extinction isocline for the subspecies (see Appendix 2). The population may have reached carrying capacity, as on San Miguel Island. Adult population estimates for Santa Cruz Island in 2010-2012 were 907, 827, and 1,073 foxes, respectively; though there was no statistical difference among those estimates (Table 1; Coonan et al. in press; C. Boser, The Nature Conservancy, pers. comm. 2014).

4. Santa Catalina Island fox

Like the Santa Cruz Island fox subspecies, Santa Catalina Island foxes recovered fairly quickly from the population lows in 1999-2000, with a total population estimate (including pups) in 2013 of 1,852 individuals, of which 1,594 were adults (Table 1; King, Santa Catalina Island Conservancy, pers. comm. 2014). The population has steadily increased since 2000, and may be approaching carrying capacity, with a total population estimate (including pups) of around 1,500 foxes in 2011 and 2012. As of 2014, the CIC annually monitored fox population size by transect trapping in the fall, and maintained a large sample of radio-collared foxes, including unvaccinated disease sentinels, for mortality monitoring.

Disease remains a concern for Santa Catalina Island foxes, since the island has high accessibility and a sizable human population with domestic dogs and cats, many likely unvaccinated. Furthermore, seven raccoons and two opossums were removed from the island between 2007 and 2013, typically arriving as stowaways

on boats (King and Duncan 2014). The strain of CDV that caused the catastrophic decline of Santa Catalina Island foxes in 1999-2000 was most closely related to that observed in raccoons (Timm et al. 2009).

The CIC vaccinates over 300 foxes annually (about 80 percent of all foxes captured) against CDV and rabies. Recent serology indicated low, but nonetheless present, seroprevalence to CDV in Santa Catalina Island foxes, and some seroprevalence to canine adenovirus, canine parvovirus and canine corona virus (King and Duncan 2014).

Santa Catalina Island foxes continue to have an unusually high incidence of ceruminous gland tumors, cancers occurring in the ear canal and associated with inflammation from ear mites. Treatment with ivermectin has been found to reduce inflammation and incidence of cancer (Coonan 2011).

Other sources of mortality include, but are not limited to, effects from competition with feral cats and mortality from vehicle strikes.

5. San Clemente Island fox

Recent trends in annual population estimates indicate that the San Clemente Island fox is increasing (λ is equal to 1.15), and the total population size in 2013 was estimated to be 1,000 foxes (M. Booker, U.S. Navy, pers. comm. 2014). Previously, Coonan (2003) had reported an estimated population size of 680 adult San Clemente Island foxes. Schmidt et al. (2005a) reported an estimated population size of 396 individuals in 2004, after applying density corrections. San Clemente Island fox survey, monitoring, and population estimate methods changed in 2007. The 2007 San Clemente Island fox population estimate, based upon the new methodologies, ranges from 302 individuals to 727 individuals (Garcia and Associates 2008). The 2009/2010 population was estimated to be 714 individuals (Coonan 2011).

6. San Nicolas Island fox

Island foxes have been monitored annually on three permanent grids on San Nicolas Island since 2000. Over that time, the population has mainly been stable, with some of the highest island fox densities ever recorded (up to 15 foxes per square kilometer). However, while individual foxes appear to be in good health, the population currently seems to be in the midst of a sustained decline which began in 2010. The island-wide population estimate in 2009 was 619 individuals; in 2012 it declined to 460 individuals; and in 2013, declined again to 341 individuals (Ferrara, pers. comm. 2014; Hudgens and Garcelon 2014). Most likely, the decline may be due to negative density-dependence causing the population to correct after exceeding carrying capacity or it is a result of extended drought conditions limiting resources. Feral cats were removed from the island in

Recovery Plan for Four Subspecies of Island Fox

2009-2010, but cat trapping activities had no effect on island fox survival. Additionally, a disease outbreak has not been ruled out. Serology results from 2013 indicated that 62 percent of tested foxes had measurable titers for CAV; 30 percent were high titers indicating recent exposure. None of the 27 previously unvaccinated foxes tested positive for CDV, but 7 had titers of 1:8 but less than 1:16 which were considered suspect (Hudgens and Garcelon 2014). Canine parvovirus was not detected in 2013, though it was detected in 16 percent of 2009 samples and in nearly 99 percent of samples from 2001-2003.

7. Summary

On the northern Channel Islands, golden eagle removal and captive breeding programs with reintroductions have reduced the risk of extinction for the San Miguel Island fox, Santa Rosa Island fox, and Santa Cruz Island fox and have allowed the re-establishment of wild populations on Santa Rosa and San Miguel Islands. Although there is still some predation by golden eagles, there have been no golden eagle nests on the northern Channel Islands since 2006, which has likely been attributable to the efforts associated with golden eagle capture and translocation, feral pig and ungulate eradication, and reintroduction of bald eagles. Although mainland animals are not permitted to be transferred to the northern islands, the possibility of unintentional or illegal introduction exists. Thus the introduction of infectious disease remains a potential threat to Santa Cruz, Santa Rosa, and San Miguel Island foxes.

On the southern Channel Islands, following captive breeding and disease mitigation efforts, the Santa Catalina Island fox population is increasing. Disease remains a concern for Santa Catalina Island foxes, since the island has high accessibility and a sizeable human population. Of the two non-listed subspecies (San Clemente Island fox and San Nicolas Island fox), San Clemente Island foxes appear to be stable, while the San Nicolas Island foxes currently appear to be in the midst of a sustained decline which began in 2010. Potential threats to Santa Catalina, San Nicolas and San Clemente Island foxes include competition with feral cats, vehicle strikes, and the introduction of infectious disease.

Recovery Plan for Four Subspecies of Island Fox

II. Recovery Strategy

The two primary known threats that resulted in the listing of the four subspecies of island fox (San Miguel Island fox, Santa Rosa Island fox, Santa Cruz Island fox, and Santa Catalina Island fox) as endangered were predation by golden eagles and the transmission of disease. Additionally, because the population size of each island fox subspecies is small, they are threatened by stochastic events and the effects of low genetic diversity. Recovery of each subspecies will be achieved by removing, or substantially reducing, known threats, such as predation by golden eagles and disease-related mortality, and increasing populations to viable levels for long-term survival of each subspecies. The strategy of this recovery plan is to continue the current recovery efforts and to improve and expand recovery actions as necessary. Recent and ongoing island fox recovery efforts include: removing golden eagles from the northern Channel Islands; reducing the threat of disease; breeding island foxes in captivity and reintroducing them to the wild; monitoring wild island fox populations; reintroducing bald eagles; and the removal of non-native species (e.g., non-native herbivores).

Long-term conservation of the subspecies will benefit from: conducting research on behavioral ecology and reproductive biology; increasing island fox education and outreach activities to reduce anthropogenic impacts; restoring island habitat; and assessing the **demographic** impact of other threats such as mortality from vehicles, competition with feral cats, and emerging disease issues (e.g., ear cancer). These are addressed under the Long-term Conservation Strategy section.

All known golden eagles have been removed from the northern Channel Islands and yet predation by golden eagles remains a potential threat to the long-term recovery of island fox populations, including the southern Channel Islands. Even one pair of nesting golden eagles appears to put significant pressure on island fox populations, whereas island fox populations can sustain some predation by transient birds, as occurred in 2007-2013 (see Sections 1.B.5. and 1.E). Thus, golden eagle monitoring and the removal of nesting birds need to continue, and management agencies need to be prepared to respond if and when new golden eagles nest on any of the Channel Islands. The successful reintroduction of bald eagles has resulted in the re-establishment of the first bald eagle nests on the northern Channel Islands in over 50 years and their continued presence is expected to be a long-term deterrent to the potential recolonization of the islands by golden eagles.

A subset of island foxes is vaccinated for CDV and rabies on all islands each year. However, resident and transient domestic dogs remain in contact with Santa Catalina Island foxes, and the potential for new disease transmission from dogs, cats, and other anthropogenic sources exists throughout the Channel Islands. Therefore, reducing the threat of disease will require avoiding introduction of new

Recovery Plan for Four Subspecies of Island Fox

pathogens or novel strains of existing pathogens to the Channel Islands and continued implementation of the epidemic response plans.

Captive breeding and reintroduction of all four endangered island fox subspecies has occurred on the Channel Islands as a means to provide a safe haven from predators and to augment the wild populations. Increasing the wild populations to levels with **vital rates** that minimize the risk of extinction is integral to island fox recovery. On-island captive breeding and reintroduction were conducted from 1999-2008 and ceased due to the success of reintroductions and the rapid growth of recovering populations. All foxes that were in captivity on Santa Catalina Island and the three northern Channel Islands have been reintroduced to the wild.

Timely threat detection and assessment of appropriate management actions is critical to maintain island foxes, which exist in a landscape that has been fundamentally altered by human impacts. Bakker and Doak (2009), provide guidance on the monitoring intensity needed to detect threats imposed by golden eagles and disease before those threats unduly impact island fox populations. We encourage frequent communication among the land managers in an effort to achieve the most cost-effective and rapid recovery of each island fox subspecies while standardizing recovery efforts as much as possible using the best available science and peer review. Management activities need to include:

- Monitoring island foxes over the long-term;
- Adapting as new information is gathered;
- Ensuring that population declines can be detected rapidly;
- Determining causes of decline; and
- Eliminating causes of decline as rapidly as is feasible.

The Channel Islands' ecosystems have been significantly altered and degraded over the past 2 centuries as a result of the introduction of non-native plant and animal species, unsustainable livestock grazing, and other anthropogenic activities (e.g., chemical pollution). Additional and increasing human impacts on all islands, such as additional proposed visitation on NPS land (NPS 2013), and increasing use of islands by private boaters who sometimes bring dogs and stowaway animals, may increase the likelihood of transferring diseases to the islands via intentional or unintentional introduction of mainland species. Due to the increasing risk of disease introduction on the islands, it is important that landowners and land managers retain the ability to use motorized vehicles to access, capture, and quarantine island foxes in remote areas.

Managed efforts to restore ecosystems (e.g., removal of invasive species on some islands, including deer, elk, pigs, sheep, rats, and cattle), will likely continue to affect the island ecosystems, with both positive and negative effects on island fox recovery.

III. Recovery Goals, Objectives, and Criteria

A. RECOVERY GOAL

The goal of this recovery plan is to recover the San Miguel Island fox, the Santa Rosa Island fox, the Santa Cruz Island fox, and the Santa Catalina Island fox so they can be delisted (removed from the List of Threatened and Endangered Species) when existing threats to each respective subspecies have been ameliorated such that their populations have been stabilized and have increased. The interim goal is to recover these subspecies to the point that they can be downlisted from endangered to threatened status.

B. RECOVERY OBJECTIVES

Recovery objectives are discrete targets that, when taken together, comprise the conditions that indicate a species may be warranted for delisting. Recovery objectives identify mechanisms for measuring progress toward and achieving the recovery goal.

Achieving the recovery goal requires: 1) increasing the population size and demographic rates to self-sustaining levels; and 2) reducing or eliminating the current threats to the survival of each subspecies.

1. Recovery Objective 1:

Each federally listed subspecies of island fox exhibits demographic characteristics consistent with long-term viability.

2. Recovery Objective 2:

Land managers are able to respond in a timely fashion to predation by nesting golden eagles or significant predation rates by transient golden eagles, to potential or incipient disease outbreaks, and to other identified threats using the best available technology.

For an island fox subspecies to be considered for downlisting from endangered to threatened status, recovery objective 1 is met.

For an island fox subspecies to be considered for delisting, recovery objective 1 and recovery objective 2 are met.

Each listed subspecies may be considered for downlisting or delisting independently of the other subspecies.

C. RECOVERY CRITERIA

An endangered species is defined in the Endangered Species Act as a species that is in danger of extinction throughout all or a significant portion of its range. A threatened species is one that is likely to become endangered within the foreseeable future throughout all or a significant portion of its range. When we evaluate whether or not a species warrants downlisting or delisting, we consider whether the species meets either of these definitions. A recovered species is one that no longer meets the Act's definitions of either threatened or endangered. Determining whether a species should be downlisted or delisted requires consideration of the of the same five categories of threats (*i.e.*, the five threat factors, A-E) which were considered when the species was listed and which are specified in section 4(a)(1) of the Endangered Species Act.

The island fox recovery criteria are measurable standards for determining whether an island fox subspecies has achieved its recovery objectives and may be considered for downlisting or delisting. The recovery criteria presented in this recovery plan represent our best assessment of the conditions that would most likely result in a determination that downlisting and/or delisting of the San Miguel Island fox, Santa Rosa Island fox, Santa Cruz Island fox, or the Santa Catalina Island fox is warranted. Achieving the prescribed recovery criteria is an indication that the species is no longer threatened or endangered. Because an actual change in status (downlisting or delisting) requires a separate rulemaking process that is based on a status assessment, including an analysis of the same five factors that were analyzed at listing, the Recovery Criteria below pertain to and are organized by these factors. Each Recovery Criterion applies to all four subspecies, except where noted otherwise.

Factor A: The present destruction, modification or curtailment of its habitat or range.

We believe that, if the threats under factors C and E are ameliorated, then the improvements in the habitat that are expected to occur with removal of herbivores responsible for habitat degradation may be a long-term benefit to the island fox, but is not necessary for recovery. Therefore, we are not proposing recovery criteria under this factor.

Factor B: Overutilization for commercial, scientific or educational purposes.

Overutilization is not currently known to be a threat for this species. Therefore, no recovery criteria are necessary for this factor.

Factor C: Disease or predation.

To address recovery objective 2, disease and predation pressures must be reduced. This will have been accomplished if the following have occurred:

C/1: Golden eagle predation:

a. To reduce the threat of extinction to the San Miguel Island fox, the Santa Rosa Island fox, and the Santa Cruz Island fox, the rate of golden eagle predation is reduced and maintained at a level that is no longer considered a threat to island fox recovery through development of a golden eagle management strategy. The strategy will be developed by the land manager(s) in consultation with the FWS and will include review by the appropriate IRT Technical Expertise Group or the equivalent. This strategy includes:

- Response tactics (including the use of helicopters and net-guns) to capture nesting golden eagles and any transient golden eagle responsible for significant island fox predation per the golden eagle response strategy.
- Tactics to minimize the establishment of successful nesting golden eagles;
- An established island fox monitoring program that is able to detect an annual island fox predation rate caused by golden eagles of 2.5 percent or greater, averaged over 3 years (Bakker and Doak 2009); and
- An established mortality rate or population size threshold that, if reached due to golden eagle predation, would require the land manager(s) to bring island foxes into captivity for safety.

b. The golden eagle prey base of deer and elk is removed from Santa Rosa Island.

At present, golden eagles are not known to prey upon Santa Catalina Island foxes. If mortality as a result of golden eagle predation becomes a threat to the Santa Catalina Island fox, implement the above measures as necessary.

C/2: A disease management strategy is developed, approved, and implemented by the land manager(s) in consultation with the FWS and includes review by the appropriate IRT TEG or the equivalent. This strategy includes:

Recovery Plan for Four Subspecies of Island Fox

- Identification of a portion of each population that will be vaccinated against diseases posing the greatest risk for which vaccines are safe and effective. Vaccinations to be provided and numbers vaccinated will be developed in consultation with appropriate subject-matter experts;
- Identification of actual and potential pathogens of island foxes, and the means by which these can be prevented from decimating fox populations;
- Disease prevention;
- A monitoring program that provides for timely detection of a disease outbreak, and an associated emergency response strategy as recommended by the appropriate subject-matter experts; and
- A process for updating the disease strategy as new information arises.

Factor D: Inadequacy of existing regulatory mechanisms.

We believe that if the threats under factors C and E are ameliorated, then additional regulatory mechanisms (beyond existing ones) are not necessary. Therefore, we are not proposing Recovery Criteria under this factor.

Factor E: Other natural or manmade factors affecting its continued existence.

To address recovery objective 1 for each of the four subspecies, the subspecies must be protected from other natural or manmade factors known to affect their continued existence. This will have been accomplished if the following has occurred:

E/1: An island fox subspecies has no more than 5 percent risk of quasi-extinction over a 50-year period (addresses objective 1). This risk level is based on the following:

- Quasi-extinction is defined as a population size of ≤ 30 individuals.
- The risk of extinction is calculated based on the combined lower 80 percent confidence interval for a 3 year running average of population size estimates, and the upper 80 percent confidence interval for a 3 year running average of mortality rate estimates.
- This 5 percent (or less) risk level is sustained for at least 5 years, during which time the population trend is not declining. A declining

Recovery Plan for Four Subspecies of Island Fox

trend is defined as the 3-year risk-level being greater in year 5 than year 1.

This risk-based recovery criterion is based on models developed separately for each listed subspecies. A description of the models can be found in Appendix 2.

Recovery Plan for Four Subspecies of Island Fox

IV. Recovery Program

A. RECOVERY ACTION NARRATIVE

The actions identified below are those that, in our opinion, are necessary to bring about the recovery of island foxes. These actions are subject to modification as dictated by new findings, changes in species status, and the completion of recovery actions. Each action has been assigned a priority as follows:

Priority 1: An action that is taken to prevent extinction or to prevent the species from declining irreversibly.

Priority 2: An action that is taken to prevent a significant decline in species population/habitat quality or some other significant negative impact short of extinction.

Priority 3: All other actions necessary to provide for full recovery of the species.

1. Reduce mortality and maintain productivity for each subspecies of island fox to sustainable levels.

Two major mortality factors have been identified, golden eagle predation and disease. Therefore, most actions identified below address these two factors.

1.1. Reduce rate of golden eagle predation and maintain at a level that is no longer considered a threat to island fox recovery. Implement and maintain an active monitoring/response program for golden eagles as needed.

1.1.1. Develop and implement a formal golden eagle management strategy.

This should include plans for monitoring, control, removal, and contingency in case of golden eagle return after removal. The golden eagle management strategy should have the flexibility to adapt to new information and changing conditions, and to evaluate all known means of capturing golden eagles, including helicopter use in remote areas, or suppressing their ability to prey on island foxes.

1.1.1.1. Develop and implement a formal golden eagle management strategy for San Miguel Island (Priority 1).

1.1.1.2. Develop and implement a formal golden eagle management strategy for Santa Rosa Island (Priority 1).

Recovery Plan for Four Subspecies of Island Fox

- 1.1.1.3. Develop and implement a formal golden eagle management strategy for Santa Cruz Island (Priority 1).

- 1.1.2. Monitor for golden eagle activity.

Conduct annual monitoring at minimum during the nesting season to detect any resident golden eagles. Monitoring should include aerial and ground surveys as needed and training for all field staff to identify and report all eagle sightings. Maximize all opportunities to locate golden eagles whenever any field activities are undertaken.

- 1.1.2.1. Monitor for golden eagle activity on San Miguel Island (Priority 2).

- 1.1.2.2. Monitor for golden eagle activity on Santa Rosa Island (Priority 2).

- 1.1.2.3. Monitor for golden eagle activity on Santa Cruz Island (Priority 2).

- 1.1.3. Remove golden eagles to maintain the Channel Islands free of resident golden eagles.

If golden eagles are seen or signs are found of their presence, steps should be taken to determine whether capture and removal to the mainland is necessary. Continue to consult with eagle experts for additional techniques to capture and/or manage golden eagles. Due to the proven difficulty of capturing golden eagles (Latta et al. 2005) all options for capturing golden eagle should be considered. Improve golden eagle capture by developing and implementing new methods and new technologies.

Continue golden eagle trapping and removal efforts until all resident golden eagles have been removed from the northern Channel Islands.

- 1.1.3.1. Complete initial removal of golden eagles from northern Channel Islands.

Continue golden eagle trapping and removal efforts until all resident golden eagles have been removed from the northern Channel Islands.

- 1.1.3.1.1. Complete initial removal of golden eagles from San Miguel Island (Priority 1).

Recovery Plan for Four Subspecies of Island Fox

- 1.1.3.1.2. Complete initial removal of golden eagles from Santa Rosa Island (Priority 1).
- 1.1.3.1.3. Complete initial removal of golden eagles from Santa Cruz Island (Priority 1).
- 1.1.3.2. Control resident golden eagles on the Channel Islands, as needed, after 1.1.3.1. above is complete to sustain island fox populations.
 - 1.1.3.2.1. Control resident golden eagles on the Channel Islands, as needed, to sustain island fox populations on San Miguel Island (Priority 1).
 - 1.1.3.2.2. Control resident golden eagles on the Channel Islands, as needed, to sustain island fox populations on Santa Rosa Island (Priority 1).
 - 1.1.3.2.3. Control resident golden eagles on the Channel Islands, as needed, to sustain island fox populations on Santa Cruz Island (Priority 1).
- 1.1.3.3. Identify and manage any activities or food sources that are attractants for golden eagles. Minimize the availability of food resources for golden eagles to inhibit successful establishment of territories/reproduction and to direct eagles toward capture baits. Conduct additional removals of golden eagles from any island as needed.
 - 1.1.3.3.1. On San Miguel Island, identify and manage any activities or food sources that are attractants for golden eagles. Minimize the availability of food resources for golden eagles to inhibit successful establishment of territories/reproduction and to direct eagles toward capture baits. Conduct additional removals of golden eagles as needed (Priority 3).
 - 1.1.3.3.2. On Santa Rosa Island, identify and manage any activities or food sources that are attractants for golden eagles. Minimize the availability of food resources for golden eagles to inhibit successful establishment of territories/reproduction and to direct eagles toward capture baits. Conduct additional

Recovery Plan for Four Subspecies of Island Fox

removals of golden eagles as needed (Priority 3).

1.1.3.3.3. On Santa Cruz Island, identify and manage any activities or food sources that are attractants for golden eagles. Minimize the availability of food resources for golden eagles to inhibit successful establishment of territories/reproduction and to direct eagles toward capture baits. Conduct additional removals of golden eagles as needed (Priority 3).

1.1.3.4. Conduct research needed to understand and eliminate golden eagle residency on the Channel Islands.

Such research could include food habit studies and genetic analyses to determine how frequently golden eagles immigrate from the mainland.

1.1.3.4.1. Conduct research needed to understand and eliminate golden eagle residency on San Miguel Island (Priority 3).

1.1.3.4.2. Conduct research needed to understand and eliminate golden eagle residency on Santa Rosa Island (Priority 3).

1.1.3.4.3. Conduct research needed to understand and eliminate golden eagle residency on Santa Cruz Island (Priority 3).

1.2. Avoid introduction of new pathogens, or novel strains of existing pathogens, to the Channel Islands by restricting or regulating movements of wild and domestic animals to the islands.

The small size of island fox populations and genetic homogeneity means that infectious disease has an unusually high potential to cause population crashes or even extinction. Island foxes have a history of exposure to infectious disease, but may be immunologically naïve to pathogen strains that are endemic to the mainland but absent from the Channel Islands.

Additional details and guidance for this recovery action provided by the island fox health TEG can be found in Appendix 3.

Recovery Plan for Four Subspecies of Island Fox

- 1.2.1. The ban on bringing pets to Channel Islands National Park, and to TNC land on Santa Cruz Island, should be well-publicized and strictly enforced.
 - 1.2.1.1. The ban on bringing pets to Channel Islands National Park-San Miguel Island should be well-publicized and strictly enforced (Priority 1).
 - 1.2.1.2. The ban on bringing pets to Channel Islands National Park-Santa Rosa Island should be well-publicized and strictly enforced (Priority 1).
 - 1.2.1.3. The ban on bringing pets to Channel Islands National Park, and to TNC land on Santa Cruz Island, should be well-publicized and strictly enforced (Priority 1).
- 1.2.2. Where there is a clear benefit to bringing domestic dogs to the northern Channel Islands, the quarantine guidelines established for dogs brought to Santa Cruz Island to assist with pig eradication efforts should be followed (see Appendix 6).
 - 1.2.2.1. Where there is a clear benefit to bringing domestic dogs to San Miguel Island, the quarantine guidelines established for dogs brought to Santa Cruz Island to assist with pig eradication efforts need to be followed (see Appendix 6) (Priority 1).
 - 1.2.2.2. Where there is a clear benefit to bringing domestic dogs to Santa Rosa Island, the quarantine guidelines established for dogs brought to Santa Cruz Island to assist with pig eradication efforts need to be followed (see Appendix 6) (Priority 1).
 - 1.2.2.3. Where there is a clear benefit to bringing domestic dogs to Santa Cruz Island, the quarantine guidelines established for dogs brought to Santa Cruz Island to assist with pig eradication efforts need to be followed (see Appendix 6) (Priority 1).
- 1.2.3. Movement of non-native species or carcasses to the northern Channel Islands should be avoided wherever possible.
 - 1.2.3.1. Movement of other mammals or carcasses to San Miguel Island should be avoided wherever possible (Priority 3).
 - 1.2.3.2. Movement of other mammals or carcasses to Santa Rosa Island should be avoided wherever possible (Priority 3).

Recovery Plan for Four Subspecies of Island Fox

- 1.2.3.3. Movement of other mammals or carcasses to Santa Cruz Island should be avoided wherever possible (Priority 3).
- 1.2.4. The potential for pathogen introduction to Santa Catalina Island from the movement of wild and domestic mammals should be reduced to the extent practicable (Priority 3).
- 1.2.5. Develop a management strategy for responding to new introductions of animals to the Channel Islands.
 - 1.2.5.1. Develop a management strategy for responding to new introductions of animals to San Miguel Island (Priority 3).
 - 1.2.5.2. Develop a management strategy for responding to new introductions of animals to Santa Rosa Island (Priority 3).
 - 1.2.5.3. Develop a management strategy for responding to new introductions of animals to Santa Cruz Island (Priority 3).
 - 1.2.5.4. Develop a management strategy for responding to new introductions of animals to Santa Catalina Island (Priority 3).
- 1.3. Implement preventative management to avoid extinction or quasi-extinction of wild populations in the event of devastating epidemics.

PVA models suggest that the probability of extinction in the face of a rabies or CDV epidemic could be substantially reduced by maintaining a “vaccinated core” of animals. This approach involves maintaining a small number of animals protected from infection by vaccination. These animals act as a “safety net,” intended to survive any epidemics that occur and then to form a founder group from which subsequent recovery may occur. PVA models suggest that, assuming vaccination is 100 percent protective, maintaining a “vaccinated core” of 80 to 100 vaccinated individuals per island fox subspecies dramatically reduces the probability of population extinction, even when there is a comparatively high (10 percent) probability of a rabies epidemic in any one year (Schwemm 2007).

Additional details and guidance for this recovery action provided by the island fox health TEG can be found in Appendix 3.

- 1.3.1. Test safety of, and antibody response to, vaccination in captive island foxes under appropriate research protocols.

Recovery Plan for Four Subspecies of Island Fox

- 1.3.1.1. Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on island foxes during a single vaccination event.
 - 1.3.1.1.1. Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on San Miguel Island foxes during a single vaccination event (Priority 1).
 - 1.3.1.1.2. Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on Santa Rosa Island foxes during a single vaccination event (Priority 1).
 - 1.3.1.1.3. Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on Santa Cruz Island foxes during a single vaccination event (Priority 1).
 - 1.3.1.1.4. Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on Santa Catalina Island foxes during a single vaccination event (Priority 1).
- 1.3.1.2. Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in island foxes.
 - 1.3.1.2.1. Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in San Miguel Island foxes (Priority 1).
 - 1.3.1.2.2. Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in Santa Rosa Island foxes (Priority 1).
 - 1.3.1.2.3. Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in Santa Cruz Island foxes (Priority 1).
 - 1.3.1.2.4. Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in Santa Catalina Island foxes (Priority 1).

Recovery Plan for Four Subspecies of Island Fox

- 1.3.1.3. Vaccines against canine parvovirus and adenovirus should be tested on island foxes.
 - 1.3.1.3.1. Vaccines against canine parvovirus and adenovirus should be tested on San Miguel Island foxes (Priority 3).
 - 1.3.1.3.2. Vaccines against canine parvovirus and adenovirus should be tested on Santa Rosa Island foxes (Priority 3).
 - 1.3.1.3.3. Vaccines against canine parvovirus and adenovirus should be tested on Santa Cruz Island foxes (Priority 3).
 - 1.3.1.3.4. Vaccines against canine parvovirus and adenovirus should be tested on Santa Catalina Island foxes (Priority 3).

- 1.3.2. On each island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes.

On islands where wild fox populations number fewer than 100 individuals, all island foxes should be vaccinated. On Santa Cruz Island, vaccination should be focused in one or two localized areas, but specifically in primary access corridors, such as Prisoner's Harbor up canyon to the central valley. On Santa Catalina Island, vaccination efforts should be concentrated around the city of Avalon (where disease introduction is most likely to occur) and around the isthmus (where infection could potentially pass between the eastern and western subpopulations). However, because of the many points of access to Santa Catalina Island, island-wide vaccination is preferred.

When a vaccine is first introduced, a proportion of vaccinated individuals should be radio-collared on each island to allow determination of whether vaccination has any negative consequences for island foxes in the absence of an epidemic.

- 1.3.2.1. On San Miguel Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes (Priority 1).
- 1.3.2.2. On Santa Rosa Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes (Priority 1).

Recovery Plan for Four Subspecies of Island Fox

- 1.3.2.3. On Santa Cruz Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes (Priority 1).
 - 1.3.2.4. On Santa Catalina Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes (Priority 1).
- 1.4. Establish monitoring and response strategies to detect and manage infectious disease threats to island fox population persistence.

Additional details and guidance for this recovery action provided by the island fox health TEG can be found in Appendix 3 and 4.

1.4.1. Monitor to detect disease-related mortality.

- 1.4.1.1. Using radio-telemetry, monitor a sample of foxes on each island to detect fox mortalities.

At least 10 island foxes selected for radio-collars as part of routine monitoring (as opposed to after a vaccine is first introduced) should not be vaccinated; this allows them to act as sentinels of infection, allowing early detection of future epidemics.

See Recovery Action Section 3.0 and Appendices 3 and 4 for more details.

- 1.4.1.1.1. Using radio-telemetry, monitor a sample of foxes on San Miguel Island to detect fox mortalities (Priority 1).
 - 1.4.1.1.2. Using radio-telemetry, monitor a sample of foxes on Santa Rosa Island to detect fox mortalities (Priority 1).
 - 1.4.1.1.3. Using radio-telemetry, monitor a sample of foxes on Santa Cruz Island to detect fox mortalities (Priority 1).
 - 1.4.1.1.4. Using radio-telemetry, monitor a sample of foxes on Santa Catalina Island to detect fox mortalities (Priority 1).
- 1.4.1.2. Any island foxes that are found dead should be collected and shipped or frozen immediately for necropsy.

Recovery Plan for Four Subspecies of Island Fox

- 1.4.1.2.1. Any San Miguel Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy (Priority 1).
- 1.4.1.2.2. Any Santa Rosa Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy (Priority 1).
- 1.4.1.2.3. Any Santa Cruz Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy (Priority 1).
- 1.4.1.2.4. Any Santa Catalina Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy (Priority 1).
- 1.4.1.3. Any island fox appearing ill or acting in an abnormal manner should be reported immediately, quarantined, and closely monitored.
 - 1.4.1.3.1. Any San Miguel Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored (Priority 1).
 - 1.4.1.3.2. Any Santa Rosa Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored (Priority 1).
 - 1.4.1.3.3. Any Santa Cruz Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored (Priority 1).
 - 1.4.1.3.4. Any Santa Catalina Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored (Priority 1).
- 1.4.1.4. Other carnivores sick or dead should be reported and closely monitored (if alive) or collected for necropsy (if dead).
 - 1.4.1.4.1. Other carnivores on San Miguel Island found sick or dead from causes other than trauma should be reported and closely monitored (if

Recovery Plan for Four Subspecies of Island Fox

- alive) or collected for necropsy (if dead) (Priority 3).
 - 1.4.1.4.2. Other carnivores on Santa Rosa Island found sick or dead from causes other than trauma should be reported and closely monitored (if alive) or collected for necropsy (if dead) (Priority 3).
 - 1.4.1.4.3. Other carnivores on Santa Cruz Island found sick or dead from causes other than trauma should be reported and closely monitored (if alive) or collected for necropsy (if dead) (Priority 3).
 - 1.4.1.4.4. Other carnivores on Santa Catalina Island found sick or dead from causes other than trauma should be reported and closely monitored (if alive) or collected for necropsy (if dead) (Priority 3).
- 1.4.2. Annually collect blood samples from a proportion of island foxes on all islands to evaluate ongoing disease risks to island fox populations and conduct a serosurvey for antibodies to CDV, CPV, CAV, and Toxoplasma as often as yearly, but no less often than every 5 years.
- 1.4.2.1. Annually collect blood samples from a proportion of island foxes on San Miguel Island to evaluate ongoing disease risks to island fox population (Priority 2).
 - 1.4.2.2. Annually collect blood samples from a proportion of island foxes on Santa Rosa Island to evaluate ongoing disease risks to island fox population (Priority 2).
 - 1.4.2.3. Annually collect blood samples from a proportion of Santa Cruz Island foxes to evaluate ongoing disease risks to island fox population (Priority 2).
 - 1.4.2.4. Annually collect blood samples from a proportion of island foxes on Santa Catalina Island to evaluate ongoing disease risks to the island fox population (Priority 2).
- 1.4.3. Develop strategies for responding to island fox deaths from infectious diseases known to represent serious threats to the persistence or recovery of the wild populations.

Recovery Plan for Four Subspecies of Island Fox

- 1.4.3.1. All island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes.
 - 1.4.3.1.1. San Miguel Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes (Priority 1).
 - 1.4.3.1.2. Santa Rosa Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes (Priority 1).
 - 1.4.3.1.3. Santa Cruz Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes (Priority 1).
 - 1.4.3.1.4. Santa Catalina Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes (Priority 1).
- 1.4.3.2. A single case of rabies, confirmed by pathology or virus isolation, should trigger management response strategy.
 - 1.4.3.2.1. A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for San Miguel Island foxes (Priority 1).
 - 1.4.3.2.2. A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for Santa Rosa Island foxes (Priority 1).
 - 1.4.3.2.3. A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for Santa Cruz Island foxes (Priority 1).
 - 1.4.3.2.4. A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for Santa Catalina Island foxes (Priority 1).

Recovery Plan for Four Subspecies of Island Fox

- 1.4.3.3. A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response strategy.
 - 1.4.3.3.1. A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for San Miguel Island foxes (Priority 1).
 - 1.4.3.3.2. A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for Santa Rosa Island foxes (Priority 1).
 - 1.4.3.3.3. A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for Santa Cruz Island foxes (Priority 1).
 - 1.4.3.3.4. A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for Santa Catalina Island foxes (Priority 1).
- 1.4.3.4. A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead island foxes.
 - 1.4.3.4.1. A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead San Miguel Island foxes (Priority 2).
 - 1.4.3.4.2. A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead Santa Rosa Island foxes (Priority 2).
 - 1.4.3.4.3. A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead Santa Cruz Island foxes (Priority 2).

Recovery Plan for Four Subspecies of Island Fox

- 1.4.3.4.4. A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead Santa Catalina Island foxes (Priority 2).
- 1.5. Conduct research to understand and evaluate the threats to island foxes posed by other infectious and noninfectious diseases, and develop management strategies.
 - 1.5.1. Complete on-going investigations of the demographic consequences and etiology of the ear tumors prevalent in Santa Catalina Island foxes to determine whether this disease poses a significant threat to this fox population (Priority 2).
 - 1.5.2. Expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted.
 - 1.5.2.1. On San Miguel Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted (Priority 2).
 - 1.5.2.2. On Santa Rosa Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted (Priority 2).
 - 1.5.2.3. On Santa Cruz Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted (Priority 2).
 - 1.5.2.4. On Santa Catalina Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted (Priority 2).
 - 1.5.3. Conduct further research as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of

Recovery Plan for Four Subspecies of Island Fox

pathological and demographic analyses, to threaten island fox populations.

- 1.5.3.1. Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the San Miguel Island fox population (Priority 2).
- 1.5.3.2. Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the Santa Rosa Island fox population (Priority 2).
- 1.5.3.3. Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the Santa Cruz Island fox population (Priority 2).
- 1.5.3.4. Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the Santa Catalina Island fox population (Priority 2).

2. Manage captive island fox populations for recovery.

On-island captive breeding and reintroduction were conducted from 1999-2008 and ceased due to the success of reintroductions and the rapid growth of recovering populations. Captive populations of island foxes were critical to the species' conservation and can continue to be important in the recovery and long-term conservation (see Long-term Conservation section) of the four listed subspecies. Captive breeding must be conducted in accordance with the FWS Captive Propagation Policy. Foxes were initially brought into captivity to prevent extinction from the threat of golden eagle predation on the northern Channel Islands, and the threat of disease on Santa Catalina Island. Captive reproduction has ensured that the island foxes did not go extinct and has contributed to recovery of wild populations by providing individuals for release as the current threats have been brought under control. Captive breeding can be phased out as wild populations recover, but the long-term persistence of island foxes (see Long-term Conservation section) may benefit from redundant, genetically diverse, and sustainable mainland populations of one or two subspecies. Manage the on-island captive populations of island foxes to augment the wild populations.

Recovery Plan for Four Subspecies of Island Fox

- 2.1. Continue captive management of island foxes as necessary to provide individuals for release.
 - 2.1.1. Continue captive management of San Miguel Island foxes as necessary to provide individuals for release (Priority 3).
 - 2.1.2. Continue captive management of Santa Rosa Island foxes as necessary to provide individuals for release (Priority 3).
 - 2.1.3. Continue captive management of Santa Cruz Island foxes as necessary to provide individuals for release (Priority 3).
 - 2.1.4. Continue captive management of Santa Catalina Island foxes as necessary to provide individuals for release (Priority 3).
- 2.2. Assuming golden eagle predation and disease risks remain low, continue annual release of island foxes from the captive facilities until such releases are no longer necessary to augment wild populations.
 - 2.2.1. Assuming golden eagle predation and disease risks remain at low levels, continue annual release of San Miguel Island foxes from the captive facility until such releases are no longer necessary to augment the wild population (Priority 3).
 - 2.2.2. Assuming golden eagle predation and disease risks remain at low levels, continue annual release of Santa Rosa Island foxes from the captive facility until such releases are no longer necessary to augment the wild population (Priority 3).
 - 2.2.3. Assuming golden eagle predation and disease risks remain at low levels, continue annual release of Santa Cruz Island foxes from the captive facility until such releases are no longer necessary to augment the wild population (Priority 3).
 - 2.2.4. Assuming golden eagle predation and disease risks remain at low levels, continue annual release of Santa Catalina Island foxes from the captive facility until such releases are no longer necessary to augment the wild population (Priority 3).
- 2.3. Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining island fox captive populations is retained while captive breeding is ongoing.
 - 2.3.1. Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which San Miguel Island fox individuals to release and

Recovery Plan for Four Subspecies of Island Fox

which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing (Priority 3).

2.3.2. Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which Santa Rosa Island fox individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing (Priority 3).

2.3.3. Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which Santa Cruz Island fox individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing (Priority 3).

2.3.4. Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which Santa Catalina Island fox individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing (Priority 3).

2.4. Continue to monitor released island foxes and use this information to modify release strategies (e.g., release locations and timing).

2.4.1. Continue to monitor released San Miguel Island foxes and use this information to modify release strategies (e.g., release locations and timing) (Priority 2).

2.4.2. Continue to monitor released Santa Rosa Island foxes and use this information to modify release strategies (e.g., release locations and timing) (Priority 2).

2.4.3. Continue to monitor released Santa Cruz Island foxes and use this information to modify release strategies (e.g., release locations and timing) (Priority 2).

2.4.4. Continue to monitor released Santa Catalina Island foxes and use this information to modify release strategies (e.g., release locations and timing) (Priority 2).

2.5. Maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat.

Recovery Plan for Four Subspecies of Island Fox

- 2.5.1. On San Miguel Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat (Priority 3).
 - 2.5.2. On Santa Rosa Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat (Priority 3).
 - 2.5.3. On Santa Cruz Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat (Priority 3).
 - 2.5.4. On Santa Catalina Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat (Priority 3).
- 2.6. While any on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes.
- 2.6.1. While San Miguel Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes (Priority 3).
 - 2.6.2. While Santa Rosa Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes (Priority 3).
 - 2.6.3. While Santa Cruz Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes (Priority 3).
 - 2.6.4. While Santa Catalina Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes (Priority 3).
- 2.7. Continue to maintain an island fox studbook and continue to use the studbook to aid decisions on fox pairing and release.

Recovery Plan for Four Subspecies of Island Fox

- 2.7.1. Continue to maintain a San Miguel Island fox studbook and continue to use the studbook to aid decisions on fox pairing and release (Priority 2).
- 2.7.2. Continue to maintain a Santa Rosa Island fox studbook and continue to use the studbook to aid decisions on fox pairing and release (Priority 2).
- 2.7.3. Continue to maintain a Santa Cruz Island fox studbook and continue to use the studbook to aid decisions on fox pairing and release (Priority 2).

3. Establish island fox monitoring strategies.

Monitoring of island fox populations has been, and will continue to be, a crucial activity. Given the inherent risk of small insular populations, robust monitoring of island fox populations and their threats is a key component of recovery and long-term conservation. Such long-term monitoring strategies should incorporate the best established methods to track population dynamics and to detect and understand the causes of population declines in a timely manner. To that end, an effective monitoring strategy should be able to address each of the following monitoring objectives:

- Tracking the status of island fox recovery, particularly relative to recovery criteria;
 - Guiding island-specific management decisions in a timely manner;
 - Refining parameter estimates for population viability analyses and facilitating cross-island comparisons; and
 - Monitoring to detect a potential future and/or current catastrophic population decline.
- 3.1. Develop and implement a monitoring strategy for each listed island fox subspecies to detect population declines and determine population trends.

Monitoring parameters need to be targeted for the purpose of tracking and determining recovery. These parameters include:

- Mortality rates (with associated cause-specific mortality rates);
- Population trend (e.g., λ , lambda); and
- Population size, possibly relative to a required minimum population size as informed by PVA.

These parameters would provide information necessary for the evaluation of extinction risk based upon the PVA and aid in determining the current

Recovery Plan for Four Subspecies of Island Fox

level of risk to an island fox population (Recovery Criterion 1) as well as aiding in determining the progress towards recovery. These parameters were presented and reviewed at the second PVA Workshop convened at University of California, Davis in December 2006, including refinement of parameters in the context of recovery criteria, to identify desired precision levels. Based on this workshop, the following parameters and associated precision levels have been chosen for the purpose of tracking and determining recovery:

- Annual estimate of island-wide population size, with an 80 percent confidence interval.
 - Annual estimate of mortality, with an 80 percent confidence interval and cause-specific mortality rates sufficient to detect a rate of eagle predation of 2.5 percent or greater (Bakker and Doak 2009). In addition, these data would provide a means of monitoring for disease outbreaks and facilitate health research and vaccine efficacy tests.
 - Estimate of trend in population size, which can be estimated either from annual abundance estimates or from population models. This estimate has no targeted precision; rather the precision of the trend estimate would be determined by the precision of the population estimates and possibly by precision of mortality rates (see Appendix 5).
- 3.1.1. Develop and implement a monitoring strategy for San Miguel Island fox subspecies to detect population declines and determine population trends (Priority 1).
 - 3.1.2. Develop and implement a monitoring strategy for Santa Rosa Island fox subspecies to detect population declines and determine population trends (Priority 1).
 - 3.1.3. Develop and implement a monitoring strategy for Santa Cruz Island fox subspecies to detect population declines and determine population trends (Priority 1).
 - 3.1.4. Develop and implement a monitoring strategy for Santa Catalina Island fox subspecies to detect population declines and determine population trends (Priority 1).
- 3.2. Ensure island fox population information is comparable across the islands to the greatest extent possible.

Recommendations for achieving this include collecting, storing, and managing data using standardized protocols.

Recovery Plan for Four Subspecies of Island Fox

- 3.2.1. Ensure San Miguel Island fox population information is comparable across islands to the greatest extent possible (Priority 3).
- 3.2.2. Ensure Santa Rosa Island fox population information is comparable across islands to the greatest extent possible (Priority 3).
- 3.2.3. Ensure Santa Cruz Island fox population information is comparable across islands to the greatest extent possible (Priority 3).
- 3.2.4. Ensure Santa Catalina Island fox population information is comparable across islands to the greatest extent possible (Priority 3).

Recovery Plan for Four Subspecies of Island Fox

V. Implementation Schedule

The following implementation schedule outlines actions and estimated costs for this recovery plan. It is a guide for meeting the objectives discussed in Parts II, III, and IV of this recovery plan. This schedule describes and prioritizes actions, provides an estimated timetable for performance of actions, indicates the responsible parties, and estimates costs of performing actions. These actions when accomplished, should further recovery and conservation of the covered subspecies.

Key to terms and acronyms used in Implementation Schedule:

Definition of Action Durations:

- Number: The predicted duration of the action in years or the cost of the action.
Ongoing: An action that is currently being implemented and will continue throughout the recovery period.
Continual: An action that is not currently being implemented but will be implemented continuously throughout the recovery period once begun.
Unknown: Either action duration or associated costs are not known at this time.

Definition of Subspecies Benefitting:

- SMIF: San Miguel Island Fox
SRIF: Santa Rosa Island Fox
SCZIF: Santa Cruz Island Fox
SCIF: Santa Catalina Island Fox

Responsible Parties:

- AZA Association of Zoos and Aquariums
CDFW California Department of Fish and Wildlife
CIC Santa Catalina Island Conservancy
FWS U.S. Fish and Wildlife Service
NPS National Park Service
TNC The Nature Conservancy
UNIV University or academic researchers

Responsible parties are those entities who may voluntarily participate in any aspect of implementation of a particular tasks listed within this recovery plan. Responsible parties may willingly participate in project planning, funding, staff time, or any other means of implementation.

Recovery Plan for Four Subspecies of Island Fox

Table 3. Implementation schedule for the recovery plan for four subspecies of island fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	1.1.1.1.	Develop and implement a formal golden eagle management strategy for San Miguel Island.	SMIF	Ongoing	FWS NPS	14	14				
1	1.1.1.2.	Develop and implement a formal golden eagle management strategy for Santa Rosa Island.	SRIF	Ongoing	FWS NPS	14	14				
1	1.1.1.3.	Develop and implement a formal golden eagle management strategy for Santa Cruz Island.	SCZIF	Ongoing	FWS NPS TNC	14	14				
2	1.1.2.1.	Monitor for golden eagle activity on San Miguel Island.	SMIF	Ongoing	FWS NPS	10	2	2	2	2	2
2	1.1.2.2.	Monitor for golden eagle activity on Santa Rosa Island.	SRIF	Ongoing	FWS NPS	10	2	2	2	2	2
2	1.1.2.3.	Monitor for golden eagle activity on Santa Cruz Island.	SCZIF	Ongoing	FWS NPS TNC	10	2	2	2	2	2
1	1.1.3.1.1.	Complete initial removal of golden eagles from San Miguel Island.	SMIF	Ongoing	FWS NPS						
1	1.1.3.1.2.	Complete initial removal of golden eagles from Santa Rosa Island.	SRIF	Ongoing	FWS NPS						
1	1.1.3.1.3.	Complete initial removal of golden eagles from Santa Cruz Island.	SCZIF	Ongoing	FWS NPS TNC						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	1.1.3.2.1.	Control resident golden eagles on the Channel Islands, as needed, to sustain island fox populations on San Miguel Island.	SMIF	Ongoing	FWS NPS	17.5	3.5	3.5	3.5	3.5	3.5
1	1.1.3.2.2.	Control resident golden eagles on the Channel Islands, as needed, to sustain island fox populations on Santa Rosa Island.	SRIF	Ongoing	FWS NPS	17.5	3.5	3.5	3.5	3.5	3.5
1	1.1.3.2.3.	Control resident golden eagles on the Channel Islands, as needed, to sustain island fox populations on Santa Cruz Island.	SCZIF	Ongoing	FWS NPS TNC	17.5	3.5	3.5	3.5	3.5	3.5
3	1.1.3.3.1.	On San Miguel Island, identify and manage any activities or food sources that are attractants for golden eagles. Minimize the availability of food resources for golden eagles to inhibit successful establishment of territories/reproduction and to direct eagles toward capture baits. Conduct additional removals of golden eagles as needed	SMI	Ongoing	FWS NPS						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	1.1.3.3.2.	On Santa Rosa Island, identify and manage any activities or food sources that are attractants for golden eagles. Minimize the availability of food resources for golden eagles to inhibit successful establishment of territories/reproduction and to direct eagles toward capture baits. Conduct additional removals of golden eagles as needed	SRI	Ongoing	FWS NPS						
3	1.1.3.3.3.	On Santa Cruz Island, identify and manage any activities or food sources that are attractants for golden eagles. Minimize the availability of food resources for golden eagles to inhibit successful establishment of territories/reproduction and to direct eagles toward capture baits. Conduct additional removals of golden eagles as needed	SCZIF	Ongoing	FWS NPS TNC						
3	1.1.3.4.1.	Conduct research needed to understand and eliminate golden eagle residency on San Miguel Island.	SMIF	Ongoing	FWS NPS UNIV						
3	1.1.3.4.2.	Conduct research needed to understand and eliminate golden eagle residency on Santa Rosa Island.	SRIF	Ongoing	FWS NPS UNIV						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	1.1.3.4.3.	Conduct research needed to understand and eliminate golden eagle residency on Santa Cruz Island.	SCZIF	Ongoing	FWS NPS TNC UNIV						
1	1.2.1.1.	The ban on bringing pets to Channel Islands National Park-San Miguel Island should be well-publicized and strictly enforced.	SMIF	Ongoing	NPS FWS	12.5	2.5	2.5	2.5	2.5	2.5
1	1.2.1.2.	The ban on bringing pets to Channel Islands National Park-Santa Rosa Island should be well-publicized and strictly enforced.	SRIF	Ongoing	NPS FWS	12.5	2.5	2.5	2.5	2.5	2.5
1	1.2.1.3.	The ban on bringing pets to Channel Islands National Park, and to TNC land on Santa Cruz Island, should be well-publicized and strictly enforced.	SCZIF	Ongoing	NPS TNC FWS	12.5	2.5	2.5	2.5	2.5	2.5
1	1.2.2.1.	Where there is a clear benefit to bringing domestic dogs to San Miguel Island, the quarantine guidelines established for dogs brought to Santa Cruz Island to assist with pig eradication efforts need to be followed (see Appendix 6).	SMIF	Unknown	NPS FWS	TBD					
1	1.2.2.2.	Where there is a clear benefit to bringing domestic dogs to Santa Rosa Island, the quarantine guidelines established for dogs brought to Santa Cruz Island to assist with pig eradication efforts need to be followed (see Appendix 6).	SRIF	Unknown	NPS FWS	TBD					

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	1.2.2.3.	Where there is a clear benefit to bringing domestic dogs to Santa Cruz Island, the quarantine guidelines established for dogs brought to Santa Cruz Island to assist with pig eradication efforts need to be followed (see Appendix 6).	SCZIF	Unknown	NPS TNC FWS	TBD					
3	1.2.3.1.	Movement of other mammals or carcasses to San Miguel Island should be avoided wherever possible.	SMIF	Ongoing	CDFW FWS NPS	TBD					
3	1.2.3.2.	Movement of other mammals or carcasses to Santa Rosa Island should be avoided wherever possible.	SRIF	Ongoing	CDFW FWS NPS	TBD					
3	1.2.3.3.	Movement of other mammals or carcasses to Santa Cruz Island should be avoided wherever possible.	SCZIF	Ongoing	CDFW FWS NPS TNC	TBD					
3	1.2.4.	The potential for pathogen introduction to Santa Catalina Island from the movement of wild and domestic mammals should be reduced to the extent practicable.	SCIF	Ongoing	CDFW CIC FWS	20	2	2	2	2	2
3	1.2.5.1.	Develop a management strategy for responding to new introductions of animals to San Miguel Island.	SMIF	Ongoing	CDFW FWS NPS	22.5	12.5	2.5	2.5	2.5	2.5
3	1.2.5.2.	Develop a management strategy for responding to new introductions of animals to Santa Rosa Island.	SRIF	Ongoing	CDFW FWS NPS	22.5	12.5	2.5	2.5	2.5	2.5

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	1.2.5.3.	Develop a management strategy for responding to new introductions of animals to Santa Cruz Island.	SCZIF	Ongoing	CDFW FWS NPS TNC	22.5	12.5	2.5	2.5	2.5	2.5
3	1.2.5.4.	Develop a management strategy for responding to new introductions of animals to Santa Catalina Island.	SCIF	Ongoing	CDFW CIC FWS	45	12.5	2.5	2.5	2.5	2.5
1	1.3.1.1.1.	Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on San Miguel Island foxes during a single vaccination event.	SMIF	One year	FWS NPS UNIV	1.25	1.25				
1	1.3.1.1.2.	Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on Santa Rosa Island foxes during a single vaccination event.	SRIF	One year	FWS NPS UNIV	1.25	1.25				
1	1.3.1.1.3.	Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on Santa Cruz Island foxes during a single vaccination event.	SCZIF	One year	FWS NPS TNC UNIV	1.25	1.25				
1	1.3.1.1.4.	Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on Santa Catalina Island foxes during a single vaccination event.	SCIF	One year	CIC FWS UNIV	1.25	1.25				

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	1.3.1.2.1.	Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in San Miguel Island foxes.	SMIF	Ongoing	FWS NPS UNIV	12	9.5	2.5			
1	1.3.1.2.2.	Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in Santa Rosa Island foxes.	SRIF	Ongoing	FWS NPS UNIV	12	9.5	2.5			
1	1.3.1.2.3.	Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in Santa Cruz Island foxes.	SCZIF	Ongoing	FWS NPS TNC UNIV	12	9.5	2.5			
1	1.3.1.2.4.	Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in Santa Catalina Island foxes.	SCIF	Ongoing	CIC FWS UNIV	12	9.5	2.5			
3	1.3.1.3.1.	Vaccines against canine parvovirus and adenovirus should to be tested on San Miguel Island foxes.	SMIF	Continual	FWS UNIV	2.5	2.5				
3	1.3.1.3.2.	Vaccines against canine parvovirus and adenovirus should to be tested on Santa Rosa Island foxes.	SRIF	Continual	FWS UNIV	2.5	2.5				
3	1.3.1.3.3.	Vaccines against canine parvovirus and adenovirus should to be tested on Santa Cruz Island foxes.	SCZIF	Continual	FWS UNIV	2.5	2.5				
3	1.3.1.3.4.	Vaccines against canine parvovirus and adenovirus should to be tested on Santa Catalina Island foxes.	SCIF	Continual	FWS UNIV	2.5	2.5				

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	1.3.2.1.	On San Miguel Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes.	SMIF	Ongoing	FWS NPS	170	34	34	34	34	34
1	1.3.2.2.	On Santa Rosa Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes.	SRIF	Ongoing	FWS NPS	170	34	34	34	34	34
1	1.3.2.3.	On Santa Cruz Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes.	SCZIF	Ongoing	FWS NPS TNC	170	34	34	34	34	34
1	1.3.2.4.	On Santa Catalina Island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes.	SCIF	Ongoing	CIC FWS	430	43	43	43	43	43
1	1.4.1.1.1.	Using radio-telemetry, monitor a sample of foxes on San Miguel Island to detect fox mortalities.	SMIF	Ongoing	NPS	220	44	44	44	44	44
1	1.4.1.1.2.	Using radio-telemetry, monitor a sample of foxes on Santa Rosa Island to detect fox mortalities.	SRIF	Ongoing	NPS	220	44	44	44	44	44
1	1.4.1.1.3.	Using radio-telemetry, monitor a sample of foxes on Santa Cruz Island to detect fox mortalities.	SCZIF	Ongoing	NPS TNC	220	44	44	44	44	44
1	1.4.1.1.4.	Using radio-telemetry, monitor a sample of foxes on Santa Catalina Island to detect fox mortalities.	SCIF	Ongoing	CIC	620	62	62	62	62	62
1	1.4.1.2.1.	Any San Miguel Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy.	SMIF	Ongoing	NPS	31.25	6.25	6.25	6.25	6.25	6.25

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	1.4.1.2.2.	Any Santa Rosa Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy.	SRIF	Ongoing	NPS	31.25	6.25	6.25	6.25	6.25	6.25
1	1.4.1.2.3.	Any Santa Cruz Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy.	SCZIF	Ongoing	NPS TNC	31.25	6.25	6.25	6.25	6.25	6.25
1	1.4.1.2.4.	Any Santa Catalina Island foxes that are found dead should be collected and shipped or frozen immediately for necropsy.	SCIF	Ongoing	CIC	100	10	10	10	10	10
1	1.4.1.3.1.	Any San Miguel Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored.	SMIF	Ongoing	NPS	1.5	.5	.25	.25	.25	.25
1	1.4.1.3.2.	Any Santa Rosa Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored.	SRIF	Ongoing	NPS	1.5	.5	.25	.25	.25	.25
1	1.4.1.3.3.	Any Santa Cruz Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored.	SCZIF	Ongoing	NPS TNC	1.5	.5	.25	.25	.25	.25
1	1.4.1.3.4.	Any Santa Catalina Island foxes appearing ill or acting in an abnormal manner should be reported immediately, quarantined and closely monitored.	SCIF	Ongoing	CIC	3	.5	.25	.25	.25	.25

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	1.4.1.4.1.	Other carnivores on San Miguel Island found sick or dead from causes other than trauma should be reported and closely monitored (if alive) or collected for necropsy (if dead).	SMIF	Continual	NPS	2.5	.25	.25	.25	.25	.25
3	1.4.1.4.2.	Other carnivores on Santa Rosa Island found sick or dead from causes other than trauma should be reported and closely monitored (if alive) or collected for necropsy (if dead).	SRIF	Continual	NPS	2.5	.25	.25	.25	.25	.25
3	1.4.1.4.3.	Other carnivores on Santa Cruz Island found sick or dead from causes other than trauma should be reported and closely monitored (if alive) or collected for necropsy (if dead).	SCZIF	Continual	NPS TNC	2.5	.25	.25	.25	.25	.25
3	1.4.1.4.4.	Other carnivores on Santa Catalina Island found sick or dead from causes other than trauma should be reported and closely monitored (if alive) or collected for necropsy (if dead).	SCIF	Continual	CIC	5	.25	.25	.25	.25	.25
2	1.4.2.1.	Annually collect blood samples from a proportion of island foxes on San Miguel Island to evaluate ongoing disease risks to island fox population.	SMIF	Ongoing	NPS	100	20	20	20	20	20
2	1.4.2.2.	Annually collect blood samples from a proportion of island foxes on Santa Rosa Island to evaluate ongoing disease risks to island fox population.	SRIF	Ongoing	NPS	100	20	20	20	20	20

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
2	1.4.2.3.	Annually collect blood samples from a proportion of Santa Cruz Island foxes to evaluate ongoing disease risks to island fox population.	SCZIF	Ongoing	NPS TNC	100	20	20	20	20	20
2	1.4.2.4.	Annually collect blood samples from a proportion of island foxes on Santa Catalina Island to evaluate ongoing disease risks to the island fox population.	SCIF	Ongoing	CIC	400	40	40	40	40	40
1	1.4.3.1.1.	San Miguel Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes.	SMIF	Ongoing	NPS						
1	1.4.3.1.2.	Santa Rosa Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes.	SRIF	Ongoing	NPS						
1	1.4.3.1.3.	Santa Cruz Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes.	SCZIF	Ongoing	NPS TNC						
1	1.4.3.1.4.	Santa Catalina Island managers should develop an emergency response strategy for dealing with disease incidents relevant to island foxes.	SCIF	Ongoing	CIC						
1	1.4.3.2.1.	A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for San Miguel Island foxes.	SMIF	Ongoing	NPS	TBD					

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	1.4.3.2.2.	A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for Santa Rosa Island foxes.	SRIF	Ongoing	NPS	TBD					
1	1.4.3.2.3.	A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for Santa Cruz Island foxes.	SCZIF	Ongoing	NPS TNC	TBD					
1	1.4.3.2.4.	A single case of rabies, confirmed by pathology or virus isolation, should trigger management response for Santa Catalina Island foxes.	SCIF	Ongoing	CIC	TBD					
1	1.4.3.3.1.	A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for San Miguel Island foxes.	SMIF	Ongoing	NPS	TBD					
1	1.4.3.3.2.	A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for Santa Rosa Island foxes.	SRIF	Ongoing	NPS	TBD					
1	1.4.3.3.3.	A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for Santa Cruz Island foxes.	SCZIF	Ongoing	NPS TNC	TBD					
1	1.4.3.3.4.	A single case of canine distemper, confirmed by pathology or virus isolation, should trigger management response for Santa Catalina Island foxes.	SCIF	Ongoing	CIC	TBD					

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
2	1.4.3.4.1.	A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead San Miguel Island foxes.	SMIF	Ongoing	NPS	TBD					
2	1.4.3.4.2.	A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead Santa Rosa Island foxes.	SRIF	Ongoing	NPS	TBD					
2	1.4.3.4.3.	A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead Santa Cruz Island foxes.	SCZIF	Ongoing	NPS TNC	TBD					
2	1.4.3.4.4.	A single case of disease caused by parvovirus or adenovirus, confirmed by pathology or virus isolation, should prompt more intensive monitoring for sick or dead Santa Catalina Island foxes.	SCIF	Ongoing	CIC	TBD					
2	1.5.1.	Complete on-going investigations of the demographic consequences and etiology of the ear tumors prevalent in Santa Catalina Island foxes should be completed to determine whether this disease poses a significant threat to this fox subspecies.	SCIF	Ongoing	CIC FWS UNIV	90	30	10	3	1	1

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
2	1.5.2.1.	On San Miguel Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted.	SMIF	Continual	FWS NPS UNIV	18.75	12.5	2.5	1.25	1.25	1.25
2	1.5.2.2.	On Santa Rosa Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted.	SRIF	Continual	FWS NPS UNIV	18.75	12.5	2.5	1.25	1.25	1.25
2	1.5.2.3.	On Santa Cruz Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted.	SCZIF	Continual	FWS NPS TNC UNIV	18.75	12.5	2.5	1.25	1.25	1.25
2	1.5.2.4.	On Santa Catalina Island, expand research on the role of co-pathogens and viral strain variation to provide better insights into the circumstances of a disease outbreak under which management interventions are, and are not, warranted.	SCIF	Continual	CIC FWS UNIV	37.5	12.5	2.5	1.25	1.25	1.25

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefiting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
2	1.5.3.1.	Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the San Miguel Island fox population.	SMIF	Continual	FWS NPS UNIV	37.5	12.5	6.25	6.25	6.25	6.25
2	1.5.3.2.	Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the Santa Rosa Island fox population.	SRIF	Continual	FWS NPS UNIV	37.5	12.5	6.25	6.25	6.25	6.25
2	1.5.3.3.	Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the Santa Cruz Island fox population.	SCZIF	Continual	FWS NPS TNC UNIV	37.5	12.5	6.25	6.25	6.25	6.25
2	1.5.3.4.	Further research should be conducted as appropriate on other infectious and noninfectious diseases that appear likely, on the basis of pathological and demographic analyses, to threaten the Santa Catalina Island fox population.	SCIF	Continual	CIC FWS UNIV	75	12.5	6.25	6.25	6.25	6.25
3	2.1.1.	Continue captive management of San Miguel Island foxes as necessary to provide individuals for release.	SMIF	Ongoing	NPS						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	2.1.2.	Continue captive management of Santa Rosa Island foxes as necessary to provide individuals for release.	SRIF	Ongoing	NPS						
3	2.1.3.	Continue captive management of Santa Cruz Island foxes as necessary to provide individuals for release.	SCZIF	Ongoing	NPS						
3	2.1.4.	Continue captive management of Santa Catalina Island foxes as necessary to provide individuals for release.	SCIF	Ongoing	CIC						
3	2.2.1.	Assuming golden eagle predation and disease risks remain at low levels, continue annual release of San Miguel Island foxes from the captive facility until such releases are no longer necessary to augment the wild population.	SMIF	Ongoing	NPS						
3	2.2.2.	Assuming golden eagle predation and disease risks remain at low levels, continue annual release of Santa Rosa Island foxes from the captive facility until such releases are no longer necessary to augment the wild population.	SRIF	Ongoing	NPS						
3	2.2.3.	Assuming golden eagle predation and disease risks remain at low levels, continue annual release of Santa Cruz Island foxes from the captive facility until such releases are no longer necessary to augment the wild population.	SCZIF	Ongoing	NPS						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	2.2.4.	Assuming golden eagle predation and disease risks remain at low levels, continue annual release of Santa Catalina Island foxes from the captive facility until such releases are no longer necessary to augment the wild population.	SCIF	Ongoing	CIC						
3	2.3.1.	Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which San Miguel Island fox individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing.	SMIF	Ongoing	AZA NPS						
3	2.3.2.	Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which Santa Rosa Island fox individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing.	SRIF	Ongoing	AZA NPS						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefiting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	2.3.3.	Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which Santa Cruz Island fox individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing.	SCZIF	Ongoing	AZA NPS						
3	2.3.4.	Use genetic, demographic, and appropriate behavioral and physiological characteristics, together with established PVA models, to determine which Santa Catalina Island fox individuals to release and which to retain annually, such that an appropriate level of genetic diversity of the remaining captive population is retained while captive breeding is ongoing.	SCIF	Ongoing	AZA						
2	2.4.1.	Continue to monitor released San Miguel Island foxes and use this information to modify release strategies (e.g., release locations and timing).	SMIF	Ongoing	NPS						
2	2.4.2.	Continue to monitor released Santa Rosa Island foxes and use this information to modify release strategies (e.g., release locations and timing).	SRIF	Ongoing	NPS						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
2	2.4.3.	Continue to monitor released Santa Cruz Island foxes and use this information to modify release strategies (e.g., release locations and timing).	SCZIF	Ongoing	NPS						
2	2.4.4.	Continue to monitor released Santa Catalina Island foxes and use this information to modify release strategies (e.g., release locations and timing).	SCIF	Ongoing	CIC						
3	2.5.1.	Once captive breeding is no longer necessary on San Miguel Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat.	SMIF	Ongoing	NPS	6.25	1.25	1.25	1.25	1.25	1.25
3	2.5.2.	Once captive breeding is no longer necessary on Santa Rosa Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat.	SRIF	Ongoing	NPS	6.25	1.25	1.25	1.25	1.25	1.25

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	2.5.3.	Once captive breeding is no longer necessary on Santa Cruz Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat.	SCZIF	Ongoing	NPS	6.25	1.25	1.25	1.25	1.25	1.25
3	2.5.4.	Once captive breeding is no longer necessary on Santa Catalina Island, maintain captive facilities such that a predetermined number of island foxes could be recaptured and maintained in captivity in the event of a new catastrophic threat.	SCIF	Ongoing	CIC	12.5	1.25	1.25	1.25	1.25	1.25
3	2.6.1.	While San Miguel Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes.	SMIF	Ongoing	NPS						
3	2.6.2.	While Santa Rosa Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes.	SRIF	Ongoing	NPS						

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	2.6.3.	While Santa Cruz Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes.	SCZIF	Ongoing	NPS						
3	2.6.4.	While Santa Catalina Island on-island captive populations still exist, continue to identify and implement improved husbandry practices to ensure the health of captive island foxes, improve reproductive success, and enhance the success of released foxes.	SCIF	Ongoing	CIC						
2	2.7.1.	Continue to maintain a San Miguel Island fox studbook and continue to use the studbook to aid decisions on fox pairing and release.	SMIF	Ongoing	AZA NPS	8.5	1.7	1.7	1.7	1.7	1.7
2	2.7.2.	Continue to maintain a Santa Rosa Island fox studbook and continue to use the studbook to aid decisions on fox pairing and release.	SRIF	Ongoing	AZA NPS	8.5	1.7	1.7	1.7	1.7	1.7
2	2.7.3.	Continue to maintain a Santa Cruz Island fox studbook and continue to use the studbook to aid decisions on fox pairing and release.	SCZIF	Ongoing	AZA NPS	8.5	1.7	1.7	1.77	1.7	1.7

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
1	3.1.1.	Develop and implement a monitoring strategy for San Miguel Island fox subspecies to detect population declines and determine population trends.	SMIF	Ongoing	CDFW FWS NPS	385	77	77	77	77	77
1	3.1.2.	Develop and implement a monitoring strategy for Santa Rosa Island fox subspecies to detect population declines and determine population trends.	SRIF	Ongoing	CDFW FWS NPS	385	77	77	77	77	77
1	3.1.3.	Develop and implement a monitoring strategy for Santa Cruz Island fox subspecies to detect population declines and determine population trends.	SCZIF	Ongoing	CDFW FWS NPS TNC	385	77	77	77	77	77
1	3.1.4.	Develop and implement a monitoring strategy for Santa Catalina Island fox subspecies to detect population declines and determine population trends.	SCIF	Ongoing	CDFW CIC FWS	770	77	77	77	77	77
3	3.2.1.	Ensure San Miguel Island fox population information is comparable across islands to the greatest extent possible.	SMIF	Ongoing	CDFW FWS NPS	2.5	.5	.5	.5	.5	.5
3	3.2.2.	Ensure Santa Rosa Island fox population information is comparable across islands to the greatest extent possible.	SRIF	Ongoing	CDFW FWS NPS	2.5	.5	.5	.5	.5	.5
3	3.2.3.	Ensure Santa Cruz Island fox population information is comparable across islands to the greatest extent possible.	SCZIF	Ongoing	CDFW FWS NPS TNC	2.5	.5	.5	.5	.5	.5

Recovery Plan for Four Subspecies of Island Fox

Action Priority	Action Number	Action Description	Species Benefitting	Action Status or Duration	Responsible Parties	Total Cost Estimate (in \$1,000 units)					
						Total	Year 1	Year 2	Year 3	Year 4	Year 5
3	3.2.4.	Ensure Santa Catalina Island fox population information is comparable across islands to the greatest extent possible.	SCIF	Ongoing	CDFW CIC FWS	5	.5	.5	.5	.5	.5

VI. Long-term Conservation Strategy

The long-term conservation strategy identifies actions that would further the conservation of the island fox. At this time, these activities are not essential for preventing extinction and are not required for downlisting or delisting a particular island fox subspecies; however, these activities could substantially enhance the long-term conservation of the species and may also increase our scientific understanding of the island fox. In the event that an island fox subspecies is recovered and delisted, completion of these actions could provide conservation benefits that could prevent future decline of the species.

We have identified the following long-term conservation actions:

- Establish a mainland captive island fox population on which to conduct research to better understand fox behavior, ecology and reproduction, and disease and vaccine efficacy.
- Increase public awareness to reduce potential threats from anthropogenic activities.
- Assess the demographic impact of other threats such as mortality from vehicle strikes and competition with feral cats.
- Restore island habitat.
- Establish conservation agreements.

1. Establish a mainland captive island fox population on which to conduct research to better understand fox behavior, ecology and reproduction, and disease and vaccine efficacy.

The establishment of a mainland captive island fox population could contribute to island fox conservation through improved opportunities for research and increased opportunities for educating and affecting public attitudes towards the island fox which could result in greater support for island fox conservation programs. A mainland population might also provide a source population for re-colonization should the subspecies become extinct. Note: unless a subspecies is extinct in the wild, any island fox housed for any reason on the mainland will not be released back to their respective island and will remain on the mainland.

A mainland captive island fox population can serve as an accessible source of individuals for research. There are still many unanswered questions concerning the best husbandry and management methods for successful captive breeding. Mainland facilities are more accessible to veterinary and husbandry experts and more efficient in terms of costs and logistics. The ability to conduct research trials on the islands is limited. Furthermore, such trials are currently not possible on the islands because existing captive breeding facilities have closed. The

Recovery Plan for Four Subspecies of Island Fox

source for a mainland captive population can come from any of the existing island fox subspecies, including non-listed subspecies, although TAR 3.6 “Assessment of the potential benefits and costs of long-term captive populations on the mainland and/or islands” suggests that the Santa Cruz Island fox population would be the best choice, because this subspecies has the most genetic diversity and the island population is recovering rapidly. Note: Unless otherwise approved by the FWS and CDFW, only injured, un-releasable foxes or abandoned pups that are incapable of surviving in the wild will be considered for transfer to zoos.

Given that space and resources are limited to establish redundant populations for each of the four endangered subspecies of island fox it is unrealistic to expect to have a redundant population of each subspecies. In the event of a catastrophic loss, two alternatives exist to repopulate an island: 1) use individuals from another existing wild island fox subspecies; or 2) use individuals from an established mainland captive population where one, or at most two, subspecies would be maintained. As noted above, the Santa Cruz Island fox population has been identified as the best choice to have individuals represented on the mainland for long-term captive populations because this subspecies has the most genetic diversity and the island population is recovering rapidly. Therefore, every effort should be made to provide opportunities to establish this mainland program, including healthy animals that might not otherwise meet the criteria of injured, un-releasable foxes or abandoned pups that are incapable of surviving in the wild.

Below is a list of activities that would benefit island fox research:

1.1 Develop a captive mainland island fox population for research and educational outreach purposes.

- Develop a long-range strategy for establishing a mainland captive island fox population in accordance with the FWS Captive Propagation Policy and CDFW Policy.

This strategy should strive to maximize the genetic and demographic viability of the mainland populations while avoiding or minimizing any detrimental impacts to wild populations. See Appendix 7 for steps necessary in establishing a mainland captive population.

- Identify and prioritize research questions that could be addressed using the captive mainland population.

This might be achieved by forming a standing advisory committee to review proposals, prioritize projects, and help identify funding sources. Selected research questions include: 1) determine the best management practices for husbandry to maximize reproduction and ensure animal welfare (e.g., mate selection, housing requirements); 2) biomedical research on

captive populations to help eliminate and/or control disease threats to the wild and captive island fox populations; and 3) development of management and husbandry techniques to maximize fox survival post-release (see Appendix 8 for details).

2. Establish, expand and continue island fox education and outreach programs.

The main objectives of the education and outreach programs include: reduce threats that are under the control of island managers, residents, visitors, and regulatory agencies; increase public support for existing and future programs dealing with island fox population recovery, threat abatement, habitat improvement and sustainable use of habitats; and development of long-term funding support, including fund-raising activities, for island fox recovery efforts.

Below is a list of such activities that would aid in island fox education and outreach:

2.1 Develop and establish on-island education programs.

- Provide island fox information to residents, staff, and visitors to the Channel Islands.
- On each island with foxes, develop self-guided kiosks, exhibits, and/or programs to provide current information about island foxes and recovery efforts.

2.2 Develop and establish mainland education programs.

- Identify educational and outreach opportunities that could be addressed using the captive mainland population.
- Develop strong collaborations with existing organizations (e.g., Friends of the Island Fox, Inc.) working on mainland education programs.
- All zoos that house island fox populations, particularly those in southern California, should include an education program with an assessment requirement of the education programs effectiveness.
- Mainland island fox exhibits should provide accurate and timely information on the status of and threats to island foxes.
- Develop island fox presentations, traveling exhibits and publications to be presented or deployed in mainland schools, symposia, meetings and other venues.

Recovery Plan for Four Subspecies of Island Fox

- Develop school curriculum materials on island foxes, consistent with California education standards that can be used in mainland classrooms prior to student field trips to an island.

2.3 Develop cost effective methods for enhancing public awareness and support for island fox recovery.

- Utilize the media to enhance public awareness and support for island fox recovery programs.
- Develop an appropriate set of professional evaluation tools (Measures of Success) to help managers and agencies evaluate the effectiveness of the general and island-wide education and awareness programs.
- Develop an effective set of communication venues for island fox researchers and land managers.

2.4 Continue and expand, as appropriate, the annual island fox conference, and develop a web-based literature depository and/or a regular newsletter or list-server to enhance communication.

- Establish a web-based island fox literature library where educators and researchers can access information about the island fox.
- Develop and publish an annual report on island fox recovery and conservation efforts.

3. Assessing the demographic impact of other threats such as mortality from vehicle strikes, competition with feral cats, and emerging disease issues (e.g., ear cancer).

The threat from anthropogenic sources such as vehicles, competition with feral cats, emerging diseases and other mortality factors may impact island fox population dynamics. Research into the effects of these potential threats will help the design of long-term conservation strategies.

4. Restoring island habitat.

Ultimately the long-term survival and viability of the island fox may depend on maintaining and restoring some of the composition, structure, and function of native ecosystems on the islands that have been altered in the past 150 years. These actions include promoting ecological conditions that dissuade golden eagle use of the Channel Islands.

Preserving and restoring native ecosystem structure and function while preserving significant cultural resources and providing for recreational opportunities are

Recovery Plan for Four Subspecies of Island Fox

explicit management goals of the NPS; TNC and CIC share similar goals. While island foxes were no doubt components of formerly intact ecosystems, there is no assurance that those intact ecosystems were optimal in terms of sustaining the largest populations of foxes. For example, the conversion of some shrublands to grasslands during the ranching period may have provided for increased populations of island foxes. On the other hand, in the face of aerial predation, shrublands may confer differential advantage to foxes. Consequently, ecosystem restoration activities will need to respond adaptively to the response of island fox populations.

The goal of ecosystem restoration is to maintain and restore native ecosystem composition, structure, and function in a manner that does not compromise island fox recovery. Ultimately, activities will restore native ecosystem integrity in a manner that enhances island fox recovery and long-term conservation, while protecting other listed and sensitive species. Because each of the Channel Islands differs significantly in their native ecosystem structure and composition, maintenance and restoration should be tailored to each island individually.

Below is a list of activities that would aid in ecosystem recovery:

- 4.1 Identify non-native plants and animals that may compromise island fox viability and evaluate their impact on fox populations. Removal or control should be conducted if impacts are significant or potentially significant and the means for practicable removal or control exists.
- 4.2 Minimize the likelihood of new non-native species introductions through the use of education, regulation, sanitation, and best management practices.
- 4.3 Reintroduce or enhance native ecosystem elements and processes that have been lost or compromised as a result of anthropogenic activities.
- 4.4 Prevent excessive human-caused impacts to island ecosystems' native structure and function to the extent practicable.
- 4.5 Minimize, to the extent feasible, mechanical, chemical, or acoustic impacts to island foxes and den sites during restoration activity, especially during the breeding and pup-rearing seasons.
- 4.6 Protect natural water supplies in island fox habitat from damage, and avoid eliminating island fox water sources.
- 4.7 Monitor island fox food resources during restoration efforts, including native animal prey populations and plant resources.

Recovery Plan for Four Subspecies of Island Fox

- If these food resources change adversely during restoration activity, manage adaptively to provide for adequate food resources during the restoration period.
- 4.8 Monitor and adaptively manage distribution of habitat types on each island during restoration activities to assure sufficient ecosystem services for island foxes, such as hunting habitat, resting habitat, and protective cover (against predation).
- Although island foxes have historically occurred in nearly all vegetation types, maintaining the native array of these types provides a buffer against unanticipated ecological catastrophe.
- 4.9 Where non-native species may represent a supportive habitat function (food, cover), plan ecosystem restoration actions to assure alternate prey or other resources provided by the non-native species are available and sufficient during the restoration period.
- 4.10 As naturally-ignited landscape fire on the Channel Islands is rare and most ecosystem elements, including island foxes, have not been selected for fire resilience, minimize the likelihood of anthropogenic fire.

5. Establishment of Conservation Agreements.

Even with successful mitigation of current threats and the recovery of island fox subspecies to viable population levels, the intrinsically small population sizes of the subspecies and their insular vulnerabilities subject the different subspecies to the continued threat of catastrophic decline from any number of causes.

To reduce the potential for future catastrophic population declines and the consequent need to relist the San Miguel Island fox, the Santa Rosa Island fox, the Santa Cruz Island fox, or the Santa Catalina Island fox post-recovery:

A Cooperative Management Agreement should be developed between the land manager(s) and the FWS to address long-term conservation needs. The agreement should be designed to respond effectively to any future significant population decline and include:

- The land manager's strategy and commitment to continue monitoring island fox subspecies such that any substantial population decline is detected in a timely manner; and
- The land manager's strategy to address the long-term conservation of island foxes at the time of proposed delisting.

VII. Literature Cited

- Aguilar, A., G. Roemer, S. Debenham, M. Binns, D. Garcelon, and R.K. Wayne. 2004. High MHC diversity maintained by balancing selection in an otherwise genetically monomorphic mammal. *Proceedings of the National Academy of Sciences* 101(10):3490-3494.
- Anon. 1931. Catalina cats and quail. *California Fish and Game* 17(4):450-51.
- Appel, M., B.E. Sheffy, D.H. Percy, and J.M. Gaskin. 1974. Canine distemper virus in domesticated cats and pigs. *American Journal of Veterinary Research* 35:803-806.
- Backlin, A.R., S.L. Compton, Z.B. Kahancza, and R.N. Fisher. 2005. Baseline biodiversity survey for Santa Catalina Island: herpetofauna and ants with remarks on small mammals and others. Final Report. U.S. Geological Survey Western Ecological Research Center, San Diego, California.
- Baird, S.F. 1857. General report upon the zoology of the several Pacific Railroad Routes. I. Mammals. In: *Reports of Explorations and Surveys to Ascertain the Most Practicable and Economical Route for a Railroad from the Mississippi River to the Pacific Ocean*, Vol. 8. Washington, D.C.
- Bakker, V. J. and D. F. Doak. 2009. Population viability management: Ecological standards to guide adaptive management for rare species. *Frontiers in Ecology and the Environment* 7(3):158-165.
- Bakker, V.J., D.F. Doak, G.W. Roemer, D.K. Garcelon, T.J. Coonan, S.A. Morrison, C. Lynch, K. Ralls, and M.R. Shaw. 2009. Incorporating ecological drivers and uncertainty into a demographic population viability analysis for the Island Fox (*Urocyon littoralis*). *Ecological Monographs* 79(1):77-108.
- Biteman, D. S., L. S. Brown, A. S. Bridges, D. K. Garcelon. 2010. Predator Research and Management in Support of the Recovery of the San Clemente Loggerhead Shrike on San Clemente Island. Final Report - 2009. U. S. Navy, Naval Facilities Engineering Command Southwest, San Diego, California, USA. 72 pp.
- Biteman, D. S., T. W. Higgins, A. S. Bridges, D. K. Garcelon. 2012. Predator Research and Ecosystem Management in Support of the Recovery of the San Clemente Loggerhead Shrike on San Clemente Island. Final Annual Report - 2011. Natural Resources Office, Naval Base Coronado, San Diego, California. 75 pp.
- Biteman, D. S., A. S. Bridges, D. K. Garcelon. 2013. Predator Research and Ecosystem Management in Support of the Recovery of the San Clemente

Recovery Plan for Four Subspecies of Island Fox

- Loggerhead Shrike on San Clemente Island. Final Annual Report - 2012. Natural Resources Office, Naval Base Coronado, San Diego, California. 84 pp.
- Boser, C. 2011. Annual Report for the Island Fox Recovery Actions Conducted under Permit TE163610-0. The Nature Conservancy, California. 20 pp.
- Bridges, A.S. S.N.D. King, D.K. Garcelon. 2014. San Clemente Island Fox Monitoring and Demography. Final report prepared for Naval Base Coronado, Public Works Office, Code N45, Building 3, 2nd Floor. PO Box 357088 San Diego, California 92135-7088. 29 pp.
- Brito, A.F., L.C. Souza, A.V. da Silva, and H. Langoni. 2002. Epidemiological and serological aspects in canine toxoplasmosis in animals with nervous symptoms. *Memórias do Instituto Oswaldo Cruz* 97:31-35.
- Brumbaugh, R.W. 1980. Recent geomorphic and vegetal dynamics on Santa Cruz Island, California. Pages 139-158 in: Power, D. M. (ed.), *The California Islands: Proceedings of a Multidisciplinary Symposium*. Santa Barbara Museum of Natural History, Santa Barbara, California.
- California Department of Fish and Game. 1987. Five-year status report on the island fox. Unpublished report, California Department of Fish and Game, Sacramento, California.
- Cayan, D., M. Dettinger, I. Stewart, and N. Knowles. 2005. Recent changes towards earlier springs: early signs of climate warming in western North America? U.S. Geological Survey, Scripps Institution of Oceanography, La Jolla, California.
- Clark, R.A., W.L. Halvorson, A.A. Sawdo, and K.C. Danielsen. 1990. Plant communities of Santa Rosa Island, Channel Islands National Park. Tech. Report No. 42. Cooperative National Park Resources Studies Unit, University of California, Davis.
- Clifford, D.L. 2006. Health threats affecting Island Fox (*Urocyon littoralis*) population recovery on the California Channel Islands. Dissertation, University of California, Davis. 198 pp.
- Clifford, D.L., J.A.K. Mazet, E.J. Dubovi, D.K. Garcelon, T.J. Coonan, P.J. Conrad, and L. Munson. 2006. Pathogen exposure in endangered island fox (*Urocyon littoralis*) populations: implications for conservation management. *Biological Conservation* 131:230-243.
- Coblentz, B. 1980. Effects of feral goats on the Santa Catalina Island ecosystem. Pages 167-172 in Power, D. M., (ed.), *The California Islands: Proceedings*

Recovery Plan for Four Subspecies of Island Fox

of a Multidisciplinary Symposium. Santa Barbara Museum of Natural History, Santa Barbara, California.

- Collins, P.W. 1980. Food habits of the island fox (*Urocyon littoralis littoralis*) on San Miguel Island, California. In Proceedings of the 2nd Conference on Scientific Research in the National Parks. Volume 12: Terrestrial Biology and Zoology. National Park Service, Washington D.C. NTIS.P881-100133.
- Collins, P. 1982. Origin and differentiation of the island fox: a study of evolution in insular populations. M. A. thesis. University of California, Santa Barbara. 303 pp.
- Collins, P.W. 1991a. Interaction between island foxes (*Urocyon littoralis*) and Indians on islands off the coast of southern California: I. Morphologic and archaeological evidence of human assisted dispersal. Journal of Ethnobiology 11(1):51-81.
- Collins, P.W. 1991b. Interaction between island foxes (*Urocyon littoralis*) and Native Americans on islands off the coast of southern California: II. Ethnographic, archaeological, and historical evidence of human assisted dispersal. Journal of Ethnobiology 11(2):205-229.
- Collins, P.W. 1993. Taxonomic and biogeographic relationships of the island fox (*Urocyon littoralis*) and gray fox (*U. cinereoargenteus*) from Western North America. Pages 351-390 in Hochberg, F. G., (ed.), Third California Islands Symposium: Recent Advances in Research on the California Islands. Santa Barbara Museum of Natural History, Santa Barbara, California. 661 pp.
- Collins, P.W. and B.C. Latta. 2006. Nesting season diet of golden eagles on Santa Cruz and Santa Rosa Islands, Santa Barbara County, California. Santa Barbara Museum of Natural History Technical Reports – No. 3. 40 pp.
- Collins, P.W. and B.C. Latta. 2009. Food habits of nesting golden eagles (*Aquila chrysaetos*) on Santa Cruz and Santa Rosa Islands, California. Pages 255-268 in: Damiani, C.C. and D.K. Garcelon (eds.). Proceedings of the Seventh California Islands Symposium. Institute for Wildlife Studies, Arcata, California.
- Collins, P.W. and L.L. Laughrin. 1979. Vertebrate zoology: the island fox on San Miguel Island. Pages 12.1-12.52 in: Power, D. M., (ed.), Natural Resources Study of the Channel Islands National Monument, California. Prepared for the National Park Service, Denver Service Center by the Santa Barbara Museum of Natural History, Santa Barbara, California.
- Collins, P.W. and T.D. Martin. 1985. A review of the population status of the Santa Catalina Island shrew, (*Sorex ornatus willetti*). U.S.D.I., Fish and

Recovery Plan for Four Subspecies of Island Fox

Wildlife Serv., Endangered Species Office, Sacramento, California, Final Report., 10181_5693 (BW) '85, SE_0018_85, 92 pp.

- Collins, P.W., D.A. Guthrie, T.C. Rick, and J.M. Erlandson. 2005. Analysis of prey remains excavated from a historic bald eagle nest site on San Miguel Island, California. Pages 103-120 in D.K. Garcelon and C.A. Schwemm, eds., Proceedings of the sixth California Islands Symposium, Ventura, CA, December 2003. National Park Service Technical Publication CHIS-05-01, Institute for Wildlife Studies, Arcata, California
- Coonan, T.J. 2003. Recovery strategy for island foxes (*Urocyon littoralis*) on the northern Channel Islands. National Park Service, Ventura California. 81 pp.
- Coonan, T.J. 2010. Summary Report: Twelfth annual meeting of the Island Fox Working Group, June 15-17, 2010. Unpublished Report on file at headquarters, Channel Islands National Park, Ventura, California. 52 pp.
- Coonan, T.J. 2011. 2010 annual report for island fox recovery actions conducted under Permit TE 0862267-0 by the National Park Service, Channel Islands National Park. Submitted to Ventura Fish and Wildlife Office, April 1, 2011. 33pp.
- Coonan, T.J. 2013. 2012 annual report for island fox recovery actions conducted under Permit TE 0862267-0 by the National Park Service, Channel Islands National Park. Submitted to Ventura Fish and Wildlife Office, April 22, 2013. 34pp
- Coonan, T.J. 2014. 2013 annual report for island fox recovery actions conducted under Permit TE-108507-2 by the National Park Service, Channel Islands National Park. Submitted to Ventura Fish and Wildlife Office, January 31, 2014. 54pp.
- Coonan, T.J. and A. Guglielmino. 2012. Island fox recovery program, 2011 annual report. Natural Resources Report NPS/MEDN/NRR – 2012/483. National Park Service, Fort Collins, Colorado. 42 pp.
- Coonan, T.J. and K.A. Rutz. 2001. Island fox captive breeding program, 1999-2000 annual report. Technical Report 01-01. National Park Service. 38 pp.
- Coonan, T.J. and K.A. Rutz. 2002. Island fox captive breeding program, 2001 annual report. Technical Report 02-01. National Park Service. 55 pp.
- Coonan, T.J. and K.A. Rutz. 2003. Island fox captive breeding program, 2002 annual report. Technical Report 03-01. National Park Service. 52 pp.

Recovery Plan for Four Subspecies of Island Fox

- Coonan, T.J. and C.A. Schwemm. 2009. Factors contributing to success of island fox reintroductions on San Miguel and Santa Rosa Islands, California. Pp. 363-376 in, Damiani, C.C and D.K. Garcelon, eds., Proceedings of the Seventh California Islands Symposium, Oxnard, California, February 5-8, 2008. Institute for Wildlife Studies, Arcata, California.
- Coonan, T.J. and C.A. Schwemm. In Review. Reintroduction of island foxes into unoccupied habitat on San Miguel and Santa Rosa Islands, California. Biological Conservation.
- Coonan, T.J., G. Austin, and C.A. Schwemm. 1998. Status and trend of island fox, San Miguel Island, Channel Islands National Park. Technical Report 98-01. National Park Service. 27 pp.
- Coonan, T.J., C.A.Schwemm, and D.K. Garcelon. 2010. Decline and recovery of the island fox: a case study for population recovery. Cambridge University Press, Cambridge, United Kingdom. 212 pp.
- Coonan, T.J., C.A. Schwemm, G.W. Roemer, and G. Austin. 2000. Population decline of island foxes (*Urocyon littoralis littoralis*) on San Miguel Island. Pages 289-297 in Browne, D. K., K. L. Mitchell, and H. W. Chaney, (eds.), Proceedings of the 5th California Islands Symposium. U.S. Department of the Interior, Minerals Management Service, Pacific OCS Region.
- Coonan, T.J., C.A. Schwemm, G.W. Roemer, D.K. Garcelon, and L. Munson. 2005c. Decline of an island fox subspecies to near extinction. Southwestern Naturalist 50(1):32-41.
- Coonan, T. J., V. Bakker, B. Hudgens, C. L. Boser, D. K. Garcelon, and S. A. Morrison. In Press. On the fast track to recovery: island foxes on the northern Channel Islands. Western North American Naturalist.
- Coonan, T.J., K. McCurdy, K.A. Rutz, M. Dennis, S. Provinsky, and S. Coppelli. 2005b. Island fox recovery program 2004 annual report. Technical Report 05-07. National Park Service. 63 pp.
- Coonan, T. J., K.A. Rutz, D. K. Garcelon, B. C. Latta, M. M. Gray, and E. T. Aschehoug. 2005a. Progress in island fox recovery efforts on the northern Channel Islands. Pages 263-273 in: Garcelon, D.K., and C.A. Schwemm, (eds.), Proceedings of the Sixth California Islands Symposium. National Park Service Technical Publication CHIS-05-01, Institute for Wildlife Studies, Arcata, California.
- Coonan, T.J., K.A. Rutz, K. McCurdy, D.K. Garcelon, B.C. Latta, and L. Munson. 2004. Island fox recovery program, 2003 annual report. Technical Report 04-02. National Park Service. 65 pp.

Recovery Plan for Four Subspecies of Island Fox

- Crooks, K.R. and D. Van Vuren. 1995. Resource utilization by two insular endemic mammalian carnivores, the island fox and island spotted skunk. *Oecologia* 104:301-307.
- Crooks, K.R. and D. Van Vuren. 1996. Spatial organization of the island fox (*Urocyon littoralis*) on Santa Cruz Island, California. *Journal of Mammalogy* 77(3):801-806.
- Crowell, H.E. 2001. Food habits and prey availability of the threatened San Miguel Island fox, *Urocyon littoralis littoralis*, during a decline in population, 1993–1998. Thesis, California Polytechnic State University, San Luis Obispo, California
- Davidson, W.R., V.F. Nettles, L.E. Hayes, E.W. Howerth, and C.E. Couvillion. 1992. Diseases diagnosed in gray foxes (*Urocyon cinereoargenteus*) from the southeastern United States. *Journal of Wildlife Diseases* 28:28–33.
- Dennis, M., K. Randall, G. Schmidt, and D. Garcelon. 2001. Island fox (*Urocyon littoralis santacruzae*) distribution, abundance and survival on Santa Cruz Island, California. Progress report: May through October 2001. Institute for Wildlife Studies, Arcata, California. 32 pp.
- Dennis, M., K. Randall, G. Schmidt, and D. Garcelon. 2002. Island fox (*Urocyon littoralis santacruzae*) distribution, abundance and survival on Santa Cruz Island, California. Progress report: for work conducted 08 Feb 2002 through 31 Mar 2002. Institute for Wildlife Studies, Arcata, California. 24 pp.
- Department of the Navy. 1963. Memorandum of agreement between the Department of the Navy and the Department of the Interior relating protection of natural values and historic and scientific objects on San Miguel and Prince Islands, California.
- Department of the Navy. 1985. Interagency agreement supplementing memorandum of agreement (dated 7 May 1963) and amended (20 October 1976) between the Departments of Interior and Navy regarding San Miguel Island.
- Department of the Navy. 1999. Environmental Assessment on Predator Management Activities, San Clemente Island, California.
- Dubey, J.P., J.L. Carpenter, M.J. Topper, and A. Uggla. 1989. Fatal toxoplasmosis in dogs. *Journal of the American Animal Hospital Association* 25:659-664.
- Faulkner, C.T., S. Patton, L. Munson, E.M. Johnson, and T.J. Coonan. 2001. *Angiocaulus gubernaculatus* in the island fox (*Urocyon littoralis*) from the

Recovery Plan for Four Subspecies of Island Fox

- California Channel Islands and comments on the diagnosis of Angiostrongylidae nematodes in canid and mustelid hosts. *Journal of Parasitology* 87(5):1174-1176.
- Fausett, L.L. 1982. Activity and movement patterns of the island fox, *Urocyon littoralis* Baird 1857 (Carnivora: Canidae). Ph. D. dissertation, University of California, Los Angeles. 132 pp.
- Fausett, L.L. 1993. Activity and movement of the island fox, *Urocyon littoralis*, on Santa Cruz Island, California. Pages 391-404 in Hochberg, F.G., (ed.), Third California Islands Symposium: Recent Advances in Research on the California Islands. Santa Barbara Museum of Natural History: Santa Barbara, California.
- Field, C.B., G.C. Daily, F.W. Davis, S. Gaines, P.A. Matson, J. Melack, and N.L. Miller. 1999. Confronting climate change in California. Ecological impacts on the Golden State. A report of the Union of Concerned Scientists, Cambridge, Massachusetts, and the Ecological Society of America, Washington, DC.
- Garcelon, D.K. 1999. Island fox population analysis and management recommendations. Unpublished report submitted to: Southwest Division, Naval Facilities Engineering Command, San Diego, California.
- Garcelon, D.K. and G.A. Schmidt. 2005. Island fox monitoring and demography on San Nicolas Island – 2004. Unpublished report prepared by the Institute for Wildlife Studies, Arcata, California. 21 pp + appendix.
- Garcelon, D.K., G.W. Roemer, and G.C. Brundige. 1991. The biology of the island fox on San Clemente and Santa Catalina islands, California. A final report submitted to the Nongame Bird and Mammal Section, Wildlife Management Branch, California Department of Fish and Game, Contract No. FG-9402.
- Garcelon, D.K., K.P. Ryan, and P.T. Schuyler. 2005. Application of techniques for feral pig eradication on Santa Catalina Island, California. Pages 331-340 in Garcelon, D. K., and C. A. Schwemm, (eds.), Proceedings of the Sixth California Islands Symposium. National Park Service Technical Publication CHIS-05-01, Institute for Wildlife Studies, Arcata, California.
- Garcelon, D.K., G.W. Roemer, R.B. Philips, and T.J. Coonan. 1999. Food provisioning by island foxes, *Urocyon littoralis*, to conspecifics caught in traps. *The Southwestern Naturalist* 44(1):83-86.
- R.K. Wayne, and B.J. Gonzales. 1992. A serologic survey of the island fox (*Urocyon littoralis*) on the Channel Islands, California. *Journal of Wildlife Management* 28(2):223-229.

Recovery Plan for Four Subspecies of Island Fox

- Garcelon, D.K., L. Munson, T.W. Vickers and D. Simmons. 2008. Research study on the pathology and veterinary services for island fox on San Clemente Island. Unpublished report prepared for Naval Facilities Engineering Command, Southwest, San Diego, California. Institute for Wildlife Studies, Arcata, California. 26pp + appendix.
- Garcia and Associates. 2008. Island Fox (*Urocyon littoralis clemente*) Monitoring and Research on Naval Auxiliary Land Field, San Clemente Island, California. Unpublished report. 34 pp.
- George, S.B. and R.K. Wayne. 1991. Island foxes. A model for conservation genetics. *Terra* 30:18-23.
- Gilbert, D.A., N. Lehman, S.J. O'Brien, and R.K. Wayne. 1990. Genetic fingerprinting reflects population differentiation in the California Channel Island fox. *Nature* 344:764-767.
- Goldstein, D.B., G.W. Roemer, D.A. Smith, D.E. Reich, A. Bergman, and R.K. Wayne. 1999. The use of microsatellite variation to infer population structure and demographic history in a natural model system. *Genetics* 151:797-801.
- Gray, M.M. 2002. Impact of a bottleneck on genetic diversity in the Channel Island fox (*Urocyon littoralis*) and genetic assessment of relatedness among individuals in a captive breeding program. Unpublished master's thesis, University of California, Los Angeles. 79 pp.
- Gray, M.M., G.W. Roemer, and E. Torres. 2001. Genetic assessment of relatedness among individuals in the island fox (*Urocyon littoralis*) captive breeding program. Unpublished manuscript on file at park headquarters, Channel Islands National Park, Ventura, California. 23 pp.
- Grinnell, J., J.S. Dixon, and J.M. Linsdale. 1937. Fur-bearing mammals of California: their natural history, systematic status, and relations to man. University of California Press, Berkeley, California, 2:452-471.
- Guthrie, D.A. 1993. New information on the prehistoric fauna of San Miguel Island, Pages 405-416 in Hochberg, F. G., (ed.), Third California Islands Symposium: Recent Advances in Research on the California Islands. Santa Barbara Museum of Natural History, Santa Barbara, California. 661 pp.
- Guttilla, D.A. 2007. Effects of sterilization on movement, home range behavior, and habitat use of feral cats on Santa Catalina Island, California. A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science in Biological Science, California State University, Fullerton.

Recovery Plan for Four Subspecies of Island Fox

- Guttilla, D.A. and P. Stapp. 2010. Effects of sterilization on movements of feral cats at a wildland-urban interface. *Journal of Mammalogy*, 91(2):482-489.
- Hall, E.R. and K.R. Kelson. 1959. *Mammals of North America, Volume II*. Ronald Press Company, New York.
- Hallbrooks, R.D., L.J. Swango, P.R. Schnurrenberger, F.E. Mitchell, and E.P. Hill. 1981. Response of gray foxes to modified live-virus canine distemper vaccines. *Journal of the American Veterinary Medicine Association* 179:1170-1174.
- Hanson, C.C. 2012. The removal of feral cats from San Nicolas Island, California to Protect Native and Endemic Species: 2011 annual report. Unpublished report, Island Conservation, Santa Cruz, California. 12 pp.
- Hochberg, M., S. Junak, R. Philbrick, and S. Timbrook. 1979. Botany. Pages 5.1-5.91 in: Power, D.M., (ed.), *Natural Resources Study of the Channel Islands National Monument, California*. Report submitted to the National Park Service. Santa Barbara Museum of Natural History, Santa Barbara, California.
- Hoover, E.A. and J.I. Mullins. 1991. Feline leukemia virus infection and diseases. *Journal of the American Veterinary Medical Association* 199:1287-1297.
- Hudgens, B.R. and J. Sanchez. 2009. Island fox monitoring on Santa Cruz Island, California, June 2008 - March 2009. Unpublished report by the Institute for Wildlife Studies, Arcata, California, for The Nature Conservancy, Santa Cruz Island Project, Ventura, California. 23 pp.
- Hudgens, B. R., and D. K. Garcelon. 2014. Island Fox Monitoring and Demography on San Nicolas Island-2013. Unpublished report prepared by the Institute for Wildlife Studies, Arcata, California. 28 pp.
- Hudgens, B. R., T.W. Vickers, D. K. Garcelon, and J. N. Sanchez. 2011. Epidemic Response Plan for San Clemente Island Foxes. Unpublished report prepared for the U. S. Navy by the Institute for Wildlife Studies, Arcata, California.
- Institute for Wildlife Studies. 2006. 2005 Annual Report to the U.S. Fish and Wildlife Service for Activities Conducted on Island Foxes (*Urocyon littoralis*) Under Permit No. TE-744878-14. Unpublished report prepared by the Institute for Wildlife Studies, Arcata, CA. 39pp.
- Institute for Wildlife Studies. 2007. 2006 annual report to the U.S. Fish and Wildlife Service for activities conducted on island foxes (*Urocyon littoralis*)

Recovery Plan for Four Subspecies of Island Fox

under permit no. TE-744878-14. Unpublished report by the Institute for Wildlife Studies, Arcata, CA. 51pp

- Ikeda, Y., K. Nakamura, T. Miyazawa, M.-C. Chen, T.-F. Kuo, J.A. Lin, T. Mikami, C. Kai, and E. Takahashi. 2001. Seroprevalence of canine distemper virus in cats. *Clinical and Diagnostic Laboratory Immunology* 8:641–644.
- [IPCC] Intergovernmental Panel on Climate Change. 2007. *Climate change 2007: the physical science basis. Summary for policymakers. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, IPCC Secretariat, World Meteorological Organization and United Nations Environment Programme, Geneva, Switzerland.*
- Jackman, R.E., W.G. Hunt, D.E. Driscoll, and F.J. Lapsansky. 1994. Refinements to selective trapping techniques: a radio-controlled bow net and power snare for bald and golden eagles. *Journal of Raptor Research*. 28:268-273.
- Johnson, D.L. 1978. The origin of island mammoths and the Quaternary land bridge history of the northern Channel Islands, California. *Quaternary Research* 10:204-225.
- Johnson, D.L. 1980. Episodic vegetation stripping, soil erosion, and landscape modification in prehistoric and recent historic time, San Miguel Island, California. Pages 103-122 in: Power, D.M. (ed.), *The California Islands: Proceedings of a Multidisciplinary Symposium*. Santa Barbara Museum of Natural History, Santa Barbara, California.
- Kelly, T.R. and J.M. Sleeman. 2003. Morbidity and mortality of red foxes (*Vulpes vulpes*) and gray foxes (*Urocyon cinereoargenteus*) admitted to the Wildlife Center of Virginia, 1993–2001. *Journal of Wildlife Diseases* 39:467–469.
- Kiff, L.F. 1980. Historical changes in resident populations of California islands raptors. Pages 651-674 in: Power, D.M., (ed.), *The California Islands: Proceedings of a Multidisciplinary Symposium*. Santa Barbara Museum of Natural History, Santa Barbara, California.
- King, J.L. and C.L. Duncan. 2008. 2007 annual report to the U.S. Fish and Wildlife Service for activities conducted by the Santa Catalina Island Conservancy on Catalina Island foxes (*Urocyon littoralis catalinae*) under permit no. TE-090990-0. Unpublished report on file at USFWS Regional Office Carlsbad, CA. 37pp.

Recovery Plan for Four Subspecies of Island Fox

- King, J.K. and C.L. Duncan. 2012. 2011 Annual Report to the U.S. Fish and Wildlife Service for Activities Conducted by the Catalina Island Conservancy on Catalina Island. 45 pp.
- King, J.K., and C.L. Duncan. 2013. Catalina Island Recovery Program: 2012 Annual Report. Conservation Technical Report CON/2014-02. Catalina Island Conservancy, Avalon, California 41. Pp.
- King, J.L. and C.L. Duncan. 2014. Catalina Island fox recovery program: 2013 Annual Report. Conservation Technical Report CON/2014-02. Catalina Island Conservancy, Avalon, California.
- Klinger, R.C., P.T. Schuyler, and J.D. Sterner. 1994. Vegetation response to the removal of feral sheep from Santa Cruz Island. Pages 341-350 in: Halvorson, W.L, and G.J. Maender, (eds.), The Fourth California Channel Islands Symposium: Update on the Status of Resources. Santa Barbara Museum of Natural History, Santa Barbara, California.
- Kohlmann, S.G., G.A. Schmidt, and D.K. Garcelon. 2005. A population viability analysis for the island fox on Santa Catalina Island, California. *Ecological Modeling* 183:77-94.
- Kovach, S.D. and R.J. Dow. 1981. Status and ecology of the island fox on San Nicolas Island, 1980. Unpublished report. Pacific Missile Test Center, Point Mugu, California 93042.
- Kovach, S.D. and R.J. Dow. 1982. Ecology of island fox and feral cats on San Nicolas Island. Pages 439-453 in Dow, R.J. (ed.), Biennial Mugu Lagoon/San Nicolas Island Ecological Research Symposium. Pacific Missile Test Center, Point Mugu, California, October 28-29, 1981.
- Kovach, S.D. and R.J. Dow. 1985. Island fox research, San Nicolas Island, Ventura County, California: 1984 Annual Report. Unpublished report to the California Department of Fish and Game. 19 pp.
- Latta, B. 2001. Channel Island Golden Eagle Research and Translocation Project. 1999-2001 Progress Report. Santa Cruz Predatory Bird Research Group.
- Latta, B.C., D.E. Driscoll, J.L. Linthicum, R.E. Jackman, and G. Doney. 2005. Capture and translocation of golden eagles from the California Channel Islands to mitigate depredation of endemic island foxes. Pages 341-350 in Garcelon, D. K., and C. A. Schwemm, (eds.), Proceedings of the Sixth California Islands Symposium. National Park Service Technical Publication CHIS-05-01, Institute for Wildlife Studies, Arcata, California.

Recovery Plan for Four Subspecies of Island Fox

- Laughrin, L.L. 1971. Preliminary account of the island fox. Supported in part by Federal Aid in Wildlife Restoration, project W-54-R, Special Wildlife Investigations.
- Laughrin, L.L. 1973. California island fox survey. State of California Department of Fish and Game, Wildlife Management Branch Administrative Report 73-3.
- Laughrin, L.L. 1977. The island fox: a field study of its behavior and ecology. Ph.D. dissertation, University of California, Santa Barbara. 83 pp.
- Laughrin, L.L. 1978. Status report on the San Nicolas Island Fox. University of California, Santa Barbara.
- Laughrin, L.L. 1980. Population and status of the island fox. Pages 745-750 in Power, D. M., (ed.), The California Islands: proceedings of a multidisciplinary symposium. Santa Barbara Museum of Natural History, Santa Barbara, California.
- Lin, D., D.D. Bowman, and R.H. Jacobson. 1992. Immunological changes in cats with concurrent *Toxoplasma gondii* and feline immunodeficiency virus infections. *Journal of Clinical Microbiology* 30:17-24.
- MacArthur, R.H. and E.O. Wilson. 1967. The Theory of Island Biogeography. Princeton University Press. 203 pp.
- Macdonald, N. and K. Walker. 2007. A New Approach for Ungulate Eradication: A Case Study for Success. Prohunt Incorporated, Ventura, California. 60 pp.
- Menke, A.S. and D.R. Miller. 1985. Entomology of the California Channel Islands. Santa Barbara Museum of Natural History, Santa Barbara, California 93105.
- Merriam, C.H. 1888. Description of a new fox from southern California. *Proceedings of the Biological Society of Washington* 4:135-138.
- Merriam, C.H. 1903. Eight new mammals from the United States. *Proceedings of the Biological Society of Washington* 16:73-78.
- Miller, P.S., G.W. Roemer, J. Laake, C. Wilcox, and T.J. Coonan. 2001. Draft island fox demographic workshop report. Unpublished manuscript on file at park headquarters, Channel Islands National Park, Ventura, California.
- Moore, C.M. and P.W. Collins. 1995. *Urocyon littoralis*. *Mammalian Species* No. 489:1-7.

Recovery Plan for Four Subspecies of Island Fox

- Moretti, L.D., A.V. da Silva, M.G. Ribeiro, A.C. Paes, and H. Langoni. 2006. *Toxoplasma gondii* genotyping in a dog co-infected with distemper virus and ehrlichiosis rickettsia. *Revista do Instituto de Medicina Tropical de São* 48:359-363.
- Morrison, S.A., N. Macdonald, K. Walker, L. Lozier, and M.R. Shaw. 2007. Facing the dilemma at eradication's end: uncertainty of absence and the Lazarus effect. *Frontiers in Ecology and the Environment* 5(5):271-276.
- Munson, L. 2010. Diseases of island foxes. Pp. 129-143 in: Coonan, Schwemm and Garcelon.
- National Park Service. 1985. General management plan for Channel Islands National Park, Vol. 1. National Park Service, Ventura, California.
- National Park Service. 1991. Natural Resources Management Guidelines. NPS-77. National Park Service, Washington, D.C.
- National Park Service. 1998. Final supplement to the final environmental impact statement: resources management plan for improvement of water quality and conservation of rare species and their habitats on Santa Rosa Island. National Park Service, Channel Islands National Park, Ventura, California. 34 pp.
- National Park Service. 2013. Draft General Management Plan/Wilderness Study/Environmental Impact Statement. National Park Service, Channel Islands National Park, Ventura, California. 491 pp.
- Newsome, S.D., P.W. Collins, T.C. Rick, D.A. Guthrie, J.M. Erlandson and M. L. Fogel. 2010. Pleistocene to historic shifts in bald eagle diets on the Channel Islands, California. *Proceedings of the National Academy of Sciences* 107:9246-9251.
- O'Brien, S.J. and J.F. Evermann. 1988. Interactive influence of infectious disease and genetic diversity in natural populations. *Trends in Ecology and Evolution* 3:254-59.
- O'Gara, B.W. and D.C. Getz. 1986. Capturing golden eagles using a helicopter and net gun. *Wildlife Society Bulletin* 14:400-402.
- O'Malley, P.G. 1994. Animal husbandry on the three southernmost Channel Islands: a preliminary overview, 1820-1950. Pp. 157-164 in: Halvorson, W.L., and G.J. Maender, (eds.), *The Fourth California Channel Islands Symposium: Update on the Status of Resources*. Santa Barbara Museum of Natural History, Santa Barbara, California.

Recovery Plan for Four Subspecies of Island Fox

- Parkes, J.P., D.S.L. Ramsey, N. MacDonald, K. Walker, S. McKnight, B.S. Cohen, and S.A. Morrison. 2010. Rapid eradication of feral pigs (*Sus scrota*) from Santa Cruz Island Biological Conservation.
- Peart, D., D.T. Patten, and S.L. Lohr. 1994. Feral pig disturbance and woody species seedling regeneration and abundance beneath coast live oaks (*Quercus agrifolia*) on Santa Cruz Island, California. Pages 313-322 in Halvorson, W.L, and G.J. Maender, (eds.), The Fourth California Channel Islands Symposium: Update on the Status of Resources. Santa Barbara Museum of Natural History, Santa Barbara, California.
- Pedersen, N.C. and J.E. Barlough. 1991. Clinical overview of feline immunodeficiency virus. *Journal of the American Veterinary Medical Association* 199:1298-1305.
- Phillips, R.B. and R.H. Schmidt. 1997. Feral cat management plan for Naval Auxiliary Landing Field, San Clemente Island, California. Submitted to the Natural Resources Office, NAS North Island, U.S. Navy by the Department of Fisheries and Wildlife, Utah State University, Logan, Utah.
- Phillips, R.B., C.S. Winchell, and R.H. Schmidt. 2007. Dietary overlap of an alien and native carnivore on San Clemente Island, California. *Journal of Mammalogy* 88(1):173-180.
- Propst, B. 1975. A population survey of the Santa Catalina Island fox. A report to the California Department of Fish and Game, Nongame Wildlife Investigation Project W-54-R-8, Job I-1.10. 8 pp.
- Rick, T.C., J.M. Erlandson, R.L. Vellanoweth, T.J. Braje, P.W. Collins, D.A. Gunthrie, and T.W. Stafford Jr. 2009. Origins and antiquity of the island fox (*Urocyon littoralis*) on California's Channel Islands. *Quaternary Research* 71 (2009) 93-98.
- Roemer, G.W. 1999. The ecology and conservation of the island fox (*Urocyon littoralis*). Ph.D. dissertation, University of California, Los Angeles. 229 pp.
- Roemer, G.W., T.J. Coonan, D.K. Garcelon, C.H. Starbird, and J.W. McCall. 2000. Spatial and temporal variation in the seroprevalence of canine heartworm antigen in the island fox. *Journal of Wildlife Diseases* 36:723-728.
- Roemer, G.W. and R.K. Wayne. 2003. Conservation in conflict: the tale of two endangered species. *Conservation Biology* 17(5):1251-1260.

Recovery Plan for Four Subspecies of Island Fox

- Roemer, G.W., C J. Donlan, and F. Courchamp. 2002. Golden eagles, feral pigs and insular carnivores: how exotic species turn native predators into prey. *Proceedings of the National Academy of Sciences* 99:791-796.
- Roemer, G.W., D.K. Garcelon, T.J. Coonan, and C.A. Schwemm. 1994. The use of capture-recapture methods for estimating, monitoring, and conserving island fox populations. Pages 387-400 in: Halvorson, W.L, and G.J. Maender, (eds.), *The Fourth California Channel Islands Symposium: Update on the Status of Resources*. Santa Barbara Museum of Natural History, Santa Barbara, California.
- Roemer, G.W., D.A. Smith, D.K. Garcelon, and R.K. Wayne. 2001a. The behavioral ecology of the island fox (*Urocyon littoralis*). *Journal of Zoology* 255:1-14.
- Roemer, G.W., T.J. Coonan, D.K. Garcelon, J. Bascompte, and L. Laughrin. 2001b. Feral pigs facilitate hyperpredation by golden eagles and indirectly cause the decline of the island fox. *Animal Conservation* 4:307-318.
- Rubin, E.S., V.J. Bakker, M.G. Efford, B.S. Cohen, J.A. Stallcup, W.D. Spencer, and S.A. Morrison. 2007. A population monitoring framework for five subspecies of island fox (*Urocyon littoralis*). Prepared by the Conservation Biology Institute and The Nature Conservancy for the Recovery Coordination Group of the Integrated Recovery Team. 145 pp + maps + app.
- Schmidt, G.A., J. Fox, and D.K. Garcelon. 2007. Island fox recovery on Santa Cruz Island, California, December 2000 - January 2007. Unpublished report by the Institute for Wildlife Studies, Arcata, California, for The Nature Conservancy, Santa Cruz Island Project, Ventura, California. 63 pp.
- Schmidt, G.A., B.R. Hudgens, and D.K. Garcelon. 2005a. Island fox monitoring and research on Naval Auxiliary Landing Field, San Clemente Island, California, Final Report: May 2004–June 2005. Unpublished report prepared by the Institute for Wildlife Studies, Arcata, California. 31pp.
- Schmidt, G.A., T.W. Vickers, and D.K. Garcelon. 2005b. Island fox recovery efforts on Santa Catalina Island, California, 1 April 2004 – 30 April 2005. Unpublished report by the Institute for Wildlife Studies, Arcata, California for the Ecological Restoration Department, Santa Catalina Island Conservancy, Avalon, CA. 38pp.
- Schmidt, G. A., C. M. Steele, T. W. Vickers, and D. K. Garcelon. 2004. Island fox recovery efforts on Santa Catalina Island, California, October 2002-October 2003. Unpublished report prepared by the Institute for Wildlife Studies, Arcata, CA. 38pp.

Recovery Plan for Four Subspecies of Island Fox

- Schwemm, C.A. 2007. Summary Report: Ninth annual meeting of the Island Fox Working Group, June 19-21, 2007. Unpublished Report on file at headquarters, Channel Islands National Park, Ventura, California. 70 pp.
- Schwemm, C.A. 2008. Summary Report: Tenth annual meeting of the Island Fox Working Group, June 24-26, 2008. Unpublished Report on file at headquarters, Channel Islands National Park, Ventura, California. 45pp.
- Sharpe, P. B. 2014. Bald Eagle Restoration on the California Channel Islands, January-December 2013, 12th Annual Report. Unpublished report prepared by the Institute for Wildlife Studies, Arcata, California for the National Park Service, Ventura, California. 25 pp.
- Sillero-Zubiri, C. and D.W. Macdonald. 2004. Foxes, wolves, jackals and dogs: status survey and conservation action plan. 2nd edition. IUCN, Gland, Switzerland.
- Smith, A.W., D.E. Skilling, N. Cherry, J.H. Mead, and D.O. Matson. 1998. Calicivirus emergence from ocean reservoirs: zoonotic and interspecies movements. *Emerging Infectious Diseases* 4:13-20.
- Sohn, R.A. and N. Thomas. 2005. Captive island fox parasite assessment and control on San Miguel, Santa Rosa and Santa Cruz Islands. Unpublished report prepared by U. S. Geological Survey, National Wildlife Health Center, Madison, Wisconsin. 67 pp.
- Sonsthagen, S.A., T.J. Coonan, B.C. Latta, G.K. Sage and S.L. Talbot. 2012. Genetic diversity of a newly established population of golden eagles on the Channel Islands, California. *Biological Conservation* 146:879-894.
- Spencer, W., E. Rubin, J. Stallcup, V. Bakker, B. Cohen, S. Morrison, and R. Shaw. 2006. Framework Monitoring Plan for the San Clemente Island Fox: With Specific Recommendations for the 2006 Field Season. Prepared for the United States Navy Region Southwest. 88 pp plus Appendices.
- Talbot, S.L., S.A. Sonsthagen, G.K. Sage, L.Pajot, B. Latta, T. Coonan, H. fitting and C.McIntyre. In preparation. Species verification and individual identification of eagle feathers dropped at Chanel Island fox depredation sites.
- Templeton, A.R. 1994. Biodiversity at the molecular genetic level: experiences from disparate macroorganisms. *Philosophical Transactions of the Royal Society of London Biological Sciences* 345(1311):59-64.
- Tenter, A.M., A.R. Heckerroth, and L.M. Weiss. 2000. *Toxoplasma gondii*: from animal to humans. *International Journal of Parasitology* 30:1217-1258.

Recovery Plan for Four Subspecies of Island Fox

- Thompson, C.M., E.L. Stackhouse, G.W. Roemer, and D.K. Garcelon. 1998. Home range and density of the island fox in China Canyon, San Clemente Island, California. U. S. Navy, Natural Resources Management Branch, Southwest Div., Nav. Fac. Eng. Command, San Diego, California. 31 pp.
- Timm, S.F., J.M. Stokely, T.B. Gehr, R.L. Peebles, and D.K. Garcelon. 2000. Investigation into the decline of island foxes on Santa Catalina Island. Report prepared for the Ecological Restoration Department, Santa Catalina Island Conservancy.
- Timm, S.F., W.D. Barker, S.A. Johnson, J.H. Sewell, P.B. Sharpe, G.A. Schmidt, and D.K. Garcelon. 2002. Island fox recovery efforts on Santa Catalina Island, California, September 2000-October 2001 annual report. Unpublished report prepared by the Institute for Wildlife Studies for the Ecological Restoration Department, Santa Catalina Island Conservancy. 52 pp.
- U.S. Fish and Wildlife Service. 2004. Endangered and threatened wildlife and plants; listing the San Miguel Island fox, Santa Rosa Island fox, Santa Cruz island fox, and Santa Catalina Island fox as endangered. Federal Register 69(44):10335-10353.
- U.S. Fish and Wildlife Service. 2005. Endangered and threatened wildlife and plants; final determination concerning critical habitat for the San Miguel Island fox, Santa Rosa Island fox, Santa Cruz island fox, and Santa Catalina Island fox as endangered. Federal Register 70(216):67924-67929.
- Vellanoweth, R.L. 1998. Earliest island fox remains on the southern Channel Islands: evidence from San Nicolas Island, California. *Journal of California and Great Basin Anthropology* 20(1):100-108.
- von Bloeker, Jr, J.C. 1967. Land mammals of the southern California islands. Pages 245-264 in: Philbrick, R.N., (ed.), *Proceedings of the Symposium on the Biology of the California Islands*. Santa Barbara Botanic Garden, Santa Barbara, California.
- Wayne, R.K., S.B. George, D. Gilbert, and P.W. Collins. 1991a. The Channel Island fox (*Urocyon littoralis*) as a model of genetic change in small populations. Pp. 639-649 in Dudley, E. C., (ed.), *The Unity of Evolutionary Biology: Proceedings of the Fourth International Congress of Systematic and Evolutionary Biology*, Vol. II. Dioscorides Press, Portland, Oregon.
- Wayne, R.K., S.B. George, D. Gilbert, P.W. Collins, S.D. Kovach, D. Girman, and N. Lehman. 1991b. A morphologic and genetic study of the island fox, *Urocyon littoralis*. *Evolution* 45(8):1849-1868.

Recovery Plan for Four Subspecies of Island Fox

Wilson, R.L. 1976. The status of the island fox, San Clemente Island. Progress Report 1. Natural Resources Program, Naval Undersea Center, San Diego, California.

PERSONAL COMMUNICATION

Booker, Melissa. 2013. U.S. Navy. Coronado, California.

Boser, Christina. 2014. The Nature Conservancy. Ventura, California

Coonan, Tim, J. 2006, 2007, 2011, 2014. National Park Service, Channel Islands National Park. Ventura, California

Ferrara, Francesca. 2014. U.S. Navy. Naval Base Ventura County, California.

Garcelon, Dave, K. 2014. Institute for Wildlife Studies. Arcata, California

Guttilla, Darcee. 2007. Santa Catalina Island Conservancy. Santa Catalina Island, California

King, Julie, K. 2014. Santa Catalina Island Conservancy. Santa Catalina Island, California

Sillett, Scott. 2004. Smithsonian Migratory Bird Center. Washington, D.C.

Schuyler, Peter. 2006. Independent Biologist, Santa Barbara, California

Trail, Pepper. 2014. Senior Forensic Scientist/Ornithologist, National Fish and Wildlife Forensics Laboratory, Ashland, Oregon. E-mail to Robert McMorran, U.S. Fish and Wildlife Service, Ventura, California, dated April 29, 2014.

Talbot, Sandra. 2014. Research Wildlife Geneticist, U.S. Geological Survey, Anchorage, Alaska, E-mail to Robert McMorran, U.S. Fish and Wildlife Service, Ventura, California, dated April 23, 2014.

Vickers, Winston. 2014. Institute for Wildlife Studies. Arcata, California

Walton, Brian. 2004. Santa Cruz Predatory Bird Research Group. Santa Cruz, California

Wolstenholme, Rachel. 2007. The Nature Conservancy. Ventura, California

VIII. Appendices

The following appendices provide information related to island fox recovery efforts. Recommendations provided in some of these the appendices were based upon the best information available at the time they were developed; however, we are aware that as new information arises, a previous recommendation may need to be revised or new recommendations may need to be developed.

These appendices are also provided to compile many sources of information related to island fox recovery into one location.

Recovery Plan for Four Subspecies of Island Fox

A. APPENDIX 1: CURRENT AND FORMER MEMBERS OF THE ISLAND FOX RECOVERY TEAM AND PRIMARY AUTHORS OF THIS RECOVERY PLAN

Island Fox Recovery Coordination Group

Brian Cypher	California State University Stanislaus, Endangered Species Recovery Program
Carlos de la Rosa	Santa Catalina Island Conservancy
David Graber	National Park Service
Devra Kleiman*	Zoo-Logic, LLC
Robert McMorran	U.S. Fish and Wildlife Service
Eric Morrissette	U.S. Fish and Wildlife Service
Peter Schuyler	Independent Biologist
Rebecca Shaw	The Nature Conservancy
Dale Steele	California Department of Fish and Wildlife
Rosie Woodroffe	University of California at Davis, Department of Wildlife Conservation Biology

*Deceased

Chairs of the Island Fox Technical Expertise Groups

Captive Population Management:

Peter Siminski The Living Desert

Ecosystem Restoration:

Lyndal Laughrin Santa Cruz Island Reserve

Fox Health:

Linda Munson* University of California at Davis

Genetics:

Colleen Lynch AZA Population Management Center

Golden Eagle:

Sandy Vissman U.S. Fish and Wildlife Service

Population Modeling:

Kathy Ralls Smithsonian National Zoological Park

Reintroduction:

Dan Blumstein University of California at Los Angeles

Wild population management:

Scott Morrison The Nature Conservancy

*Deceased

Recovery Plan for Four Subspecies of Island Fox

Integrated Island Fox Recovery Team

Andelt, William	Coppelli, Susan	Gabriel, Mourad
Asa, Cheryl	Cory, Coleen	Galipeau, Russell
Bakker, Vickie	Cypher, Brian	Garcelon, David
Baldwin, Sandra	Daily, Marla	Graber, Dave
Bauman, Karen	Daugharty, Kirin	Gray, Melissa
Blumenshine, Karen	Davis, Heather	Green, Michele
Blumstein, Dan	Dearborn, Keri	Guglielmino, Angela
Boyce, Jennifer	de la Rosa, Carlos	Guieb, Ruben
Breen, Kevin	Denney, Richard	Guttilla, Darcee
Bremner-Harrison, Sam	Dennis, Mitchell	Haight, Bob
Brock, Kelly	deSpain, Forrest	Hall, Lynn
Calkins, Betsy	Detrich, Phil	Hanna, Shari
Carlstead, Kathy	Doak, Dan	Hill, Karl
Charlton, Josh	Drake, Lisa	Horiszny, Sheri
Chatfield, Jennifer	Dratch, Peter	Hudgens, Brian
Christianson, Matt	Duncan, Calvin	Johnson, Heather
Clifford, Deana	Efford, Murray	Keitt, Brad
Collins, Paul	Faulkner, Kate	Kimble, Katie
Comrack, Lyann	Ferrara, Francesca	King, Julie
Coonan, Tim	Fox, Jodi	Kleiman, Devra*

Recovery Plan for Four Subspecies of Island Fox

Kunkel, Kyran	Patton, Sharon	Smith, Grace
Latta, Brian	Power, Paula	Smith, Tessa
Laughrin, Lyndal	Powers, Robyn	Sovada, Marsha
Lea, Amanda	Provinsky, Stephanie	Steele, Dale
Leslie, Elaine	Ralls, Katherine	Swarts, Hilary
Little, Annie	Randall, Kara	Thomas, Nancy
Littlefield, Ben	Ruane, Martin	Varsik, Alan
Lynch, Colleen	Roemer, Gary	Vermeer, Lotus
Mazet, Jonna	Rubin, Esther	Vickers, Winston
McCrary, Mike	Sanchez, Jessica	Vissman, Sandy
McMorran, Robert	Sandhaus, Estelle	Watson, Debbie
Meyer, Pat	Schuyler, Peter	Wayne, Bob
Miller, Phil	Schwemm, Cathy	Widmer, Ali
Morrison, Scott	Scott, Eric	Wilkerson, Cynthia
Morrisette, Eric	Scott, Kim	Willett, Mark
Moxie, Jeff	Shaw, Rebecca	Williams, Ian
Munson, Linda*	Sharpe, Peter	Wolstenholme, Rachel
Orrock, John	Siminski, Peter	Woodroffe, Rosie

*Deceased

B. APPENDIX 2: DISCUSSION OF THE RISK-BASED RECOVERY CRITERION

Models were developed for the San Miguel Island fox, Santa Rosa Island fox, Santa Cruz Island fox, and Santa Catalina Island fox to gain insight into the factors influencing the risk of extinction for each subspecies population over a 50 year time period (Bakker et al. 2009). The model is a two-stage (pup v. non-pup; i.e., considering only two life stages, pups (< 1 year old) and animals referred to as adults (> 1 year old)) stochastic matrix model. Demographic rates are simulated based on established relationships with environmental conditions such as golden eagle numbers, island fox densities, and weather conditions. The model carefully incorporates uncertainty in our knowledge of island foxes into model predictions.

Based on the output of model simulations, it is possible to predict the risk of a population reaching quasi-extinction using adult mortality rate and adult population size. These risk predictions can be plotted as isoclines (see Figures 1a through 1d: a graph has been created for each subspecies). Each isocline identifies the risk of the particular subspecies reaching the determined quasi-extinction level of 30 foxes over the determined timeframe of 50 years based on current mortality rates and population sizes.

To use these graphs to assess attainment of recovery criterion 1, one plots the average adult mortality rate against the average adult population size calculated over three (3) years along with their 80% confidence intervals. From the location of this point relative to the risk isoclines, one would be able to identify the current predicted risk of quasi-extinction for the subspecies based upon the model. Recovery criterion 1 is attained when the plotted point and its mortality and population size confidence intervals lie entirely below the isocline delineating 5% risk of quasi-extinction.

Management to avoid quasi-extinction rather than true extinction (e.g., zero individuals) helps account for uncertainty in models, especially uncertainty associated with population dynamics at small sizes, and it focuses efforts on maintaining populations at levels at which management action is most feasible. This approach is used commonly in estimating risk for a population (Morris and Doak 2002), and has been used in the development of risk-based recovery criteria (e.g., Northern right, Fin, and Sperm whales). The choice of an appropriate threshold depends on a range of biological and socio-political factors (Burgman et al. 1993). A quasi-extinction threshold of 30 individuals was selected for each of the federally listed island fox subspecies populations.

Note: The isoclines provided are to serve for illustrative purposes and to provide a visual reference to estimate the risk of quasi-extinction based upon the appropriate parameters. The actual risk of quasi-extinction is to be calculated.

Recovery Plan for Four Subspecies of Island Fox

The isoclines associated with the risk of quasi-extinction for each subspecies should be adjusted with new information, as necessary.




Figure 1a: Risk of the San Miguel Island population reaching quasi-extinction using adult mortality rate and adult population size.




Figure 1b: Risk of the Santa Rosa Island population reaching quasi-extinction using adult mortality rate and adult population size. Because the PVA model was not parameterized using data from Santa Rosa Island, this contour plot assumes that foxes on Santa Rosa Island have survival rates similar to foxes on Santa Cruz and San Miguel islands and reproductive rates similar to those on San Miguel Island.




Figure 1c: Risk of the Santa Cruz Island population reaching quasi-extinction using adult mortality rate and adult population size.




Figure 1d: Risk of the Santa Catalina Island population reaching quasi-extinction using adult mortality rate and adult population size. Because the PVA model was not parameterized using data from Santa Catalina Island, this contour plot assumes that foxes on Santa Catalina Island have survival rates similar to foxes on Santa Cruz and San Miguel islands and reproductive rates similar to those on San Miguel Island. We assumed no rainfall effects on survival rates.

Literature Cited

- Bakker, V.J., D.F. Doak, G.W. Roemer, D.K. Garcelon, T.J. Coonan, S.A. Morrison, C. Lynch, K. Ralls, and M.R. Shaw. 2009. Incorporating ecological drivers and uncertainty into a demographic population viability analysis for the Island Fox (*Urocyon littoralis*). *Ecological Monographs* 79(1):77-108.
- Burgman, M.A., S. Ferson, and H.R. Akcakaya. 1993. Risk assessment in conservation biology. Chapman and Hall, London, UK.
- Morris, W.F. and D.F. Doak. 2002. Quantitative conservation biology: theory and practice of population viability analysis.

Description of the model used to calculate the risk-based recovery criterion

This appendix is intended to briefly describe the methods and results of the updated island fox PVA upon which some of the recovery planning is based. The manuscript describing demographic analyses and population simulations making up the PVA is listed below:

- Bakker, V.J., D.F. Doak, G.W. Roemer, D.K. Garcelon, T.J. Coonan, S.A. Morrison, C. Lynch, K. Ralls, and M.R. Shaw. 2009. Incorporating ecological drivers and uncertainty into a demographic population viability analysis for the Island Fox (*Urocyon littoralis*). *Ecological Monographs* 79(1):77-108. Appendix 3: Technical Analysis Request 2.1 Development of Population Monitoring Plans for Free-Ranging Island Foxes

C. APPENDIX 3: GUIDELINES AND RECOMMENDATIONS ON VACCINATION PROTOCOLS, COLLECTION OF HEALTH DATA, AND FUTURE RESEARCH NEEDS RELATED TO FOX HEALTH.

This appendix includes additional details associated with Recovery Actions 1.2 to 1.4 of the Recovery Plan for Four Subspecies of Island Fox and was developed in coordination with the Fox Health Technical Expertise Group.

Recovery Action 1.2 – Avoid introduction of new pathogens, or novel strains of existing pathogens, to the Channel Islands by restricting or regulating movements of wild and domestic animals to the islands.

Movement of both domestic and wild mammals to the northern Channel Islands is not permitted under Channel Islands National Park regulations, although a small number of domestic dogs have been brought to Santa Rosa and Santa Cruz Islands in association with the various forms of hunting operations. Additionally, there is a chronic risk of animals (particularly domestic dogs, cats, and rats) being brought ashore unofficially from boats.

Where there is a clear benefit to bringing such animals to the islands, veterinary experts familiar with threats to island foxes should be consulted to develop appropriate quarantine or containment protocols.

Restricting animal movements to Santa Catalina Island is more difficult than to the northern Channel Islands because Santa Catalina Island is composed entirely of private land, has thriving resident and visitor populations, and has easily accessible regularly scheduled transport to and from the mainland. Efforts to minimize animal movement to the island could include education campaigns, developing partnerships with the City of Avalon and others, and adopting regulations restricting such movement and controlling feral cats.

Education campaigns should be extended to mainland-based boaters that may inadvertently transport stowaway wildlife such as raccoons to the islands.

Management and Removal of Introduced Mammals

All islands should have protocols in place to deal with invasive animal introductions.

- Protocols should be communicated to non-fox personnel so that proper steps are taken in the event of discovery of invasive animals on boats or on an island.
- Parties involved with protocol development should be cognizant that return of wild animals to the mainland for release back into the wild may risk introduction of unique infectious agents (such as *Spirocerca* or island-evolved viral strains) to naïve mainland wildlife populations, and is not advisable.

- Protocols should address: 1) requiring return of a boat to the mainland before exit of a stowaway animal; 2) capture protocols developed with other agencies which might be involved (depending on the island) such as the California Department of Fish and Wildlife (CDFW), the U.S. Navy, or the City of Avalon; 3) preconditions, indications, and protocols for lethal removal which should be developed with input from CDFW; 4) samples such as blood and feces which should be taken from captured invasive animals; 5) protocols for determining whether animals which have been lethally removed should be necropsied or otherwise tested for disease.

Recovery Action 1.3 – Implement Prophylactic Management to avoid extinction or quasi-extinction of wild populations in the event of devastating epidemics.

Any prophylactic vaccination against canine distemper virus (CDV) would need to take into account serological evidence that a strain of CDV is currently circulating within the wild populations on Santa Cruz and Santa Catalina Islands with no documented ill effects (strains with similarly benign effects occurred on San Miguel Island and Santa Rosa Island but may have been eliminated by vaccination when all foxes were taken into captivity and vaccinated). These strains may confer immunity to more dangerous strains; their eradication through vaccination should be avoided if possible. Maintaining a “core group” of a minimum of 80-100 CDV vaccinated individuals of each subspecies should allow such viruses to continue circulating in the unvaccinated proportion of each population. Larger numbers of animals may be vaccinated within “core” areas if managers wish to increase the likelihood of higher survival percentages in the event of a CDV epidemic.

Prophylactic vaccination against rabies should take into account the public health concerns that are associated with rabies, the risks to personnel incurred when handling potentially infected foxes for the purpose of vaccination in the event of an epidemic, and the potential desire of managers to increase the likelihood of survival percentages that are greater than the minimums required to avoid quasi-extinction. Maintaining a core group of a minimum of 80 – 100 individuals vaccinated against rabies, plus as many additional individuals in the population as can be opportunistically vaccinated, is advisable in order to maximize the number of foxes that would be expected to survive an epidemic. This approach would also reduce the level of threat to human and domestic animal populations that are in contact with island foxes.

1.3.1 – Test safety and antibody response to vaccination in captive island foxes under appropriate research protocols.

Recovery Plan for Four Subspecies of Island Fox

1.3.1.1 - Conduct CDV vaccination trials by administering two vaccinations at different bodily locations on island foxes during a single vaccination event.

Purevax™ recombinant CDV vaccine has been shown to be safe for use in island foxes, and to trigger seroconversion after administration of two intramuscular doses of vaccine given two weeks apart. However, field application of the vaccine would be more practicable if protection could reliably be achieved in the course of a single handling event, recently done successfully with wild dogs.

1.3.1.2 - Assess the efficacy of standard inactivated rabies vaccines in producing an antibody response in island foxes

This may be achievable using serum already banked from captive island foxes. Further studies are likely to be necessary if initial investigations suggest a poor immune response to rabies vaccination as implemented. Recombinant subunit rabies vaccines also need to be tested for safety and ability to induce an antibody response. Additionally, rabies vaccine delivered in bait should be assessed for ability to induce an antibody response.

1.3.1.3 – Vaccines against canine parvovirus and adenovirus should be tested on island foxes

Inactivated and monovalent modified live vaccines should be considered for testing with the knowledge that inactivated vaccines may not induce a robust antibody response and modified-live vaccines may induce disease. While an epidemic of parvovirus or adenovirus is unlikely to occur in the near future because of evidence of extensive exposure in the current populations, testing vaccines would be appropriate to have knowledge of their safety and effects if future vaccination is needed.

1.3.2 – On each island, maintain vaccination cover for rabies and CDV in at least 80 to 100 island foxes

All vaccinated foxes should be permanently marked, and detailed records should be maintained on the vaccination status of all island foxes handled, whether in the wild or in captivity. This is important to avoid confusion in the future interpretation of serology results.

Given uncertainties about the immune status of island foxes given single vs. repeated doses of vaccine, and the desirability of retaining nonlethal CDV strains in circulation on the islands if possible, it would be advisable to identify (and mark) the individuals to be vaccinated, and then re-vaccinate as many as possible. Because of attrition and failure to capture vaccinated individuals, new individuals may need to be vaccinated annually to maintain a

core population of vaccinated island foxes. Less systematic approaches risk protecting too few island foxes (if single doses of vaccine do not confer protection, and only a proportion can be recaptured for boosters), or too many island foxes (if single doses are effective but assumed not to be so).

Recovery Action 1.4 – Establish monitoring and response strategies to detect and manage infectious disease threats to island fox population persistence.

1.4.1 – Monitor to detect disease-related mortality

Ensure that management activities to avoid introducing new infectious diseases to island fox populations, and to avoid complete extinction in the face of devastating epidemics, are combined with mechanisms to detect, and respond to, any outbreaks that could occur.

All staff working on the Channel Islands, irrespective of their duties and expertise, should be trained to recognize and immediately report sick or dead carnivores, including island foxes. All staff should be trained to avoid exposure to diseases.

Necropsies should be performed as soon as possible for all carnivores that die from causes other than trauma to determine if the cause was an infectious disease potentially transmissible to island foxes.

1.4.2 – Annually collect blood samples from a proportion of island foxes on all islands to evaluate ongoing disease risks to island fox populations

Decisions about whether or not clinical interventions are justified will depend not only on information on mortality but also on the immune status of the population concerned. For example, a recent study suggests that foxes on all islands have been exposed to CDV and canine parvovirus (Clifford et al. 2006), indicating some degree of population protection and therefore little need to intervene if a small number of deaths were detected. However, patterns of exposure are likely to be dynamic and in small isolated populations pathogen populations can easily die out leaving the host population entirely susceptible to reinfection a generation later.

Island fox blood samples should be collected from a representative proportion of each subspecies during each year. If achievable, these samples should be collected, at a minimum, from each island fox population during each year including:

- all radio-collared individuals;
- 25 individuals sampled in a previous year;
- up to 25 previously un-sampled adults from a variety of age classes and geographical location; and

Recovery Plan for Four Subspecies of Island Fox

- as many previously un-sampled yearlings and pups (aged >5 months to avoid maternal antibodies) as possible.

1.4.3 – Develop response strategies for responding to island fox deaths from infectious diseases known to represent serious threats to the persistence or recovery of the wild populations.

This strategy should include:

- a list of the veterinarians and other relevant experts willing and able to advise on appropriate management, and their emergency contact details;
- a protocol for expediting transportation of relevant personnel to the affected island(s);
- facilities, equipment and supplies on site for investigating an outbreak;
- facilities, equipment and supplies on site for vaccination, quarantine, or treatment if indicated;
- measures to contact and work with these individuals if a disease outbreak occurs; and
- a stockpile of traps to capture island foxes if needed for vaccination.

Intervention for rabies

Intervention in a rabies epidemic may or may not be advisable, and is dependent on many factors. These include the percentage of the population that had been previously vaccinated, the geographic extent of detected disease, the availability of oral bait-based vaccine, and the number of personnel available to implement an intervention strategy who are vaccinated against rabies and properly trained. Immediate vaccination of all wild island foxes on the affected island, preferably commencing in areas remote from the index case, is advisable if oral bait-based vaccines are available, and have been tested and shown to be safe for island foxes. Trapping and vaccinating could pose unacceptable risks to personnel, and should not be utilized unless absolutely necessary, and only after consultation with the island fox health expertise group and other experts. The intensity of monitoring should be increased island-wide to detect both sick and dead animals. Euthanasia is recommended for sick animals. Rabies vaccination of human staff on the Channel Islands would also be advisable under these circumstances. Regular training should be conducted to minimize the likelihood of inadvertent human exposure to rabies in the event of an epidemic.

Intervention for canine distemper

Intervention in a CDV epidemic may or may not be advisable, and is dependent on many factors. These include the percentage of the population

Recovery Plan for Four Subspecies of Island Fox

that had been previously vaccinated, the geographic extent of detected disease, the availability of adequate isolation, and levels of staff training. Intervention could include immediate vaccination of all wild island foxes on the affected island, preferably commencing in areas remote from the index case, however trapping and vaccinating could potentially increase the spread of the disease via contaminated equipment or personnel. The intensity of monitoring should be increased island-wide to detect both sick and dead animals. Euthanasia of sick animals may be advisable after veterinary exam or consultation if isolation from other island foxes, both in the wild and in captivity, cannot be assured. Bringing apparently healthy foxes into captivity may also be inadvisable unless strict individual quarantine measures are possible, since some apparently healthy individuals could be incubating the disease. Screening of accumulated serum samples should be commenced immediately. Further blood sampling of the wild island fox population could be used to track the epidemic; serum from juveniles (aged 5-12 months) would be particularly helpful in this regard. Maintain a stockpile of Merial Purevax Ferret™ vaccine sufficient to protect a population in the face of a CDV outbreak at a central location and use as needed on whichever of the islands may be affected. Merial Purevax Ferret™ is the only vaccine against CDV that is both safe for use in island foxes and also triggers seroconversion; however, it is only available intermittently.

Intervention for parvovirus or adenovirus

Sick island foxes may be taken into captivity, isolated, and treated if proper isolation facilities and trained personnel are available. Scat surveys should be conducted to evaluate the prevalence of virus-induced diarrhea. If the virus were to be isolated from any sample, the strain should be investigated. As for CDV, screening of accumulated serum samples should be commenced immediately, as should prospective sampling to track the progression of the epidemic. Samples from juveniles (aged 5-12 months) would be particularly valuable for the latter purpose.

Literature Cited

- Clifford, D.L., J.A.K. Mazet, E.J. Dubovi, D.K. Garcelon, T.J. Coonan, P.J. Conrad, and L. Munson. 2006. Pathogen exposure in endangered island fox (*Urocyon littoralis*) populations: implications for conservation. *Biological Conservation* 131 (2006) 230-243.

D. APPENDIX 4: GUIDELINES FOR VACCINATION AND SAMPLING OF CAPTIVE AND WILD POPULATIONS

Review of appropriateness of vaccination

- Canine distemper and rabies are the diseases most likely to cause extinction, so vaccination is still considered important.
- Canine distemper virus (CDV) and rabies vaccines are available and safe for the foxes and the ecosystem.
- Because of continued endangered status, it is desirable to provide a safety net through core population vaccination.
- As long as monitoring is continuing, vaccination does not entail significant extra effort or cost.
- A proportion of foxes in larger populations should not be vaccinated against CDV in order to allow circulation of existent endemic wild viral strains.
- There are no concerns with assuring that wild rabies viral strains circulate in the population, so vaccination of a larger proportion of the population against this disease is acceptable for maximum fox and human protection in the event of an outbreak.
- Rabies has a public health aspect that may influence vaccination numbers, epidemic response, and locations of vaccinated core populations.
- Vaccinating a greater proportion of the population than the minimum required to avoid extinction (based on modeling) is a possible alternative to direct epidemic response and / or resumption of captive population maintenance in the event of an epidemic.

Vaccination recommendations:

- The vaccines to use are Merial's Purevax® Ferret Distemper Vaccine for CDV and Merial's Imrab 3® for rabies.
- Vaccinate all captive foxes for CDV and rabies
- Vaccinate all captured foxes against rabies EXCEPT radio-collared disease sentinels
- Vaccinate all opportunistically caught animals on islands with small populations (SMI and SRI) against CDV. When the populations reach 50% of the recovery goal, transition to the long-term core vaccination program.
- On islands with larger populations, a minimum of 20 radio-collared animals should not be vaccinated so that they can serve as a disease sentinel population for early detection of an epidemic. These individuals should be juveniles (1 – 2 years old if possible), and should be distributed as randomly as possible across the island.
- On islands with larger populations, vaccinate core group(s) totaling a minimum of 80 - 100 animals against CDV and rabies in strategic geographic location(s) in perpetuity.
- The decision to vaccinate animals radio-collared for purposes other than disease sentinels depends on management goals.

Figure 1 – CDV vaccination strategy





Figure 2 – Rabies vaccination strategy




FOX VACCINATION AND BLEEDING DECISION TREE

ALL radio-collared animals should be bled when the collar is placed and in all subsequent captures

Vaccinate ALL foxes against rabies EXCEPT radio-collared disease sentinels

“Core” is a minimum of 80 – 100 foxes vaccinated against CDV and rabies divided into 2-3 groups in strategic areas on all islands. On islands with larger populations, CDV vaccinate additional foxes up to the number that managers desire to have remaining after an epidemic.



Protocols for long term health monitoring

1. Continue collection and archiving of serum and blood from selected animals for health monitoring. The most valuable animals to bleed are pups and AC I, the disease sentinels, and other unvaccinated animals that were previously bled.

If populations are large enough:

- Sample as many previously unvaccinated (against CDV) age class I and pups > 5 mo old as possible (target number 25–50, including yearlings that are being radio-collared).
- Sample all unvaccinated (against CDV) radio-collared animals (target number 25-50)
- Sample 25-50 unvaccinated (against CDV) age class II or above animals that were bled in a previous year.

On islands with smaller populations:

- Sample all unvaccinated (against CDV) foxes over 5 months of age (up to a target number of 100).

Animals found to have naturally occurring (not vaccine-induced) CDV antibodies should have serology repeated yearly if possible in order to determine persistence and fluctuation of antibodies in response to exposure to wild viral strains

Conduct a serosurvey for antibodies to CDV, canine parvovirus (CPV), canine adenovirus (CAV), and *Toxoplasma* as often as yearly, but no less often than every 5 years. Using samples collected under the protocol in #1 above, test approximately 100 - 125 samples per island. The cost per sample is approximately \$50 - 75. If insufficient funds are available, then test 25 juveniles (giving priority to radio-collared animals) and 25 previously bled adults (giving priority to radio-collared animals).

2. Collect feces from 20 wild foxes on each island per year for parasite surveillance and store in 5 - 10% formalin for future analysis.

3. Record body weight and body condition score whenever foxes are handled.

4. Record injuries or other lesions and take photographs if possible

**E. APPENDIX 5: TECHNICAL ANALYSIS REQUEST 2.1 -
DEVELOPMENT OF POPULATION MONITORING PLANS FOR
FREE-RANGING ISLAND FOXES**

The draft “Island Fox Strategy for Recovery” dated June 14, 2006, calls for long-term monitoring of all wild populations via the best established methods, to monitor population dynamics and to ensure that population declines are detected rapidly and their causes understood. A technical analysis is required to identify the specific objectives (i.e. parameters, precision) of such a monitoring program and to develop statistically robust methods to meet these objectives. The following outlines the goals of this analysis, suggests which Technical Expertise Groups should be included, and provides a generalized process by which a monitoring plan can be developed for each island:

- 1) The goals of this analyses are to:
 - a) assess management objectives and needs related to the fox population on each island, and to recommend monitoring protocols designed specifically to address these management needs.
 - b) recommend monitoring protocols to collect population parameters necessary for development and refinement of PVAs that may be used to guide management activities.
 - c) recommend monitoring protocols to collect population parameters necessary to determine if recovery criteria, as adopted in the USFWS Recovery Plan, are reached.
 - d) recommend monitoring protocols to collect population parameters necessary for cross-island comparisons to increase our knowledge about island fox population dynamics.
 - e) recommend topics of future research modules which, although not part of a long-term monitoring plan, may be complementary to long-term monitoring activities.

In addition, each monitoring plan should include recommendations that facilitate the collection of animal health measures necessary to track population health as per the recommendation of the Fox Health Technical Expertise Group.

- 2) Although the actual development of each island-specific monitoring plan is anticipated to be carried out by qualified contractors, the following Technical Expertise Groups (“TEG”) should be involved in the conceptual development of monitoring plans and participate in a regular review of the plans as they are developed:
 - a) Population Modeling
 - b) Wild Population Management
 - c) Fox Health

Recovery Plan for Four Subspecies of Island Fox

A Task Force including the Chair of each of the above TEGs and/or their designated representatives should be available for discussion or review of issues as necessary. As work progresses, additional topics may be identified and included in plan development.

- 3) The following general steps and analyses should be included in developing each plan:
 - a) Collect and review information pertinent to each island, including past and current monitoring programs, monitoring data, and ecological and physical characteristics of the islands as they relate to monitoring needs and constraints.
 - b) Identify and articulate monitoring objectives using input from managers and the Task Force.
 - c) Analyze existing protocols to evaluate whether they are generating the appropriate parameters needed to meet current monitoring objectives. For example, a representation analysis of current trapping protocols should be conducted to determine how well trapping efforts represent habitat variability (e.g. vegetation, topography, distance to shoreline, or general location on the island) and management issues (e.g. distance to roads) on the island.
 - d) Develop recommended protocols, possibly with alternative scenarios, for
 - i) monitoring survival and cause specific survival
 - ii) sampling (trapping) to collect demographic data
 - e) Obtain input from managers and Task Force on the above protocols and alternative scenarios to determine feasibility and whether desired parameters will be generated.
 - f) Obtain input from a statistician on the above protocols and alternative scenarios to determine if methods are statistically robust.
 - g) Prepare draft and final monitoring plans for each island, allowing time for review and input from managers and Task Force.

This TAR relates to San Miguel, Santa Rosa, Santa Cruz, Santa Catalina, and San Nicolas Islands.

The final Technical Analysis Request 2.1 can be found in:

Rubin, E.S., V.J. Bakker, M.G. Efford, B.S. Cohen, J.A. Stallcup, W.D. Spencer, and S.A. Morrison. 2007. A population monitoring framework for five subspecies of island fox (*Urocyon littoralis*). Prepared by the Conservation Biology Institute and The Nature Conservancy for the Recovery Coordination Group of the Integrated Recovery Team. 145pp + maps + app.

Table 3-1 Monitoring goals for each island as stated and prioritized by island managers (from Rubin et al. 2007)

San Miguel	Management goal:	Parameter needed:
	<ol style="list-style-type: none"> 1. Monitor population status and trend. 2. Monitor threats to population (influences on population size and viability). 	<ol style="list-style-type: none"> 1. Index of island-wide changes in abundance (annual change in island-wide N or density) 2a. Survival (by age, gender, year) 2b. Cause-specific mortality (predation, disease, etc.) 2c. Reproduction (annual recruitment)
Santa Catalina	Management goal:	Parameter needed:
	<ol style="list-style-type: none"> 1. Monitor threats to population (influences on population size and viability). 2. Monitor population status and trend. 3. Estimate population size 	<ol style="list-style-type: none"> 1a. Survival (by age, gender, year) 1b. Cause-specific mortality (predation, disease, etc.) 2. Index of island-wide changes in abundance (annual change in island-wide N or density) 3. Island-wide abundance estimate (N)
Santa Rosa	Management goal:	Parameter needed:
	<ol style="list-style-type: none"> 1. Monitor population status and trend. 2. Monitor threats to population (influences on population size and viability). 3. Inform land management decisions. 	<ol style="list-style-type: none"> 1. Index of island-wide changes in abundance (annual change in island-wide N or density) 2a. Survival (by age, gender, year) 2b. Cause-specific mortality (predation, disease, etc.) 2c. Reproduction (annual recruitment) 3. Density by habitat type
Santa Cruz	Management goal:	Parameter needed:
	<ol style="list-style-type: none"> 1. Monitor threats to population (influences on population size and viability). 2. Monitor population status and trend. 3. Inform land management decisions. 	<ol style="list-style-type: none"> 1a. Survival (by age, gender, year) 1b. Cause-specific mortality (predation, disease, etc.) 2. Index of island-wide changes in abundance (annual change in island-wide N or density) 3. Survival and density by habitat type

Recovery Plan for Four Subspecies of Island Fox

F. APPENDIX 6: RECOMMENDATIONS FOR DOG ENTRY AND EXIT FROM THE NORTHERN CHANNEL ISLANDS (SANTA CRUZ, SANTA ROSA, AND SAN MIGUEL) AND THE NAVY ISLANDS (SAN CLEMENTE AND SAN NICOLAS)

Northern Islands:

Pre-Movement Quarantine

All dogs destined for shipment must be placed in a quarantine facility for 30 days before transport to any of the northern Channel Islands. The purpose of this quarantine facility is to prevent infection of the dogs after they have been tested and treated for parasites and infectious diseases. The facility should be: 1) isolated from contact with other carnivores, and 2) an all-in/all-out facility (no entry of new animals during the 30 days). If possible, the dogs should be individually housed and the substrate should be concrete or other surface that can be disinfected.

Vaccination

All dogs will have a current vaccination for the following:

DHPP(LC)- Modified Live Virus Vaccines

Canine distemper virus

Canine infectious hepatitis (canine adenovirus)

Canine parainfluenza virus

Canine parvovirus

Leptospirosis

Coronavirus

Killed Vaccines:

Rabies

Bordetella (kennel cough)

Canine influenza H3N8

The entire vaccination series will be completed at least one month, but no more than six months, prior to the dog's arrival on the island. Dogs vaccinated less than one month prior to transport may shed modified vaccine virus or viruses acquired through natural exposure before being protected by vaccines.

Dogs remaining on the island must be vaccinated annually with killed or subunit vaccines only (if available). If modified live vaccines must be used, the dogs should be kept in quarantine for 30 days after vaccination.

Parasites

All dogs must be negative for heartworms (*Dirofilaria immitis*) by DiroCheck® or SNAP® tests and be screened for microfilaria six months before being

Recovery Plan for Four Subspecies of Island Fox

transported to the island. Dogs must then be placed on an appropriate heartworm preventative and kept on preventative treatments while on-island. Recommended preventative treatments are Heartgard Plus® or Interceptor®.

All dogs must test negative for endoparasites prior to transport to the island. Three consecutive fecal samples over a 5 day period must be tested for endoparasites using both zinc and sugar floatation methods. Dogs with positive fecal tests should be treated with appropriate anthelmintics and then re-tested until they have three consecutive fecal samples test negative. If dogs are not individually housed, then all contact animals must also be treated and retested. Dogs should be rechecked by the same protocol annually.

During quarantine, all dogs must be checked for ectoparasites, including *Sarcoptes*, *Demodex* and *Otodectes* mites. If positive for any mite, the dogs should be appropriately treated and rechecked until negative. If dogs are not individually housed, all contact animals should also be treated and retested. Once negative for ectoparasites, the dogs should be placed on an appropriate preventative before being transported to the island. Recommended preventative treatments are Interceptor® or Frontline®.

Health Certificate

All dogs must be given a complete physical exam by a licensed veterinarian to confirm that they are in good general health and free of evidence of any infectious diseases within ten days of being transported to the island. The examination should include confirmation of vaccination status, confirmation of negative heartworm, endoparasite, and ectoparasite tests (including ear mites) and a negative Lyme disease test.

Post-transport Quarantine

Vaccinated animals can still be subclinically infected with infectious agents, such as canine distemper, and can therefore act as a source of infection for island foxes. Most dogs that mount an effective immune response to canine distemper virus clear the virus from their system within two weeks of exposure and cease to shed the virus. Also, most subclinical diseases caused by other agents usually become apparent within three weeks. Therefore all dogs transported to the islands must be quarantined on-island for three weeks in a location and facility that is inaccessible to island foxes. Dogs should be physically assessed by a veterinarian if possible before being released from quarantine.

Feces and urine from quarantined dogs should be disposed of in such a manner that foxes are not exposed to either feces or urine or the effluent from the disposal areas (see facility recommendations below).

Periodic fecal parasite checks should be done every 3 – 6 months while dogs are on the island.

Inter-Island Travel and Re-entry Requirements

The requirements for vaccination, parasite screening and quarantine are triggered each time a dog travels from the mainland to an island or travels between islands.

Dog Holding/Quarantine Facility Recommendations

Dog holding pens should be constructed in such a way as to have both inner fences and an outer perimeter fence separated from the dog holding pens by at least a 30 foot buffer space to prevent aerosol exposure to island foxes. In addition, multiple strands of electrified wire should be strung on the top of the fence to further discourage foxes from climbing into the facility. This fence design incorporates features that have been successful in keeping wild foxes from entering the Santa Catalina Island fox breeding facility.

Storage of food and equipment should be in a secure off-site location to further deter foxes from attempting entry into the dog holding facility.

Rubber boots should be worn inside the perimeter fence and a shoe bath and brush should be provided to step in as leaving to avoid carrying feces outside the perimeter on shoe bottoms.

All fecal waste from dog holding pens will be collected and disposed of in a waste digestion tank or sewage system. Effluent from digestion tanks will discharge underground into the earth. No effluent from the digestion tank will be discharged above ground. Pens will periodically be washed down with water.

If no concrete trough is present to catch runoff from runs, especially during rains, a ditch should be dug around the outside of the dog runs (inside of perimeter fence) to prevent rain runoff from carrying dog waste through or under perimeter fence.

Recommendations for dogs leaving the island:

Because domestic dogs residing on any of the northern Channel Islands may have acquired intestinal parasites from the foxes that may not have been controlled by routine monthly parasite preventatives, and might not be detected by routine fecal parasite checks, it is recommended that the dogs have pre-departure precautionary parasite treatment.

Treatments listed are by parasite type.

For Spirocerca:

Doramectin 10 mg/ml (1%) strength injectable (Dectomax or generic)

Dose = 0.4 mg/kg subcutaneously every 2 weeks beginning 6 months before departure from the island

Examples: A 20 kg (44 lb) dog would get 0.8 ml (ml's are the same as cc's) each dosing.

25 kg – 1.0 ml each dose

30 kg – 1.2 ml each dose

35 kg – 1.4 ml each dose

Ivermectin is the alternative treatment if doramectin is not available, but DO NOT GIVE BOTH DORAMECTIN AND IVERMECTIN:

Ivermectin 10 mg/ml (1%) strength injectable (Ivomec or generic)

Dose = 0.3 mg/kg every 2 weeks on same schedule as above.

Examples: A 20 kg (44 lb) dog would get 0.6 ml each dosing

25 kg – 0.75 ml each dose

30 kg – 0.9 ml each dose

35 kg – 1.05 ml each dose

Certain dogs are overly sensitive to Doramectin or Ivermectin, and can have symptoms of toxicity after dosing. Symptoms can be mild to severe (from mild lethargy or GI symptoms to seizures and collapse). A veterinarian should be consulted immediately if any significant symptoms occur with these or any of the drugs outlined.

For Mesocestoides:

Praziquantel 56.8 mg/ml strength injectable (Droncit or generic)

Caution - avoid giving this drug on the same day as Doramectin or Ivermectin because both may cause lethargy, GI upset, or other symptoms in some dogs

Dose = 0.2 ml / 2.3 kg (5 lb) body wt subcutaneously (to a maximum of 3 cc) – give 3 doses, 3 weeks apart.

Examples: A 20 kg (44 lb) dog would get 1.8 ml subcutaneously.

25 kg – 2.2 ml each dose

30 kg – 2.6 ml each dose

35 kg – 3.0 ml each dose

For Coccidia (Isospora, possibly others):

Sulfadimethoxine 500 mg tablets (Albon or generic)

Dose = 50 mg/kg the first day orally, then 25 mg/kg daily.

Treatment should start 3 weeks before departure and continue daily till 2 weeks after arrival.

Examples: A 20 kg (44 lb) dog would get 1,000 mg (2 of the 500 mg tablets) the first day, then 1 tablet (500 mg) each day thereafter

25 kg-1,250 mg (2 ½ tablets) the first day, then 625 mg (1 ¼ tabs) each day

30 kg-1,500 mg (3 tablets) the first day, then 750 mg (1 ½ tabs) each day

35 kg -1,750 mg (3 ½ tablets) the first day, then 875 mg (1 ¾ tabs) each day

For Uncinaria, Toxocara / Toxascaris:

Fenbendazole 100 mg/ml oral suspension (Panacur)

Dose = 50 mg/kg orally once a day for 3 days, then repeat the 3 day course

3-day treatment should be given 3 times prior to departure from the island at 3 week intervals

Examples: A 20 kg dog would get 1,000 mg (10ml) orally daily for 3 days each dosing

25 kg – 1,250 mg (12.5 ml)

30 kg – 1,500 mg (15 ml)

35 kg – 1,750 mg (17.5 ml)

The suggested schedules above are approximate. Multiple drugs should not be given on the same day, so scheduled treatments should be staggered. An exception is that the daily dose of sulfadimethoxine (Albon) can be given at the same time as any one of the other drugs.

Associated fecal parasite analysis:

A set of 3 fecal samples should be collected over a 5 day period for analysis by both zinc and sugar flotation methods prior to initiation of treatment. A second 3 sample set of fecal samples should be collected and analyzed ~7 days (to allow for analysis time) before the dogs leave the island.

After departure of dogs from the island, options for sterilization of dog housing areas for the purposes of reducing or eliminating parasite eggs are:

The top 6 inches of soil can be removed and buried below 18 inches of normal soil.

Recovery Plan for Four Subspecies of Island Fox

The area can be protected from foxes – because parasite eggs are very persistent in the environment, the area should be kept protected for 10 years.

The soil (or other surface) can be steamed with a steam pressure cleaner – this works well for hard surfaces like wood or concrete, and can be effective with soil, but depth of penetration of the steam is difficult to determine with certainty.

Hydrated lime or sodium borate may be mixed with the soil to kill parasites with desiccation and heat.

Concrete areas can be treated with bleach or ammonia.

Movement of Working Military Dogs on and off the Navy Islands (Nicolas & Clemente):

It is recognized by the fox health group that military operational considerations are sometimes at odds with previously recommended protocols for dogs entering the islands, especially requirements for quarantine periods. If quarantine periods are not possible, we recommend that all feasible measures should be taken so that there is no direct exposure of foxes to working dogs and their waste materials.

We also recommend that all military working dogs taken to the islands be current on their vaccines and currently on a monthly parasite preventative.

We also strongly recommend that no military working dogs come to the island within 2 months of any overseas deployment. This is to make it more likely that any infectious diseases acquired during a deployment will have been detected clinically or through testing prior to entry to the island.

G. APPENDIX 7: GUIDELINES FOR ESTABLISHING A MAINLAND POPULATION OF ISLAND FOXES WITH TASKS, RESPONSIBLE PARTIES, AND TIME FRAMES

This table of identified steps was modified from May 2005 Technical Analysis Response 3.6.

Task	Actions	Responsible Parties
Obtain necessary ESA permits and authorizations		
	Establish an ESA and CDFW permit held by the FWS for all mainland activities, including quarantine, holding, breeding, research and animal shipments	FWS, CDFW
	Obtain and maintain the appropriate permits as required by the Department of Fish and Wildlife and the Fish and Wildlife Service	AZA-SSP, FWS, CDFW
Develop necessary shipping, quarantine, and holding protocols and guidelines		
	Develop a standardized animal shipment protocol for off island transfers	AZA-SSP
	Approval of mainland facilities will be based on existing CDFW policies, procedures, and permitting processes.	AZA-SSP, FWS, CDFW
	Develop draft quarantine protocols	Completed
	Submit draft quarantine protocols to Health TEG for review	Completed
	Finalize Quarantine protocol	Randy Junge for Health TEG
	Present Quarantine protocol to Group	Health TEG
	Create standardized protocols for preventive medicine while on mainland	Health TEG
Develop AZA Species Survival Plan (SSP) to provide oversight of mainland population		
	Establish an MOU between AZA and the FWS	Completed
	Establish an AZA Island Fox SSP	AZA Canid TAG
	Name a SSP Coordinator, Studbook Keeper and necessary Advisor(s)	AZA Canid TAG
	Recognition by the FWS and CDFW of the AZA Island Fox SSP as the knowledgeable authority on mainland captive breeding for Island foxes	FWS, CDFW
	Decide SSP goals for mainland population	AZA-SSP, TAG, FWS, NPS, TNC, CDFW
	Recruit SSP institutions	AZA-SSP
	Develop SSP Master Plan	AZA-SSP

Recovery Plan for Four Subspecies of Island Fox

Task	Actions	Responsible Parties
Address financial oversight issues and concerns		
	Set up a FWS grant agreement with SSP	FWS
	Each holding institution takes financial responsibility for animals held	AZA-SSP, Each holding facility
	Liaise with Friends of the Island Fox, Inc. (FIFI) for development activities	AZA-SSP, FIFI, FWS
	ID private sources of funding	AZA-SSP, FIFI
	ID grant sources of funding	AZA-SSP, FIFI
	Set up gifted endowment	FIFI

H. APPENDIX 8: IDENTIFIED RESEARCH NEEDS USING A MAINLAND CAPTIVE ISLAND FOX POPULATION

15 July 2007

A. Ensure that husbandry and management practices in captive facilities enhance animal welfare and maximize breeding success.

- Develop methodologies to encourage maximum breeding success, together with methods for monitoring and record keeping.
- Determine the “Best Management Practices” for housing, managing and breeding captive Island foxes through testing of different housing and management protocols. Analyze research results on the social and reproductive behavior of wild island foxes to inform captive husbandry and management.
- Using an adaptive management paradigm, ensure that captive facilities conform to the best animal care standards known for comparable species and facilities. Strive to achieve captive breeding success rates for island foxes that are comparable with the most successful canid breeding program in the AZA’s Canid Taxon Advisory Group (TAG).

B. Conduct research using captive island fox populations to enhance their welfare and breeding success in captivity, and to help identify, eliminate and control threats to the recovery and sustainability of wild island fox populations.

- Analyze mechanisms of mate choice and mate compatibility, e.g. determine the differences in behavioral characteristics of successful and unsuccessful mated pairs and the underlying cause(s).
- Analyze husbandry and management factors contributing to reproductive success and failure, e.g. determine when in the reproductive cycle failure occurs; determine factors contributing to pup loss; determine the best pen design and management practices for captive island foxes; determine the best management practices for facilitating successful mate choice and producing successful breeding pairs.
- Conduct biomedical research on captive populations to help eliminate and/or control disease threats to the wild and captive island fox populations, e.g. develop and test vaccines using captive individuals to help develop protocols and practices for vaccinations of wild island foxes.
- Conduct and perfect semen collection, cryo-preservation and semen banking for a genetic bank.

C. Develop “Best Management Practices” for rearing island foxes for release into the wild and for their reintroduction.

- Identify and conduct research on the management and husbandry factors that influence post-release survival of island foxes.
- Develop the most cost-effective methods of housing and rearing island foxes to preserve their ability to survive in the wild. Develop objectives for evaluation of success of management protocols.
- Develop “Best Management Practices” for releasing island foxes into the wild and for post-release management and monitoring. Variables that need consideration include, in priority order, feeding regimes before and after release, post-release support (e.g. food supplementation and shelter), individual differences in island fox behavior, pre-release housing and adaptation to release sites, social groupings, and medical exams. Include for each variable how, what, how much, when, and for how long.
 1. Develop standardized protocols for pre-release preparation.
 2. Develop objectives for evaluation of reintroduction success.
 3. Develop and implement standardized techniques for post-release management and monitoring. Ensure compatibility between short-term monitoring of new releases and long-term monitoring protocols for the wild island fox populations.

I. APPENDIX 9: LIST OF TECHNICAL ANALYSIS REQUESTS (TARS)

Copies of these technical reports can be obtained by contacting the Ventura Fish and Wildlife Office at 805-644-1766 or 2493 Portola Road, Suite B; Ventura, California 93003.

Note that some of the information and/or recommendations provided in these analyses may no longer be up-to-date or consistent with current island fox recovery efforts. Some approaches to recovery have changed since they were developed and may change in the future as new information arises.

Technical Analysis 1.3

Use the PVA models and supporting data to determine the conditions in the wild populations that would trigger taking further foxes into captivity (e.g., during pig eradication on Santa Cruz, or if another disease outbreak occurred).

Technical Analysis 2.1

Development of population monitoring plans for free-ranging island foxes.

Technical Analysis 3.1

Determine the target captive population size for each subspecies, building on population viability analyses for wild population and demographic and genetic data on which these models are based.

Urgent Technical Analysis Related to Analyses 3.1 and 3.3

Determine whether, how, and where to release captive-bred foxes this fall and, if no releases, develop contingency plans that may include establishing mainland populations or expanding existing on-island populations.

Technical Analysis 3.4

Develop management and husbandry plans for each subspecies, taking into account studbook data, and results from research into best husbandry practices (pen size, social structure, mate choice, etc). The focus for research and management for each captive population will depend on the size and stability of that subspecies' wild and captive populations.

Technical Analysis 3.6

Assessment of the potential benefits and costs of long-term captive populations on the mainland and/or islands.

Sub-Analysis 3.6.1:

Recovery Plan for Four Subspecies of Island Fox

Identify and describe the potential benefits, costs, and major issues associated with the following strategies (or combinations of thereof) for maintaining captive populations of island foxes:

- a. using existing on-island facilities.
- b. expanding on-island facilities.
- c. using existing space in mainland facilities (e.g., zoos).
- d. constructing new mainland facilities for island foxes.

Sub-Analysis 3.6.2:

Identify to the extent possible the necessary steps and their logical progression for establishing and managing captive populations on the mainland.

Sub-Analysis 3.6.3:

If the establishment of mainland populations is determined to be both desirable and practical, identify weight criteria to be used to prioritize subspecies of the island foxes for representation in mainland populations.

Technical Analysis 4.1

Analyze efficacy of golden eagle control and capture methods utilized to date and recommend innovative program for removal methods, taking into account the most up-to-date information on the status of the wild fox populations.

Recovery Plan for Four Subspecies of Island Fox

Table 8-1: Date that the Island Fox Recovery Coordination Group (RCG) issued their recommendation for each TAR.

TAR 1.3	September 10, 2004
TAR 2.1	The RCG did not issue a recommendation associated with this TAR. The request was sent October 20, 2006.
TAR 3.1	See “Urgent Technical Analysis Related to Analyses 3.1 and 3.3.”
Urgent Technical Analysis Related to Analyses 3.1 and 3.3	October 4, 2004
TAR 3.4	April 12, 2005
TAR 3.6	February 9, 2006
TAR 4.1	January 7, 2004

Recovery Plan for Four Subspecies of Island Fox

J. APPENDIX 10: GLOSSARY OF TERMS IN THE RECOVERY PLAN FOR FOUR SUBSPECIES OF ISLAND FOX

Age class	Foxes are aged according to tooth eruption and wear patterns on the first upper molar (Wood 1958; Roemer 1999) and are assigned to discrete age classes: pups (Age Class 0), young adults (Age Class 1: 7 months to 2 years), adults (Age Class 2: 2 to 3 years), mature adults (Age Class 3: 3 to 4 years) and old adults (Age Class 4: greater than 4 years) (Roemer 1999).
Allee effects	Allee effects occur when population growth rate decreases with declining density. Allee effects are expected to occur at very small population sizes and may arise due to mate scarcity, inbreeding, or disruption in social behavior.
Allele	An allele is one member of a pair or series of genes that occupy a specific position on a specific chromosome.
Allelic diversity	Allelic diversity is the average number of alleles per locus.
Allozyme	Allozymes are variable forms of the same enzyme. They differ because they originate from different genetic sequences. Studying allozyme variation in a population is an indirect measure of the genetic variation of the population. In this type of study, whole proteins are analyzed instead of the genes they originated from. If there is no genetic variation in a population (monomorphic = “one form”), the entire population uses just one form of the gene. If the population shows polymorphisms (“many forms”) of allozymes, this can be used to describe how much variation is present.
Anthelmintics	Anthelmintics are agents used to treat parasitic worms in animals.
Bottleneck	A population undergoes a bottleneck when a combination of environmental conditions occurs that causes a serious reduction in the size of a population. A population that has undergone a bottleneck often has reduced genetic diversity.

Recovery Plan for Four Subspecies of Island Fox

Caliciviruses	Members of the Caliciviridae family of viruses. Calicivirus infections commonly cause acute gastroenteritis, which is the inflammation of the stomach and intestines. Symptoms can include vomiting and diarrhea.
Contraindication	A contraindication is a specific situation in which a drug, procedure, or surgery should not be used, because it may be harmful to the patient.
Degloving	A type of injury where an extensive section of skin is completely torn off the underlying tissue, severing its blood supply.
Demographic	A characteristic used to describe some measurable aspect of a population, such as growth rate, age structure, birth rate, and gross reproduction rate.
Diurnal	An animal that is diurnal is active during the daytime and rests during the night, as opposed to an animal that is nocturnal, or mostly active during the nighttime.
DNA restriction fragments	A DNA restriction fragment is a DNA fragment resulting from the cutting of a DNA strand by a restriction enzyme by a process called restriction. Each restriction enzyme is highly specific, recognizing a particular short DNA sequence and cutting both DNA strands at specific points within this site. Restriction fragments can be analyzed using techniques such as gel electrophoresis or used in recombinant DNA technology.
Docile canid	A docile canid refers to any of the various widely distributed carnivorous or omnivorous mammals of the family Canidae, which includes the foxes, wolves, dogs, jackals, and coyotes, that are easily managed or handled.
Endoparasite	An endoparasite is a parasite that inhabits the internal organs or tissues of an animal or plant
Effective population size	The effective population size refers to the average number of individuals in a population that actually contribute genes to succeeding generations by breeding. This number is generally lower than the observed population size.

Recovery Plan for Four Subspecies of Island Fox

Evisceration	To remove the entrails of; disembowel.
Founder	A founder is an individual that established a population. The founders referred to in this recovery plan are the individuals that were captured from the wild and brought into captivity, establishing the captive breeding programs.
Hacking	Hacking is a technique used in the captive release of birds of prey where shelter and food is provided for a bird prior to fledging and is continued to be provided until the fledging becomes independent.
Heterozygosity	Heterozygosity is the proportion of individuals heterozygous at all loci divided by the number of loci.
Heterozygous	Heterozygous refers to an individual that possesses two different forms of a particular gene, one inherited from each parent.
Hyperpredation	Analogous to apparent competition (Holt 1977), hyperpredation occurs when a prey species that can sustain high predation rates subsidizes the extinction of another prey species by acting as an alternate food resource for a shared predator (Courchamp et al. 1999).
Hypervariable DNA	Usually used to describe the sequences of DNA that do not have a function in the organism, a.k.a. “junk” DNA. These sequences are especially useful for determining differences within the population, because these regions of DNA are not critical and can vary greatly from one individual to another with no consequence.
Inbreeding depression	Inbreeding depression is an increased expression of deleterious alleles in individuals, resulting in an overall decline in the vigor of a population, due to mating among relatives.
Intraspecific aggression	Intraspecific aggression refers to aggression among members of a single species.
Isocline	An isocline is a series of lines with the same slope. For our purposes, an isocline graph provides a way

Recovery Plan for Four Subspecies of Island Fox

to visually identify where an island fox population stands in regards to current on-island parameters.

Leptospira

Very slender aerobic spirochetes; free-living or parasitic in mammals.

Loci (plural of locus)

In genetics, the locus is a specific place on a chromosome where a gene is located.

Major Histocompatibility Complex (MHC)

Genetic loci that encode for three classes of transmembrane (cell) proteins. These proteins induce the organism's immune response. This region is especially important in genetic studies since these types of genes are found in all vertebrates and are numerous, making these regions a rich source of information regarding genetic evolutionary lineage. There can be hundreds of alleles of each MHC locus.

Metastasize

The spreading of cancer cells from their original site to other parts of the body.

Microsatellite DNA

Pieces of the same small DNA sequence which are repeated, often in non-coding genetic regions of the chromosome ("junk" DNA that does not contain any genes).

For example, if a small nucleotide sequence normally repeats within a species' genome, it may repeat a different number of times in different individuals. The number of repeats is easily detected and serves as a basis for measuring the genetic variability within the population.

Minisatellite DNA

Similar to microsatellite DNA, minisatellites are longer DNA sequences (1000 to 5000 bases long) of 20-50 repeats.

Mitochondrial DNA

The DNA within an organism's cells that is located inside the mitochondria, not inside the nucleus. Mitochondrial DNA is maternally inherited in most species and is used widely to assess taxonomic relationships and differences among populations and species. Mitochondrial DNA analysis (mtDNA) can be used to examine older biological samples that lack cellular material with a nucleus (nucleated), such as hair, bones, and teeth.

Recovery Plan for Four Subspecies of Island Fox

Morbillivirus	Morbillivirus is a genus of viruses in the family of Paramyxoviridae that includes the causative agents of measles, canine distemper virus, phocine distemper, and rinderpest.
Necropsy	A necropsy is an examination and dissection of a dead body to determine cause of death or the changes produced by disease.
Oocyst	An oocyst is the thick-walled spore phase of certain protists (sporozoans), such as <i>Cryptosporidium</i> and <i>Toxoplasma</i> .
Otitis	Otitis is a bacterial infection of the ear.
Pelage	Pelage is the wool, fur, or hair coat of a mammal.
Phylogeny	Phylogeny is the evolutionary development and history of a species or higher taxonomic grouping of organisms.
Polymorphism	Polymorphism is the existence of two or more forms of individuals within the same animal species (independent of sex differences).
Prophylactic management	Implementation of protective measures to prevent disease or extinction.
Quasi-extinction	Quasi-extinction is a drop in numbers of individuals to some very low level at which the population is expected to be critically imperiled. At quasi-extinction, population dynamics are expected to be immediately and adversely affected by factors such as Allee effects, inbreeding depression, and demographic randomness. At this point, management options are severely constrained.
Serological	A serological test or survey pertains to the measurement and characterization of antibodies, antigens, and other immunological substances in body fluids (serum), usually blood.
Serovar	A group of closely related microorganisms distinguished by a characteristic set of antigens.
Stochastic events	Chance or random events.

Recovery Plan for Four Subspecies of Island Fox

Sympatric	Sympatric refers to populations or closely related species that occupy the same or overlapping geographic areas without interbreeding.
Synergistic	Different actions combined or correlated, working together.
Toxoplasmosis	An intracellular tissue infection of a parasite from the genus <i>Toxoplasma</i> , particularly in mammals and birds.
Vital rates	Rates of those components; such as birth, marriage, fertility, and death; which indicate the nature and possible changes in a population. Even when population numbers are stable, there may be changes in the vital rates.

References for Glossary

- Courchamp, F., M. Langlais, and G. Sugihara. 1999. Control of rabbits to protect island birds from cat predation. *Biological Conservation* 89:219-225.
- Holt, R.D. 1977. Predation, apparent competition and the structure of prey communities. *Theoretical. Population Biology* 12: 197-229.
- Roemer, G.W. 1999. The ecology and conservation of the island fox (*Urocyon littoralis*). Ph.D. dissertation, University of California, Los Angeles. 229 pp.
- Wood, J. E. 1958. Age structure and productivity of a gray fox population. *Journal of Mammology* 39:74-86.

**K. APPENDIX 11: ACRONYMS AND ABBREVIATIONS IN THE
RECOVERY PLAN FOR FOUR SUBSPECIES OF ISLAND FOX**

AZA	Association of Zoos and Aquariums
CBI	Conservation Biology Institute
CDFG	California Department of Fish and Game
CDFW	California Department of Fish and Wildlife (formally called California Department of Fish and Game)
CDV	Canine distemper virus
CIC	Santa Catalina Island Conservancy
CINP	Channel Islands National Park
DDT	Dichlorodiphenyltrichloroethane
DNA	Deoxyribonucleic acid
ENSO	El Niño Southern Oscillations
ESA	Endangered Species Act of 1973, as amended
FIV	Feline immunodeficiency virus
FelV	Feline leukemia virus
FWS	U.S. Fish and Wildlife Service
GIS	Geographic Information System
IRT	Island Fox Integrated Recovery Team
IUCN	World Conservation Union (formerly called International Union for Conservation of Nature and Natural Resources)
IA	Interagency Agreement
IWS	Institute for Wildlife Studies

Recovery Plan for Four Subspecies of Island Fox

Km ²	Square kilometer
MHC	Major Histocompatibility Complex
MOA	Memorandum of Agreement
NEPA	National Environmental Policy Act
NPS	National Park Service
PMTC	Pacific Missile Test Center
PVA	Population Viability Analysis
RCG	Island Fox Recovery Coordination Group
SSC	Species Survival Commission
SSP	Species Survival Plan
TAG	Taxon Advisory Group
TAR	Technical Analysis Request
TBD	To be determined
TEG	Technical Expertise Group
TNC	The Nature Conservancy
USGS-BRD	U.S. Geological Survey - Biological Resource Discipline
UNIV	University or academic researchers

L. APPENDIX 12: SUMMARY OF COMMENTS

Summary of Comments

On September 14, 2012, we published a Notice of Availability in the Federal Register of the Draft Recovery Plan for Four Subspecies of Island Fox (Draft Plan) for a 60-day comment period ending on November 13, 2012, for Federal agencies, State and local governments, and members of the public (77 FR 56858). In accordance with our peer review policy published on July 1, 1994 (59 FR 34270), we solicited expert opinions from four knowledgeable individuals with scientific expertise that included familiarity with canids, carnivores, the geographic region in which the species occurs, and conservation biology principles relevant to the species. We received responses from two of the four peer reviewers.

This section provides a summary of general demographic information including the total number of letters from various affiliations and also provides a summary of major comments, as appropriate. All letters of comment on the draft plan are kept on file in the Ventura Fish and Wildlife Office at 2493 Portola Road, Suite B, Ventura, CA 93003.

The following is a breakdown of the number of letters received, by affiliation:

Federal agencies	2
State agencies	1
Environmental/conservation organizations	1
Academia/professional	1
Individual citizens	1

A total of six letters were received, each containing varying numbers of comments. Some of the comments were duplicated between letters. Most letters provided new information or suggestions for clarity. Where appropriate, we incorporated new information into the final version of the recovery plan. Some comments dealt with matters of opinion, which did not provide information relevant to the recovery of island fox and did not result in changes to the draft plan. A few of the comments concurred with parts of the draft plan and while these review comments were helpful, they generally did not result in changes. We did not receive any comments that we considered controversial or significant in the sense of making a difference in the fundamental way that recovery of the island fox is being approached. Information and comments not incorporated into the final version of the recovery plan were considered, noted, and are on file with the entire package of agency and public comments. Major comments that were not incorporated or that require additional clarification in addition to their incorporation are addressed below.

Summary of Comments and Our Responses

Comment: One commenter strongly disagreed with the statement on page 84, section 4.4: “Prevent disturbance of island ecosystems’ native structure and function to the extent practicable,” stating that, “island ecosystems are inherently unstable; that is the natural state of the system. Attempting to impose stability on that system may be legally necessary, but it is not ecologically appropriate and will have unintended consequences.”

Response: We recognize the issues associated with trying to maintain an ecosystem at pre-human condition and the difficulty in restoring these affected ecosystems and have modified the text accordingly. We have changed, “Prevent disturbance of island ecosystems’ native structure and function to the extent practicable” to “Prevent excessive human-caused impacts to island ecosystems’ native structure and function to the extent practicable”. We believe the decades of heavy disturbance by human associated activities, including grazing, have altered the landscape over time to such a degree that in order to maintain some stability, actions must be taken to prevent further human induced ecosystem change while stabilizing and restoring, to the extent practicable, past human induced disturbances.

Comment: One commenter expressed concern that Santa Rosa Island is grouped with the other northern Channel Islands that experienced ‘hypermortality’ as a result of golden eagle predation, yet the information provided in the document indicates that while other islands have rebounded dramatically, Santa Rosa has not, and over 4 months in late 2006, 10 foxes (12% of the total population at that time) died of something other than eagle predation indicating there may be other sources of mortality on Santa Rosa Island that need to be considered.

Response: In those cases in which the cause could be determined, the vast majority was golden eagle predation. Predation by golden eagles accounted for 33 of 77 island fox mortalities from 2003-2013 on Santa Rosa Island. The cause of mortality was not determined in 29 cases due to advanced decay of the carcass. Other mortality causes included entrapment in drain pipes (3), intestinal intussusception (collapse of one section of the intestine into another; 1 case), cholecystitis (inflammation of the gall bladder caused by gallstones, 2 cases), leptospirosis (2), and emaciation (3, following wounds).

Comment: One commenter expressed concern that the natural source and rate of mortality is nowhere to be found in the document, and that various values are periodically described as high or low, but there is no baseline to compare these numbers.

Response: There are very little data from fox populations that aren't influenced by catastrophic mortality factors, except for the recovering populations without influence by predation/disease. Survival has been very high in the recovering populations - 90% and above. The only pre-decline "background" survival rates we are aware of are those from San Miguel Island in 1993-1994, and those were close to 100% for adults and above 70% for pups (Coonan et al. 2005). On Santa

Recovery Plan for Four Subspecies of Island Fox

Cruz Island, 1993-1994, the survival rate was also close to 100% (Roemer et al. 2001).

Comment: Two commenters suggested including bald eagle reintroductions as part of the recovery program.

Response: We believe bald eagle reintroduction falls outside the scope of this document. The reintroduction of bald eagles to the Channel Islands began in 1980 on Catalina Island and in 2002 for the northern Channel Islands. Reintroductions to the northern Channel Islands have been a major focus of the Montrose Settlements Restoration Program. While bald eagle reintroductions have benefited the recovery of island foxes on the northern Channel Islands, these efforts started in 2002, prior to the Federal listing which occurred in 2004. Therefore, while we include bald eagle reintroductions as part of the ongoing recovery efforts being conducted for island foxes, we did not include the reintroduction of bald eagles as a recovery action within the recovery plan since this program was initiated prior to Federal listing.

Comment: One commenter expressed concern about the lack of monitoring within the Long-Term Conservation section given that island systems are always under threat by their isolated nature.

Response: We believe that we have appropriately addressed this concern in the recovery plan. As part of the Long-Term Conservation section, we identify the establishment of Conservation Agreements between the land manager(s) and the FWS to address long-term conservation needs, one of which is, “The land manager’s strategy and commitment to continue monitoring island fox subspecies such that any substantial population decline is detected in a timely manner.” We believe this is appropriate given that, even with successful mitigation of current threats and the recovery of island fox subspecies to viable population levels, the intrinsically small population sizes of the subspecies and their insular vulnerabilities subject the different subspecies to the continued threat of catastrophic decline from any number of causes.

Comment: One commenter asked, “What evidence is there for bald eagle effects on golden eagles (preventing golden eagles from nesting)?”

Response: We are not aware of any direct evidence or studies that address this comment. However, we believe the extirpation of bald eagles (*Haliaeetus leucocephalus*) from the Channel Islands as a result of dichlorodiphenyltrichloroethane (DDT) may have facilitated golden eagle colonization. Bald eagles historically bred on the islands and aggression by breeding bald eagles may have discouraged foraging golden eagles from establishing residence.

Comment: Two commenters suggested that we should mention in the disease section that the threat of rabies is so great that all fox subspecies are vaccinated against it.

Recovery Plan for Four Subspecies of Island Fox

Response: We have added the following text and believe it is worth mentioning here separately. “Although rabies has never been found in island wildlife, if it were to occur, there would be no way to effectively mitigate an outbreak. Hence the only defense is a vaccination program.”

Comment: One commenter expressed disagreement with the long-term conservation actions requiring maintenance of a vaccinated subpopulation because these types of actions run counter to the natural instability of island populations.

Response: Although rabies has never been found in island wildlife, if it were to occur, there would be no way to effectively mitigate an outbreak. Hence the only defenses are a vaccination program and a subset of each fox subspecies are vaccinated against it. Furthermore, reducing the threat of disease will require 1) preventing introduction of new pathogens or novel strains of existing pathogens to the Channel Islands by eliminating visitation by mainland animals, and 2) the continuation of the vaccination programs already established on each of the northern Channel Islands and Santa Catalina. Although mainland animal visitation is prohibited on the northern Channel Islands, evidence exists that dogs are frequently brought to the islands. Therefore, continued proactive vaccination of a core population of island foxes remains the only viable alternative to protect these populations from CDV and rabies. Additionally, future vaccinations for any delisted subspecies will potentially help prevent the need for relisting under the Endangered Species Act.

Comment: One commenter suggested the statement “predation by golden eagles remains a threat to the long-term recovery of wild island fox populations” should be reconsidered in the light of management actions and the resulting lack of breeding by golden eagles on the islands and greatly reduced predation by golden eagles on foxes

Response: We agree. Timely threat detection and assessment of appropriate management actions is critical to maintain responsible management of island foxes, which exist in a landscape that has been fundamentally altered by human impacts. We have modified the document to reflect this by inserting “potential.” The sentence now reads, “All known golden eagles have been removed from the northern Channel Islands, yet predation by golden eagles remains a potential threat to the long-term recovery of wild island fox populations, including in the southern Channel Islands.” We note, even one pair of nesting golden eagles appears to put significant pressure on island fox populations, whereas island fox populations can sustain some predation by transient eagles, as occurred in 2007-2013. Thus, golden eagle monitoring and the removal of nesting birds need to continue, and management agencies need to be prepared to respond if and when golden eagles nest on any of the Channel Islands.

Comment: One commenter suggested we revise Recovery Criteria 1 that states, “risk level is sustained for at least 5 years, during which time the population trend is not declining” to include text identifying what defines a declining trend.

Recovery Plan for Four Subspecies of Island Fox

Response: We agree and have included the following text to define a declining trend. For the purpose of this recovery plan, “A declining trend is defined as the 3-year risk-level being greater in year 5 than year 1.”

Comment: One commenter suggested we modify Recovery Objective 2 to reflect the reduced likelihood of predation by golden eagles.

Response: We agree and have modified the text for Recovery Objective 2, from “Land managers are able to respond in a timely fashion to potential and ongoing predation by golden eagles, to potential or incipient disease outbreaks, and to other identified threats” to “Land managers are able to respond in a timely fashion to predation by nesting golden eagles or significant predation rates by transient golden eagles, to potential or incipient disease outbreaks, and to other identified threats using the best available technology.”

Comment: One commenter suggested that it is counterproductive to develop a formal golden eagle strategy beyond the strategies that have already been successfully developed and implemented.

Response: We believe it is still necessary to have a strategy in place and have recommended the existing strategy be included as part of the epidemic response plan. We believe the guidelines set forth for detecting disease-related mortality events is applicable to detecting mortality of island foxes by golden eagles. Guidance has been provided on the monitoring intensity needed to detect threats imposed by golden eagles and disease before those threats unduly impact island fox populations.

Comment: One commenter suggested the following text, as described in the Recovery Actions Section, “Manage captive populations of island foxes for recovery” implies that the recovery action is ongoing.

Response: To address this comment, we have inserted the following text: “If required, conduct captive breeding and reintroduction of island foxes to increase population size (captive breeding and reintroduction were conducted from 1999-2008 and ceased due to the success of reintroductions and rapid growth of recovering populations).”

Comment: One commenter disagreed with text that states, “Quick and accurate detection of possible future declines is paramount to maintaining viable fox populations,” stating that “islands did manage to retain viable populations through numerous historic bottlenecks, despite documented reductions to <10 individuals in at least one case.”

Response: Based on the complexities associated with island ecosystems and the inherent risks they face, timely threat detection and assessment of appropriate management actions is critical to maintain responsible management of island foxes, which exist in a landscape that has been fundamentally altered by human impacts. Guidance has been provided for the monitoring intensity needed to detect threats imposed by golden eagles and disease before those threats unduly impact island fox populations.

Recovery Plan for Four Subspecies of Island Fox

Comment: One commenter expressed concern that the species is being relegated to permanent maintenance, given the fact that new threats to small, island populations will always arise and that once these factors are mitigated to the best of our ability, island fox populations should be allowed to pursue their natural trajectory, even if that means instability and potential extinction on individual islands.

Response: We recognize the issues associated with trying to maintain an ecosystem at pre-human conditions and the difficulty in restoring these affected ecosystems and have modified the text accordingly. We have changed, “Prevent disturbance of island ecosystems’ native structure and function to the extent practicable” to “Prevent excessive human-caused impacts to island ecosystems’ native structure and function to the extent practicable.” We believe the decades of heavy disturbance by human associated activities, including grazing, have altered the landscape over time to such a degree that in order to maintain some stability, actions must be taken to prevent future human induced ecosystem change while stabilizing and restoring, to the extent practicable, past human induced disturbances.

Comment: One commenter expressed concern that golden eagles being observed but not actively preying upon island foxes will be allowed to remain on the island.

Response: We have reflected this change in the document which now reads, “Response tactics (including the use of helicopters and net-guns) to capture nesting golden eagles and transient golden eagles responsible for significant island fox predation, per the golden eagle response strategy.”

Comment: One commenter expressed concern about vaccination on Santa Cruz and Santa Catalina Islands being focused on one or two localized areas.

Response: We agree and have updated the text to reflect this change. On islands where wild fox populations number fewer than 100 individuals, all island foxes should be vaccinated. On Santa Cruz Island, vaccination should be focused in one or two localized areas, but specifically in primary access corridors, such as Prisoner’s Harbor up canyon to the central valley. On Santa Catalina Island, vaccination efforts should be concentrated around the city of Avalon (where disease introduction is most likely to occur) and around the isthmus (where infection could potentially pass between the eastern and western subpopulations). However, because of the many points of access to Santa Catalina Island, island-wide vaccination is preferred.

Comment: One commenter suggested that we not exclude animals killed by trauma from being necropsied, stating that “disease may predispose an animal to death by other sources, including roadkill.”

Response: We agree. We have removed the language “from causes other than trauma.”

Recovery Plan for Four Subspecies of Island Fox

Comment: One commenter recommended that we note that it is not feasible to maintain more than one subspecies as a genetically diverse, sustainable population within mainland Association of Zoos and Aquariums accredited zoos due to space limitations.

Response: We believe this is already identified with the existing text, which states, “may benefit from redundant, genetically diverse, and sustainable mainland populations of one or two subspecies.”

Comment: One commenter suggested sections should be updated to reflect the fact that captive breeding and reintroduction has ceased.

Response: We agree. We have included text, where appropriate, to reflect this update. An example of text inserted is as follows: “On-island captive breeding and reintroduction were conducted from 1999-2008 and ceased due to the success of reintroductions and the rapid growth of recovering populations.”

Comment: One commenter was unable to find in the document an outline for future serology testing work. The commenter assumed CDV, CPV, and CAV will be screened for each year, but asked if any others would be screened, such as Leptospirosis and toxoplasma?

Response: Appendix 4 “Guidelines for Vaccination and Sampling of Captive and Wild Populations” of the Recovery Plan states, “Conduct a serosurvey for antibodies to CDV, canine parvovirus (CPV), canine adenovirus (CAV), and *Toxoplasma* as often as yearly, but no less often than every 5 years. At this time, while leptospirosis is not included in the serology list, it’s a pathogen that is monitored for using the post-mortem (necropsy) surveillance.

Comment: One commenter noted there were some difficulties in comparing seroprevalence across years because different labs/tests were used and inquired whether there is a particular lab that will be used each year or at least a standardized testing protocol.

Response: Difficulties in comparing seroprevalence across years may be explained in part by differences in test sensitivities in the labs used for these surveys between 1988 and 2006. Since the 2006 survey, the Island Fox Health Advisory Group recommended Cornell University serve as the single lab to be used for island fox serology. The lab at Cornell University has been used since that date so that testing is standardized and cross-island comparisons can be made.

Comment: One commenter suggested that the subspecies determination for a mainland captive island fox population be considered as soon as possible, prior to further investment being made in (and valuable spaces filled by) San Clemente Island foxes.

Response: We agree. According to the “Assessment of the potential benefits and costs of long-term captive populations on the mainland and/or islands” suggests that the Santa Cruz Island fox population would be the best choice, because this subspecies has the most genetic diversity and the island population is recovering rapidly.

Comment: One commenter suggested we add “Only injured, unreleasable foxes or abandoned pups that are incapable of surviving in the wild will be considered for transfer to zoos.”

Response: We agree that at one time this was the responsible way for moving forward as a result of the incredibly low numbers of individuals present on each island; however, given that space and resources are limited to establish redundant populations for each of the four endangered subspecies of island fox, it is unrealistic to expect to have a redundant population of each subspecies.

According to the “Assessment of the potential benefits and costs of long-term captive populations on the mainland and/or islands,” the Santa Cruz Island fox population would be the best choice, because this subspecies has the most genetic diversity and the island fox population is recovering rapidly. Therefore, based on the long-term conservation strategy that identifies actions that would further the conservation of the island fox, we do not believe it is in the best interest at this time to remove island foxes from islands other than Santa Cruz for the genetically diverse mainland captive population. We also believe that in order to establish a genetically diverse mainland captive population, we cannot rely only on those foxes that are injured, unreleasable, or abandoned pups that are incapable of surviving in the wild; healthy foxes that will best represent and meet the needs for fulfilling this activity will also be necessary.

Comment: One commenter agreed that we should establish, expand and continue island fox education and outreach programs, but we should add an assessment requirement so the effectiveness of education programs can be ensured.

Additionally, the statement that “All zoos that house island fox populations, particularly those in southern California should include a robust education program” leads to difficulty in measuring “robust.”

Response: We agree and have modified the text accordingly. All zoos that house island fox populations, particularly those in southern California, should include an education program with an assessment requirement of the education programs effectiveness. An example of an effective program is defined as a 10 percent increase in public awareness, such that 10 percent of visitors learn that island foxes are unique to the Channel Islands and the species’ could be at risk of contracting mainland diseases from mainland animals.

Literature Cited

Coonan, T.J., C.A. Schwemm, G.W. Roemer, D.K. Garcelon, and L. Munson.

2005. Decline of an island fox subspecies to near extinction.

Southwestern Naturalist 50(1):32-41.

Roemer, G.W., T.J. Coonan, D.K. Garcelon, J. Bascompte, and L. Laughrin.

2001. Feral pigs facilitate hyperpredation by golden eagles and indirectly cause the decline of the island fox. *Animal Conservation* 4:307-318.